

2023-1169

UNITED STATES COURT OF APPEALS FOR THE FEDERAL CIRCUIT

AMARIN PHARMA, INC., AMARIN PHARMACEUTICALS IRELAND LIMITED,
MOCHIDA PHARMACEUTICAL CO., LTD.,

Plaintiffs-Appellants,

v.

HIKMA PHARMACEUTICALS USA INC., HIKMA PHARMACEUTICALS PLC,

Defendants-Appellees,

HEALTH NET LLC,

Defendant

Appeal from the United States District Court for the District of Delaware
Case No. 1:20-cv-01630-RGA-JLH, Judge Richard G. Andrews

APPELLANTS' OPENING BRIEF

Nathan K. Kelley
Nathanael D. Andrews
PERKINS COIE LLP
700 13th Street NW, Suite 800
Washington, D.C. 20005
Phone: (202) 654-3343
E-mail: NKelley@perkinscoie.com

Mark T. Smith
PERKINS COIE LLP
110 N. Wacker Drive, Suite 3400
Chicago, IL 60606
Phone: (312) 324-8474
E-mail: MarkSmith@perkinscoie.com

*Counsel for Amarin Pharma, Inc., Amarin Pharmaceuticals Ireland Ltd., and
Mochida Pharmaceutical Co., Ltd.*

March 21, 2023

Claim Language

'537 patent

1. A method of reducing occurrence of a cardiovascular event in a hypercholesterolemia patient consisting of:

identifying a patient having triglycerides (TG) of at least 150 mg/DL and HDL-C of less than 40 mg/dL in a blood sample taken from the patient as a risk factor of a cardiovascular event, wherein the patient has not previously had a cardiovascular event, and administering ethyl icosapentate in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor,

wherein said 3-hydroxyl-3-methylglutaryl coenzyme A reductase inhibitor is administered to the patient at least one of before, during and after administering the ethyl icosapentate; and

wherein the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor is selected from the group consisting of pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, rosuvastatin, and salts thereof, and

wherein daily dose of the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor are 5 to 60 mg for pravastatin, 2.5 to 60 mg for simvastatin, 10 to 180 mg for fluvastatin sodium, 5 to 120 mg for atorvastatin calcium hydrate, 0.5 to 12 mg for pitavastatin calcium, 1.25 to 60 mg for rosuvastatin calcium, 5 to 160 mg for lovastatin, and 0.075 to 0.9 mg for cerivastatin sodium.

'861 patent

1. A method of reducing risk of cardiovascular death in a subject with established cardiovascular disease, the method comprising administering to said subject about 4 g of ethyl icosapentate per day for a period effective to reduce risk of cardiovascular death in the subject.
2. The method of claim 1, wherein the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL and a fasting baseline LDL-C level of about 40 mg/dL to about 100 mg/dL.

FORM 9. Certificate of Interest

Form 9 (p. 1)
July 2020

**UNITED STATES COURT OF APPEALS
FOR THE FEDERAL CIRCUIT**

CERTIFICATE OF INTEREST

Case Number 23-1169

Short Case Caption Amarin Pharma, Inc. v. Hikma Pharmaceuticals USA Inc.

Filing Party/Entity Amarin Pharma, Inc.; Amarin Pharmaceuticals Ireland; and
Mochida Pharmaceuticals Co., Ltd.

Instructions: Complete each section of the form. In answering items 2 and 3, be specific as to which represented entities the answers apply; lack of specificity may result in non-compliance. **Please enter only one item per box; attach additional pages as needed and check the relevant box.** Counsel must immediately file an amended Certificate of Interest if information changes. Fed. Cir. R. 47.4(b).

I certify the following information and any attached sheets are accurate and complete to the best of my knowledge.

Date: 3/21/2023

Signature: /s/Nathan K. Kelley

Name: Nathan K. Kelley

FORM 9. Certificate of Interest

Form 9 (p. 2)
July 2020

1. Represented Entities. Fed. Cir. R. 47.4(a)(1).	2. Real Party in Interest. Fed. Cir. R. 47.4(a)(2).	3. Parent Corporations and Stockholders. Fed. Cir. R. 47.4(a)(3).
Provide the full names of all entities represented by undersigned counsel in this case.	Provide the full names of all real parties in interest for the entities. Do not list the real parties if they are the same as the entities. <input checked="" type="checkbox"/> None/Not Applicable	Provide the full names of all parent corporations for the entities and all publicly held companies that own 10% or more stock in the entities. <input type="checkbox"/> None/Not Applicable
Amarin Pharma, Inc.		Amarin Corporation plc
Amarin Pharmaceuticals Ireland Limited		Amarin Corporation plc
Mochida Pharmaceutical Co., Ltd.		N/A

☐ Additional pages attached

FORM 9. Certificate of Interest

Form 9 (p. 3)
July 2020

4. Legal Representatives. List all law firms, partners, and associates that (a) appeared for the entities in the originating court or agency or (b) are expected to appear in this court for the entities. Do not include those who have already entered an appearance in this court. Fed. Cir. R. 47.4(a)(4).

☐ None/Not Applicable

☐ Additional pages attached

Fish & Richardson, P.C.	Jeremy Douglas Anderson	Robert M. Oakes
Deanna J. Reichel	Elizabeth M. Flanagan	Jonathan E. Singer
Michael J. Kane		

5. Related Cases. Provide the case titles and numbers of any case known to be pending in this court or any other court or agency that will directly affect or be directly affected by this court's decision in the pending appeal. Do not include the originating case number(s) for this case. Fed. Cir. R. 47.4(a)(5). See also Fed. Cir. R. 47.5(b).

☒ None/Not Applicable

☐ Additional pages attached

6. Organizational Victims and Bankruptcy Cases. Provide any information required under Fed. R. App. P. 26.1(b) (organizational victims in criminal cases) and 26.1(c) (bankruptcy case debtors and trustees). Fed. Cir. R. 47.4(a)(6).

☒ None/Not Applicable

☐ Additional pages attached

TABLE OF CONTENTS

Table of Authorities	iv
Table of Abbreviations and Conventions	vii
Related Cases	viii
Introduction	1
Jurisdiction	2
Statement of Issues.....	3
Statement of the Case.....	4
A. Vascepa® and its demonstrated efficacy for two different treatments.....	4
1. Severe hypertriglyceridemia, pancreatitis, and its treatment	4
2. Amarin initially patented the use of Vascepa® to treat severe HTG	5
3. Amarin later patented the use of Vascepa® to reduce cardiovascular risk in patients with non-severe HTG or existing cardiovascular disease.....	6
4. Mochida’s prior foundational work and patent	10
B. Hikma launched generic Vascepa®, describing it on its website, in press releases, and with a supposedly “skinny” label	12
1. Hikma submitted a Section viii statement to obtain FDA approval to sell generic Vascepa® solely to treat severe hypertriglyceridemia.....	12
2. Hikma’s website associated its generic version of Vascepa® with a method of use beyond its approved use.....	14
3. Hikma issued press releases, encouraging readers to prescribe generic Vascepa® for more than treating patients with severe hypertriglyceridemia.....	15

4.	Hikma’s “skinny” label teaches the claimed limitations and omits the prior limitation of use excluding the cardiovascular risk indication	17
C.	The district court proceeding.....	19
1.	Amarin sued Hikma for inducing infringement	19
2.	Hikma moved to dismiss under Rule 12(b)(6)	20
3.	The magistrate judge recommended denying Hikma’s motion to dismiss	21
4.	The district court granted Hikma’s motion	22
	Summary of Argument.....	25
	Argument.....	26
I.	Standard of review	26
II.	Amarin satisfied the pleading standard by pleading plausible infringement by Hikma	26
A.	Surviving a motion to dismiss requires pleading facts sufficient to state a plausible claim for relief	27
B.	Inducement requires showing direct infringement and actions taken with the intent to cause infringing conduct	29
C.	Taking the allegations together, Amarin’s claim against Hikma is plausible, and discovery is likely to further support Amarin’s case	30
III.	The district court erred by weighing the pled facts piecemeal against the plausibility pleading standard	35
IV.	The district court erred by resolving the key factual dispute of what Hikma’s conduct communicated to the market.....	39
V.	The district court erred by analogizing the wrong cases.....	41
A.	This case involves evidence showing inducement in addition to the label—it is not a “label only” case	42
B.	Previous cases consistently reached a later stage than a motion to dismiss because factual issues are central to inducement	48

C. This case involves a patented use that is broader than the off-patent use—unlike in <i>Grunenthal</i>	51
Conclusion	53
Addenda	
1. Memorandum Opinion, Order, and Final Judgment (Appx1-15)	
2. Magistrate’s Report and Recommendation (Appx1413-1430)	
3. U.S. Patent No. 9,700,537 (Appx36-47)	
4. U.S. Patent No. 10,568,861 (Appx77-129)	
Certificate of Compliance	

TABLE OF AUTHORITIES

Cases	Pages
<i>Amarin Pharma, Inc. v. Hikma Pharmaceuticals USA</i> , Case No. 20-1723 (Fed. Cir.), ECF No. 78	5
<i>Ashcroft v. Iqbal</i> , 556 U.S. 662 (2009).....	27, 28, 35
<i>AstraZeneca Pharms. LP v. Apotex Corp.</i> , 669 F.3d 1370 (Fed. Cir. 2012)	50
<i>AstraZeneca LP v. Apotex, Inc.</i> , 633 F.3d 1042 (Fed. Cir. 2010)	29, 44, 47, 49
<i>Ballentine v. United States</i> , 486 F.3d 806 (3d Cir. 2007)	26
<i>Bayer Schering Pharma AG v. Lupin, Ltd.</i> , 676 F.3d 1316 (Fed. Cir. 2012)	49, 50
<i>Bell Atl. Corp. v. Twombly</i> , 550 U.S. 544 (2007).....	27, 28, 35
<i>Bill of Lading Transmission & Processing Sys. Pat. Litig., In re</i> , 681 F.3d 1323 (Fed. Cir. 2012)	36
<i>Burlington Coat Factory Sec. Litig., In re</i> , 114 F.3d 1410 (3d Cir. 1997)	28
<i>DSU Med. Corp. v. JMS Co.</i> , 471 F.3d 1293 (Fed. Cir. 2006)	29, 30, 31
<i>GlaxoSmithKline LLC v. Teva Pharms. USA, Inc.</i> , 7 F.4th 1320 (Fed. Cir. 2021)	13, 24, 29, 40-43, 48, 49, 53
<i>Grunenthal GMBH v. Alkem Labs. Ltd.</i> , 919 F.3d 1333 (Fed. Cir. 2019)	23-24, 39, 45, 46, 49, 51, 52
<i>HZNP Meds. LLC v. Actavis Labs. UT, Inc.</i> , 940 F.3d 680 (Fed. Cir. 2019)	45-47, 49

<i>Kedra v. Schroeter</i> , 876 F.3d 424 (3d Cir. 2017)	36
<i>MEMC Elec. Materials, Inc. v. Mitsubishi Materials Silicon Corp.</i> , 420 F.3d 1369 (Fed. Cir. 2005)	29, 31
<i>Metro-Goldwyn-Mayer Studios Inc. v. Gorkster, Ltd.</i> , 545 U.S. 913 (2005).....	30, 31, 36
<i>Scheuer v. Rhodes</i> , 416 U.S. 232 (1974)	28
<i>Takeda Pharms. U.S.A., Inc. v. West-Ward Pharm. Corp.</i> , 785 F.3d 625 (Fed. Cir. 2015)	30, 47, 49
<i>Takeda Pharms. U.S.A., Inc. v. West-Ward Pharm. Corp.</i> , 188 F. Supp. 3d 367 (D. Del. 2016).....	50
<i>Takeda Pharms. U.S.A., Inc. v. West-Ward Pharm. Corp.</i> , 2016 WL 723054 (D. Del. Dec. 14, 2016)	50
<i>Tellabs, Inc. v. Makor Issues & Rts., Ltd.</i> , 551 U.S. 308 (2007).....	35, 36
<i>Visual Memory LLC v. NVIDIA Corp.</i> , 867 F.3d 1253 (Fed. Cir. 2017)	26
<i>Warner-Lambert Co. v. Apotex Corp.</i> , 316 F.3d 1348 (Fed. Cir. 2003)	30

Statutes	Pages
15 U.S.C. § 78u–4(b)(2)	35
21 U.S.C. § 355(j)(2)(A)(vii-viii).....	13
28 U.S.C. § 1295(a)(1)	3
28 U.S.C. § 1331	2
28 U.S.C. § 1338(a)	2
35 U.S.C. § 271(b)	29

35 U.S.C. § 271(e)(2)(A)	50
42 U.S.C. § 1983	36

Rules	Pages
21 C.F.R. § 314.94(a)(12)	13
Fed. R. Civ. P. 8(a)(2)	27, 35

TABLE OF ABBREVIATIONS AND CONVENTIONS

'537 patent	U.S. Pat. No. 9,700,537
'861 patent	U.S. Pat. No. 10,568,861
Appx _____	joint appendix page _____
Amarin	plaintiffs–appellants Amarin Pharma, Inc., Amarin Pharmaceuticals Ireland Ltd., and Mochida Pharmaceutical Co., Ltd., collectively
CV	cardiovascular
E-EPA	ethyl icosapentate (or icosapent ethyl)
EPA	icosapentaenoic acid
FAC	Amarin’s First Amended Complaint, Appx504-557
Hikma	defendants–appellees Hikma Pharmaceuticals USA Inc. and Hikma Pharmaceuticals PLC, collectively
HTG	hypertriglyceridemia
MTD	Hikma’s motion to dismiss first amended complaint for failure to state a claim under Fed. R. Civ. P. 12(b)(6), Appx941-968
SH	severe hypertriglyceridemia
TG	triglyceride
the asserted patents	the '537 and '861 patents
USPTO	United States Patent and Trademark Office
xx:yy-zz	column xx, lines yy-zz

RELATED CASES

No other appeals involving this civil action have been before this or any other appellate court. Appellants and their counsel are unaware of any other pending cases that will directly affect or be directly affected by the decision in this case.

INTRODUCTION

The generic drug approval scheme created by Congress permits a generic drug manufacturer like Hikma to copy the portion of a brand-name drug's label directed to a non-patented use of the drug so long as the generic manufacturer carves out any patented uses from its proposed label. If approved by the FDA, the generic manufacturer can market the drug, but only for the non-patented use indicated on the resulting "skinny label," which lists less than all the uses for which the brand-name drug has received approval. But even when the generic manufacturer proceeds with its skinny label, it still cannot encourage physicians to use the generic drug for the unauthorized, patented use that is not on the skinny label.

In this case, Hikma attempted to create a skinny label under this scheme. But Hikma, through multiple modes and media outside the label, encouraged prescribing physicians to replace Amarin's brand-name medication with Hikma's generic version for the patented use—the use for which Hikma's generic version was not approved. This was induced patent infringement.

The district court dismissed Amarin's complaint for failing to state a claim for induced infringement. In so doing, the district court misapplied the plausibility pleading standard by improperly considering Amarin's allegations in isolation instead of weighing them together; it made improper, implicit factual determinations

while bypassing the key factual dispute regarding what Hikma communicated to prescribing physicians; and it incorrectly analogized the case law.

Skinny-label precedent can involve difficult and close questions. The Court does not need to grapple with those questions here because this is not a true skinny-label case. Amarin's allegations are based not just on Hikma's label, but on its public statements made in press releases and on its website that encouraged physicians to prescribe Hikma's generic drug for an off-label use patented by Amarin. The district court erred when it considered each of those allegations one-by-one without considering whether it was at least plausible that Hikma's label together with its various public statements collectively encouraged infringement by prescribing physicians. And contrary to the district court's analysis, this Court's skinny-label precedent further supports inducement—especially *plausible* inducement—on the facts alleged. And the case law fully supports Amarin's view that the difficult questions and related factual disputes in this case cannot be resolved on a motion to a dismiss.

Reversal and remand is appropriate.

JURISDICTION

The district court had jurisdiction under 28 U.S.C. §§ 1331 and 1338(a), and entered partial final judgment pursuant to Fed. R. Civ. P. 54(b) on October 13, 2022.

Appx14-15; Appx507. Appellants filed a timely notice of appeal on November 9, 2022. Appx2069-2070. This Court has jurisdiction under 28 U.S.C. § 1295(a)(1).

STATEMENT OF ISSUES

1. Did the district court err by failing to consider the combined weight and effect of evidence demonstrating Hikma's repeated extra-label encouragement of using their generic version of Amarin's patented drug for *both* the approved skinny-label use of treating severe hypertriglyceridemia *and* the non-approved and infringing use of reducing cardiovascular risk in patients who do not suffer from severe hypertriglyceridemia when it dismissed Amarin's complaint for failure to state a claim?

2. Did the district court err by implicitly making a factual finding on the pleadings regarding what Hikma's conduct communicated to prescribing physicians, a key element of induced infringement?

3. Did the district court improperly analogize Amarin's allegations to *Grunenthal*, a label-only case where the asserted patent covered a use narrower than the generic label instructed, whereas Amarin alleged extra-label inducement activity by Hikma in the context of Amarin's patents, which are directed to a use broader than Hikma's approved use?

STATEMENT OF THE CASE

A. Vascepa[®] and its demonstrated efficacy for two different treatments

1. Severe hypertriglyceridemia, pancreatitis, and its treatment

Triglycerides, whether derived from food or made by the body, are a type of fat that circulates in human blood. While triglycerides are necessary for good health, high triglyceride levels, generally referred to as hypertriglyceridemia (“HTG”), can lead to serious conditions. Severe HTG¹ is a condition in which a patient’s blood triglyceride level (measured in milligrams of triglycerides per deciliter of blood, i.e., mg/dL) exceeds 500 mg/dL. Appx696 (FAC, Ex. K ¶1). The primary concern with severe HTG patients is pancreatitis. Appx952 (MTD, 5); Appx866 (FAC, Ex. BB at ¶7).

Vascepa[®] is a prescription drug Amarin developed and markets in the United States. Appx508 (FAC, ¶28). Its active ingredient is ethyl icosapentate (“E-EPA”) (sometimes referred to as icosapent ethyl), an ethyl ester of an omega-3 polyunsaturated fatty acid called icosapentaenoic acid (“EPA”) that occurs in and is

¹ Severe hypertriglyceridemia, or severe HTG, was referred to as “SH” before the district court, and the corresponding indication was referred to as the “SH indication.”

derived from fish oils. Appx1415 (R&R, 3); Appx40 ('537 patent, 3:14-40); Appx508 (FAC, ¶28).

Amarin initially studied Vascepa[®] in patients with severe HTG (the “Marine” trial). Appx508 (FAC, ¶30). The Marine trial demonstrated that Vascepa[®] lowers triglyceride levels in severe HTG patients without also raising levels of “bad” cholesterol, LDL-C. *Id.* That was a significant outcome because another drug indicated to reduce triglyceride levels—Lovaza[®]—can significantly increase bad cholesterol levels. Appx107 ('861 patent, 1:40-45). Following the Marine trial, the FDA approved Vascepa[®] as a treatment for severe HTG in 2012. Appx508 (FAC, ¶30). With that approval, Vascepa[®] became the first—and remains the only—medication approved for treating severe HTG that does not raise bad cholesterol levels. Appx508 (FAC, ¶30); *see* Appx107 ('861 patent, 1:40-45).

2. Amarin initially patented the use of Vascepa[®] to treat severe HTG

Amarin initially obtained patents covering only the use of Vascepa[®] to treat severe HTG (“the severe HTG patents”), but those patents are not at issue here. Amarin asserted the severe HTG patents against Hikma in 2017 after Hikma sought FDA approval to launch a generic form of Vascepa[®] for treating severe HTG. Appx980 (Opp., 6 n.2). In March 2020, the U.S. District Court for the District of Nevada found the severe HTG patents invalid as obvious, and, in September 2020, this Court affirmed without opinion. Appx948 (MTD, 1); *Amarin Pharma, Inc. v.*

Hikma Pharms. USA, Case No. 20-1723 (Fed. Cir. Sept. 3, 2020), ECF No. 78.

Hikma's generic version of Vascepa[®] was then approved by the FDA for—and only for—treating severe HTG. JA521 (FAC, ¶ 82).

3. Amarin later patented the use of Vascepa[®] to reduce cardiovascular risk in patients with non-severe HTG or existing cardiovascular disease

Non-severe HTG, or simply HTG, broadly refers to the condition in which a patient's triglyceride level is above the normal acceptable level of 150 mg/dL, which is less than one-third the 500 mg/dL triglyceride level associated with severe HTG. The primary concern when it comes to HTG, as opposed to severe HTG, is not pancreatitis. *See* Appx952 (MTD, 5); Appx866 (FAC, Ex. BB at ¶ 7). Instead, patients with HTG, like those with hypercholesterolemia (high cholesterol) or those with an established cardiovascular disease, are at risk for adverse cardiovascular events like a heart attack, i.e., myocardial infarction. *See* Appx40 ('537 patent, 3:14-40). Thus, unlike severe HTG where pancreatitis is the primary concern, the primary concern for patients with HTG and elevated LDL cholesterol levels is cardiovascular risk reduction. Appx866 (FAC, Ex. BB at ¶ 7).

After completing the Marine trial and receiving FDA approval for Vascepa[®] to treat severe HTG, Amarin continued its clinical work and studied whether Vascepa[®] could alternatively be used to treat patients with elevated triglyceride levels (200 - 500 mg/dL) and controlled LDL cholesterol levels in what was known

as the “Anchor” trial. *See* Appx509 (FAC, ¶31); Appx871 (FAC, Ex. BB at ¶18).

When the Anchor trial was designed, it was generally accepted that lowering triglyceride levels correlated with reduced cardiovascular risk. Appx509 (FAC, ¶31); Appx866 (FAC, Ex. BB at ¶7). The Anchor trial was thus designed to study whether Vascepa—as an add-on to a common cholesterol-lowering therapy using drugs called “statins” would lower triglyceride levels in patients having HTG, but not severe HTG. Appx509 (FAC, ¶31) (explaining that the Anchor trial studied patients with triglycerides levels between 200 mg/dL and 500 mg/dL). While the Anchor trial demonstrated that Vascepa[®] lowered triglyceride levels in those patients, the FDA did not approve Vascepa[®] for cardiovascular risk reduction because it concluded, based on the results of other intervening studies, that reduced triglyceride levels were *not* correlated with reduced cardiovascular risk. Appx509 (FAC, ¶32) (citing Appx863-881 (FAC, Ex. BB)); Appx871-872 (FAC, Ex. BB at ¶¶19-20).

In view of the FDA’s revised understanding that reduced triglyceride levels were not correlated with reduced cardiovascular risk, Amarin conducted a third clinical trial called the “REDUCE-IT” trial. Appx509 (FAC, ¶33). Rather than focusing on triglyceride levels, the REDUCE-IT trial studied whether Vascepa[®] could reduce cardiovascular events by following more than 8,000 patients over a median of five years. Appx509 (FAC, ¶33) (citing Appx832-843 (FAC, Ex. V)). As

in the Anchor trial, Vascepa[®] was evaluated in the REDUCE-IT trial as an add-on to statin therapy to determine its effect on reducing cardiovascular events in patients with elevated triglyceride levels (between 150 mg/dL and 499 mg/dL). Appx509 (FAC, ¶ 33); Appx832 (FAC, Ex. V at Methods). But unlike the Anchor trial, where results were based on measuring the patients' triglyceride levels, the REDUCE-IT trial directly studied whether Vascepa[®] reduced cardiovascular events by following patients with HTG and observing their clinical outcomes. *See* Appx509 (FAC, ¶ 33).

The REDUCE-IT trial was a success. The results, first announced in September 2018, showed a 25% further reduction in major cardiovascular events compared to patients on statin therapy alone. Appx509-510 (FAC ¶ 34) (citing Appx680-683 (FAC, Ex. H)). Those results were met with widespread enthusiasm and surprise in the field and were hailed as a “game changer.” Appx510 (FAC ¶ 35) (citing Appx852-853 (FAC, Ex. Y); Appx855-857 (FAC, Ex. Z)).

Based on the success of the REDUCE-IT trial, the FDA approved Vascepa[®] for a second indication: as a treatment to reduce cardiovascular risk in patients with HTG. *See* Appx509-510 (FAC, ¶ 34) (citing Appx685-689 (FAC, Ex. I)); Appx517 (FAC, ¶ 62) (citing Appx674-678 (FAC, Ex. G)). The Vascepa[®] label thus includes two approved indications, the earlier “Severe Hypertriglyceridemia Indication” for treating severe HTG and the later “CV Indication” for reducing cardiovascular risk in patients with HTG. Appx514 (FAC, ¶ 56) (citing Appx635-648 (FAC, Ex. D)).

After the second approval, Amarin removed language from its label that indicated Vascepa[®] had not been approved for cardiovascular risk reduction, i.e., the “CV Limitation of Use.” Appx514-517 (FAC, ¶¶ 60-63).

The FDA’s decision to approve Vascepa[®] for cardiovascular risk reduction was considered a “major milestone in cardiovascular prevention.” Appx518 (FAC, ¶ 66) (citing Appx685-689 (FAC, Ex. I)). Once Vascepa[®] received approval for cardiovascular risk reduction and the related limitation was removed from the Vascepa[®] label, “healthcare providers rapidly associated administration of [E-EPA] together with a statin as a method for reducing risk of cardiovascular events in patients with baseline triglycerides ≥ 150 mg/dL,” i.e., in patients with HTG. Appx519 (FAC, ¶ 67).

Based on work performed in connection with the REDUCE-IT trial, Amarin obtained U.S. Patent No. 10,568,861, which is one of the two patents at issue in this appeal and, unlike the previously litigated severe HTG patents, is directed to cardiovascular risk reduction. *See* Appx107 (’861 patent, 1:49-51); Appx128 (’861 patent, 43:7-14). Claims 1 and 2 of the ’861 patent cover methods for reducing the risk of cardiovascular death in patients with established cardiovascular disease by administering Vascepa:

1. A method of reducing risk of cardiovascular death in a subject with established cardiovascular disease, the method comprising administering to said subject about 4 g

of ethyl icosapentate per day for a period effective to reduce risk of cardiovascular death in the subject.

2. The method of claim 1, wherein the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL and a fasting baseline LDL-C level of about 40 mg/dL to about 100 mg/dL.

Appx129. On March 20, 2020, Amarin timely submitted information regarding the '861 patent to the FDA for listing in the Orange Book as covering methods of using Vascepa[®] to reduce cardiovascular risk. Appx520 (FAC, ¶ 77).

4. Mochida's prior foundational work and patent

Amarin's clinical trials related to Vascepa[®] were preceded by other work done by Mochida, a Japanese pharmaceutical manufacturer that later became Amarin's licensing partner, in the late 1990s and early 2000s. Appx510 (FAC, ¶ 36). Mochida established through its own cardiovascular outcomes trial that 1.8 grams per day of E-EPA, i.e., the same active ingredient as in Vascepa[®], suppressed certain cardiovascular risk in patients with high cholesterol, i.e., hypercholesterolemic patients. Appx510-511 (FAC, ¶ 36); Appx616 (FAC, Ex. B at Procedures). A statistical analysis of Mochida's trial results was later conducted to assess the effect of EPA on a Japanese patient population with a particular profile of risk factors for coronary artery disease, and its results were published in a 2008 article by Saito *et al.*, titled, "Effects of EPA on coronary artery disease in hypercholesterolemic patients with multiple risk factors: Sub-analysis of primary prevention cases from

the Japan EPA Lipid Intervention Study (JELIS),” 200 Atherosclerosis 135-400 (2008) (hereafter the “Saito article”). Appx511 (FAC, ¶¶37-40) (citing Appx615-620 (FAC, Ex. B)).

U.S. Patent No. 9,700,537, the other patent at issue in this appeal, is based on work performed in connection with the Saito article. The ’537 patent describes a method of reducing the risk of a cardiovascular event by administering EPA, with the active ingredient E-EPA, in combination with a statin to a patient with high cholesterol, elevated triglycerides, and reduced HDL-C (good cholesterol). Appx1416 (R&R, 4); Appx40 (’537 patent, 3:14-40). Claim 1 states:

1. A method of reducing occurrence of a cardiovascular event in a hypercholesterolemia patient consisting of:
 - identifying a patient having triglycerides (TG) of at least 150 mg/DL and HDL-C of less than 40 mg/dL in a blood sample taken from the patient as a risk factor of a cardiovascular event, wherein the patient has not previously had a cardiovascular event, and administering ethyl icosapentate in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor,
 - wherein said 3-hydroxyl-3-methylglutaryl coenzyme A reductase inhibitor is administered to the patient at least one of before, during and after administering the ethyl icosapentate; and
 - wherein the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor is selected from the group consisting of pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, rosuvastatin, and salts thereof, and
 - wherein daily dose of the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor are 5 to 60 mg for pravastatin, 2.5 to 60 mg for simvastatin, 10 to 180 mg

for fluvastatin sodium, 5 to 120 mg for atorvastatin calcium hydrate, 0.5 to 12 mg for pitavastatin calcium, 1.25 to 60 mg for rosuvastatin calcium, 5 to 160 mg for lovastatin, and 0.075 to 0.9 mg for cerivastatin sodium.

Appx46. The '537 patent is assigned to Mochida, and Amarin holds the exclusive license to it. Appx512 (FAC, ¶¶42-43). On January 9, 2020, Amarin timely submitted information regarding the '537 patent to the FDA for listing in the Orange Book as covering methods of using Vascepa[®] to reduce cardiovascular risk. Appx519 (FAC, ¶71).²

The '537 and '861 patents cover the CV Indication for Vascepa[®], and that CV Indication currently accounts for more than 90% of Vascepa[®] sales. Appx923-925 (Amarin's letter to its payer community after Hikma's generic launch); Appx540 (FAC, ¶152).

- B. Hikma launched generic Vascepa[®], describing it on its website, in press releases, and with a supposedly “skinny” label**
 - 1. Hikma submitted a Section viii statement to obtain FDA approval to sell generic Vascepa[®] solely to treat severe hypertriglyceridemia**

On September 21, 2016, Hikma (through its predecessor) submitted an abbreviated new drug application for a generic version of Vascepa[®]. Appx525 (FAC, ¶99). While Hikma's application was pending, the '537 and '861 patents issued. *See*

² The record below also involved U.S. Patent No. 8,642,077, but the parties resolved their dispute regarding that patent, and it is not relevant to this appeal.

Appx36 ('537 patent); Appx77 ('861 patent); Appx613 (FAC, Ex. A). Because Amarin listed both the '537 and '861 patents in the Orange Book before Hikma's generic version of Vascepa[®] was approved, Hikma was required to provide to the FDA either patent certifications under Section vii or a statement under Section viii. *See* 21 U.S.C. § 355(j)(2)(A)(vii-viii); 21 C.F.R. § 314.94(a)(12); Appx526 (FAC, ¶ 102). A "Section viii statement" (submitted under 21 U.S.C. § 355(j)(2)(A)(viii)), is filed when a generic applicant seeks FDA approval to label its drug only for uses *not* covered by method-of-use patents, like those at issue here. Appx524-525 (FAC, ¶ 95). Because the resulting generic's label would include less than the full label by virtue of the excluded patented indications, it is common to refer to a generic under a Section viii statement as a "skinny label" drug. *See GlaxoSmithKline LLC v. Teva Pharms. USA, Inc.*, 7 F.4th 1320, 1328 (Fed. Cir. 2021) (per curiam) (noting that Teva's label prepared under Section viii was "a so-called 'skinny label.'"). In this case, Hikma submitted a Section viii statement to the FDA with respect to the '537 and '861 patents, seeking FDA approval for only the severe HTG indication. Appx526 (FAC, ¶ 104).

After the severe HTG patents were invalidated in the Nevada action, and based on its Section viii statement, the FDA granted final approval for Hikma's ANDA on May 21, 2020. Appx506 (FAC, ¶ 11); Appx613 (FAC, Ex. A). Hikma

launched its generic version of Vascepa[®] in the United States in November 2020.

Appx506 (FAC, ¶ 13) (citing Appx715-717 (FAC, Ex. N)).


2. Hikma’s website associated its generic version of Vascepa[®] with a method of use beyond its approved use

When Hikma launched its generic version of Vascepa[®], with its approved indication limited to treating severe HTG, Hikma’s website announced that its product was in the Therapeutic Category of “Hypertriglyceridemia.” Appx532 (FAC ¶ 125) (citing Appx820 (FAC, Ex. T (Hikma’s website))). Treating patients having *severe* HTG—the approved indication—means treating patients with triglyceride levels of at least 500 mg/dL. But “hypertriglyceridemia,” i.e., HTG, as opposed to severe HTG, generally refers to patients having triglycerides of over 150 mg/dL who are at increased cardiovascular risk—the very patients studied in the REDUCE-IT trial for cardiovascular risk reduction. Hikma’s website thus promoted its generic EPA capsules as therapeutically equivalent for a use beyond treating patients with severe HTG. Appx532-533 (FAC, ¶ 126).³ Hikma has not explained why it declined to accurately describe the therapeutic category as Severe Hypertriglyceridemia.

³ Hikma knew that broader use would include the patented indication. In the Nevada trial based on Amarin’s severe HTG patents, Hikma acknowledged that there are ‘several reasons why a physician might prescribe Vascepa (or the Hikma Defendants’ ANDA Products) ... other than to treat severe hypertriglyceridemia,’ including to reduce cardiovascular risk.” Appx529 (FAC, ¶ 110) (citing Appx845-847 (FAC, Ex. W, ¶ 116)).

3. Hikma issued press releases, encouraging readers to prescribe generic Vascepa[®] for more than treating patients with severe hypertriglyceridemia

Beyond its website, Hikma also issued pre-launch press releases that carefully conveyed to readers that Hikma's product should be used for all uses for which Vascepa[®] was approved. First, the press releases indicated that Vascepa[®] is indicated only "in part" to treat patients with severe HTG. Appx709; Appx712. Hikma's press releases went further and identified its product as "Hikma's generic version" of Vascepa[®] without any qualification and in an attempt to equate Hikma's approval to the two Vascepa[®] indications. Appx709; Appx712. In fact, the later press release announced that Hikma had "received FDA approval of the product" without acknowledging that the approval was only for the severe HTG indication. Appx712. Taken together, Hikma made clear that Vascepa[®] was indicated for more than one use and then identified its own product as a generic version of Vascepa[®]. In describing its "generic version of Vascepa," both press releases further touted the value of *all* domestic Vascepa[®] sales, *id.*, even though Hikma knew more than 75% of sales were for the (patent-protected) CV Indication. Appx529 (FAC, ¶¶ 112-113); Appx531 (FAC, ¶¶ 119-120); Appx846 (¶ 115 (Hikma's proposed findings of fact from the Nevada trial)); Appx803 (FAC, Ex. Q). Hikma illustrated the lopsided sales in favor of the "separately patented" CV Indication during the Nevada trial in one of its demonstratives to downplay the commercial success of the SH Indication:



Appx803 (FAC, Ex. Q).

Amarin asserted those pre-launch press releases as evidence supporting its allegations that Hikma developed its product based on market assumptions that included the universe of Vascepa® sales, not just sales related to severe HTG treatment. Appx528 (FAC, ¶ 109). Importantly, Amarin alleged that Hikma’s press releases “communicate[d] to and instruct[ed] healthcare providers and patients that Hikma’s ‘generic version’ of VASCEPA® should be used for all the same indications as VASCEPA®, including to reduce the risk of [cardiovascular] events per the CV Indication awarded to VASCEPA®, and thus promote[d] and encourage[d] that use.” Appx530 (FAC, ¶ 115); Appx531 (FAC, ¶ 122). It was only after its launch that Hikma finally noted in a press release that its product had limited approval,

Appx715, but by that time Hikma had already primed the market with its earlier press releases.

4. Hikma’s “skinny” label teaches the claimed limitations and omits the prior limitation of use excluding the cardiovascular risk indication

Hikma’s label (Appx693-707) includes information that Amarin asserted would encourage a healthcare provider to prescribe its generic product for the patented and non-approved CV Indication. For example, the REDUCE-IT study—which was relevant solely to the CV Indication—is described in section 5.1 of Hikma’s label. Appx696. Co-administering with a statin (part of the CV Indication) is discussed in sections 12.3 and 14.2. Appx701-702. And Hikma itself recognized that the patient population for the severe HTG indication overlaps in part with the patient population for the cardiovascular risk indication. Appx803 (FAC, Ex. Q). Further, Hikma’s label identifies potential side effects, stating in part that people who have cardiovascular disease or diabetes with a risk factor for cardiovascular disease may experience heart rhythm problems when they take Hikma’s product:

Heart rhythm problems (atrial fibrillation and atrial flutter). Heart rhythm problems which can be serious and cause hospitalization have happened in people who take icosapent ethyl, especially in people who have heart (cardiovascular) disease or diabetes with a risk factor for heart (cardiovascular) disease, or who have had heart rhythm problems in the past.

Appx704-705 (FAC, Ex. K).

Hikma's label makes clear that: "Medicines are sometimes prescribed for purposes other than those listed in a Patient Information leaflet." Appx535 (FAC, ¶132); Appx705. One of Hikma's physician experts in the Nevada trial pointed to that language when explaining that "most often we use this medication for reasons other than the [severe HTG clinical trial] data, and in the patient information section it specifically tells the patients that we would potentially do that." Appx535 (FAC, ¶132) (citing Appx849-850 (FAC, Ex. X at 617)). In other words, Hikma's physician understood that the label tells readers that Hikma's product will be prescribed for reasons other than treating patients with severe HTG.

Beyond including information relevant to the CV Indication, Hikma removed a relevant Limitation of Use from its label. When Hikma submitted its ANDA (September 21, 2016), Vascepa[®] was approved only for the severe HTG indication, and the Vascepa[®] label included a statement in its Limitation of Use section, which said: "The effect of VASCEPA on cardiovascular mortality and morbidity in patients with severe hypertriglyceridemia has not been determined." Appx514-515 (FAC, ¶60) (citing Appx650-661 (FAC, Ex. E); Appx663-672 (FAC, Ex. F)). Hikma's proposed label thus included the same Limitation of Use.

The FDA's approval of Vascepa[®] for the cardiovascular indication allowed *Amarin* to add a cardiovascular risk reduction indication to, and remove the cardiovascular Limitation of Use from, the Vascepa[®] label. However, *Hikma* also

removed the cardiovascular Limitation of Use from its proposed label and did so with knowledge of the '537 and '861 patents. Appx528 (FAC, ¶108). When Hikma launched its product, its label did not include any cardiovascular Limitation of Use. *See* Appx526 (FAC, ¶107) (citing Appx694-707 (FAC, Ex. K)). Amarin alleged that Hikma removed that Limitation of Use “so that healthcare providers and patients would believe that Hikma’s generic [E-EPA] capsules could be and should be used just like VASCEPA[®], including to reduce the risk of cardiovascular events per the CV Indication awarded to VASCEPA[®].” Appx528 (FAC, ¶108). That allegation is well founded because the market understands what the presence of the cardiovascular Limitation of Use implies because another similar well-known drug, Lovaza[®], which contains EPA and lowers triglyceride levels, includes the same limitation and has not been shown to be effective for reducing cardiovascular risk. Appx516 (FAC, ¶61) (citing Appx807-818 (FAC, Ex. S)); Appx518 (FAC, ¶64).

C. The district court proceeding

1. Amarin sued Hikma for inducing infringement

On November 30, 2020, Amarin sued Hikma for inducing infringement of the '861 and '537 patents (“the asserted patents”), Appx159-166 (Compl., ¶¶120-143), and on January 25, 2021, Amarin filed its First Amended Complaint (“FAC”), Appx504-557. Amarin alleged that the combination of Hikma’s label, its press releases, and its website encouraged healthcare providers to: (i) associate Hikma’s

generic with the patented use of Vascepa[®] known to the market; and (ii) administer Hikma's generic version of Vascepa[®] to patients with non-severe HTG in order to reduce cardiovascular risk. Appx533 (FAC, ¶¶ 127-128).⁴

2. Hikma moved to dismiss under Rule 12(b)(6)

Hikma moved to dismiss the inducement claims under Federal Rule of Civil Procedure 12(b)(6) for failure to state a claim. Appx941-968 (MTD). Hikma did *not* dispute that Amarin sufficiently alleged that healthcare providers had directly infringed the '537 and '861 patents or that Hikma knew about the asserted patents and the direct infringement. And, Hikma admitted that the district court could consider Amarin's allegations about Hikma's website, press releases, and label. Appx950 (MTD, 3) ("Now that Hikma recently launched its product, Amarin can rely on information outside of the labeling to prove inducement."). Rather, Hikma's motion was limited to arguing that Amarin did not plausibly allege that Hikma took "active steps" to encourage that infringement, despite Amarin's allegations regarding Hikma's website and press releases in addition to its label. Appx945 (MTD, ToC); Appx1011 (Reply, 10).

⁴ Amarin amended its complaint once to add allegations, mostly related to Amarin's decision to add another defendant: Health Net, LLC. Appx558. Health Net separately moved to dismiss the first amended complaint, and Health Net's motion was denied. Amarin and Health Net subsequently resolved their dispute, so Health Net is not part of or related to this appeal.

3. The magistrate judge recommended denying Hikma’s motion to dismiss

The magistrate judge recommended denying Hikma’s motion to dismiss. Appx503; Appx1430 (R&R, 18). Recognizing that it could not “make factual findings about what Hikma’s label and advertisements communicate to physicians,” Appx1414 (R&R, 2), the magistrate judge explained why Amarin’s allegations could not be resolved on the pleadings. The impact of Hikma’s actions on healthcare providers raised a factual dispute that could plausibly be resolved in Amarin’s favor:

Hikma urges the Court to resolve this case at the pleadings stage, pointing out that the contents of its label and public statements are undisputed. *But there is a real dispute about what those contents communicate to others*, and I do not think it is appropriate to resolve it on a motion to dismiss. Stated another way, at this stage of the case, I am not prepared to say that Hikma’s label and public statements—as a matter of law—could never amount to instruction and encouragement to infringe the asserted patents.

Appx1427 (R&R, 15) (emphasis added).

The magistrate judge agreed with Hikma that it had no duty to actively discourage infringement, and further agreed that Hikma’s mere knowledge of direct infringement would be insufficient on its own to state a claim. Appx1426 (R&R, 14). “But [Hikma] cannot present information in a way that encourages infringement.” Appx1426 (R&R, 14). Amarin plausibly alleged that is exactly what Hikma did. The magistrate judge recognized that Amarin’s various allegations

should be considered together, not individually, and viewed in a light most favorable to Amarin. *See* Appx1424-1425 (R&R, 12-13). As the magistrate judge explained, “[t]he assessment of whether a complaint plausibly alleges inducement in a pharmaceutical case is thus no different than the analysis in any other case.” Appx1423 (R&R, 11).

4. The district court granted Hikma’s motion

The district court overruled the magistrate judge’s recommendation and granted Hikma’s motion to dismiss. Appx2. Beginning with Hikma’s label, the district court explained that it could find no instruction “as to CV risk reduction.” Appx6. In analyzing Hikma’s label, the district court focused on “CV risk reduction,” and not the claimed patient population. In so doing, the “side effect” language was “hardly instruction or encouragement” to use Hikma’s generic drug to reduce cardiovascular risk. Appx6. The district court opinion did not discuss whether the “side effect” language would be understood by healthcare providers as encouraging them to prescribe Hikma’s generic drug to the patient population claimed in the ’537 or ’861 patents. And the district court found that Hikma had no duty to discourage the patented use. Appx7. Even if removing the Limitation of Use communicated to the public that Hikma’s generic drug could be used to reduce cardiovascular risk, the district court reasoned that healthcare providers would

understand that as Hikma “merely describing” an infringing mode, not encouraging it. Appx7.

The district court then set the label aside to separately analyze the non-label allegations: “Since I find that the label does not instruct CV risk reduction, the question is whether Hikma’s public statements, including press releases and Hikma’s website, induce infringement.” *Id.*

As to Hikma’s pre-launch press releases, the district court concluded that advertising its product as the “generic equivalent” of Vascepa[®] did not expose Hikma to liability under *GlaxoSmithKline* and citing sales figures for infringing uses goes to Hikma’s “intent to induce” but does not count as an inducing act. Appx8.

Turning to Hikma’s website that advertised Hikma’s product in the “hypertriglyceridemia” therapeutic category, the district court acknowledged that “Amarin has pled that the category ‘hypertriglyceridemia’ includes infringing uses.” Appx8. Again, isolating the analysis, the district court reasoned “[t]he question is whether this is enough, without a label or other public statements instructing as to infringing use, to induce infringement,” and decided the answer was “no.” Appx8.

To justify its conclusion, the district court compared the facts here to the facts in *GlaxoSmithKline* (finding infringement) and *Grunenthal GMBH v. Alkem Labs. Ltd.*, 919 F.3d 1333 (Fed. Cir. 2019) (finding no infringement), and concluded this case was more like *Grunenthal* than *GlaxoSmithKline*. Appx8-9. The district court

attempted to distinguish Teva’s press releases in *GlaxoSmithKline*, which described the Teva product as a generic “cardiovascular agent,” a category that included infringing and non-infringing uses, from Hikma’s website. Appx9. The district court found Hikma’s website was less like the press releases in *GlaxoSmithKline* and more like the label at issue in *Grunenthal*, where the label included a broader, non-patented category, which by definition covered a narrower, patented use. Appx9. In *Grunenthal*, this Court found that listing the non-patented, broader category on the label was not a specific encouragement to use the drug for the narrower, patented purpose. *Grunenthal*, 919 F.3d at 1339; *see* Appx9. The district court did not comment on the distinction between *Grunenthal* and this case, namely, that the relationship between the non-patented use and patented use is reversed: the non-patented use here (i.e., the severe HTG indication) is *narrower* than the patented use (i.e., the cardiovascular risk reduction indication for general HTG). Hikma’s website did not list a non-patented use that included a narrower patented use like the *Grunenthal* label; instead, Hikma listed a broader general HTG therapeutic category where the majority of prescriptions would be for the patented use to reduce cardiovascular risk. *See* Appx820 (FAC, Ex. T); Appx529 (FAC, ¶¶ 112-113); Appx531 (FAC, ¶¶ 119-120).

The district court concluded that Amarin “failed to plead inducement based on Hikma’s label or public statements” and granted Hikma’s motion to dismiss the first amended complaint. Appx9.

SUMMARY OF ARGUMENT

This case involves Hikma’s so called skinny-label generic version of Amarin’s breakthrough Vascepa[®] medication, but the evidence in this case goes beyond the label, skinny or not. This is a pleadings standard case. Yes, Amarin alleged that Hikma’s skinny label was not skinny enough, i.e., that it covered patented methods of treatment. But Amarin alleged more than that.

Amarin alleged that Hikma, knowing that the “vast majority” of Vascepa[®] prescriptions were for the patented breakthrough cardiovascular risk reduction treatment, undertook a series of communications with the market with the intent that its generic be used to replace Vascepa[®] for the patented use. Hikma broadcast a broader therapeutic category (hypertriglyceridemia) than it had approval for (severe hypertriglyceridemia) and then emphasized the generic equivalency of its product with Vascepa[®]—most often used for cardiovascular risk reduction—through Hikma’s website, multiple press releases, and the language of Hikma’s label. It is more than plausible that Hikma intended these active steps to influence prescribing physicians to replace Vascepa[®] with its generic for the patented use. That was

enough to satisfy the pleading requirements and survive the motion to dismiss. Holding otherwise resulted in multiple errors.

The district court erred by: (1) weighing Amarin's allegations separately and in isolation against the plausibility pleading standard rather than considering whether, as Amarin pled, Hikma's conduct as a whole induced infringement; (2) making implicit factual findings on the key question of what Hikma's conduct communicated to prescribing physicians; and (3) misapplying skinny-label precedent. Those errors effectively and improperly elevated the pleading standard to deprive Amarin of its right to pursue a more than plausible claim for induced infringement. Reversal and remand is appropriate.

ARGUMENT

I. Standard of review

This Court reviews motions to dismiss for failure to state a claim under the law of the regional circuit. *Visual Memory LLC v. NVIDIA Corp.*, 867 F.3d 1253, 1257 (Fed. Cir. 2017). The Third Circuit reviews the district court's grant of such a motion de novo. *Ballentine v. United States*, 486 F.3d 806, 808 (3d Cir. 2007).

II. Amarin satisfied the pleading standard by pleading plausible infringement by Hikma

The requirement to state a plausible claim for relief at the pleading stage is not demanding. The specific claim here, induced infringement, requires encouragement or promotion of an infringing use. Amarin pled multiple facts that

together demonstrated how Hikma encouraged or promoted prescribing physicians to replace Amarin's Vascepa[®] with Hikma's generic for the patented use of reducing cardiovascular risk in patients with elevated triglyceride levels.

A. Surviving a motion to dismiss requires pleading facts sufficient to state a plausible claim for relief

“To survive a motion to dismiss, a complaint must contain sufficient factual matter, accepted as true, to ‘state a claim to relief that is plausible on its face.’” *Ashcroft v. Iqbal*, 556 U.S. 662, 678 (2009) (quoting *Bell Atl. Corp. v. Twombly*, 550 U.S. 544, 570 (2007)). Only “a short and plain statement of the claim showing that the pleader is entitled to relief” is required. Fed. R. Civ. P. 8(a)(2). Factual allegations are reviewed “on the assumption that all the allegations in the complaint are true (even if doubtful in fact).” *Twombly*, 550 U.S. at 555. The allegations do not need to be detailed. *Id.* Instead, “[f]actual allegations must be enough to raise a right to relief above the speculative level,” as opposed to allegations that provide a “formulaic recitation” of the claim elements. *Id.*

This was not a case where Amarin merely speculated about what doctors might do. Instead, Amarin set forth allegations of how Hikma used multiple communications on its website, press releases, and label together to encourage using its generic as a replacement for Vascepa[®] for the patented use. Those allegations easily exceeded mere speculation.

A claim is plausible when the complaint contains “factual content that allows the court to draw the reasonable inference that the defendant is liable for the misconduct alleged.” *Iqbal*, 556 U.S. at 678 (citing *Twombly*, 550 U.S. at 556). The plausibility requirement is not a “probability requirement.” *Twombly*, 550 U.S. at 556. “[I]t simply calls for enough fact to raise a reasonable expectation that discovery will reveal evidence” showing the alleged misconduct. *Id.* “[A] well-pleaded complaint may proceed even if it strikes a savvy judge that actual proof of those facts is improbable, and ‘that a recovery is very remote and unlikely.’” *Id.* (quoting *Scheuer v. Rhodes*, 416 U.S. 232, 236 (1974)). “The issue is not whether a plaintiff will ultimately prevail but whether the claimant is entitled to offer evidence to support the claims.” *In re Burlington Coat Factory Sec. Litig.*, 114 F.3d 1410, 1420 (3d Cir. 1997).

Amarin alleged facts showing how Hikma’s multiple communications plausibly encouraged infringement. Those allegations highlighted a key factual dispute over how prescribing physicians would understand Hikma’s multiple communications—a dispute that would be informed by discovery and expert testimony. But even without that discovery and testimony—prematurely cutoff by the district court—Amarin’s alleged facts were enough to create a “reasonable inference,” *Iqbal*, 556 U.S. at 678, that Hikma’s multiple communications encouraged physicians to use Hikma’s generic for the patented use.

B. Inducement requires showing direct infringement and actions taken with the intent to cause infringing conduct

Section 271(b) of Title 35 provides that “[w]hoever actively induces infringement of a patent shall be liable as an infringer.” 35 U.S.C. § 271(b). To state a claim of induced infringement under § 271(b), the complaint must plausibly allege that: (1) there has been direct infringement; (2) the defendant knowingly induced infringement; and (3) the defendant possessed the intent to encourage another’s infringement. *MEMC Elec. Materials, Inc. v. Mitsubishi Materials Silicon Corp.*, 420 F.3d 1369, 1378 (Fed. Cir. 2005).

A generic manufacturer can be liable for inducing infringement even when it has attempted to “carve out” the patented indications with a skinny label. *GlaxoSmithKline*, 7 F.4th at 1338; *see AstraZeneca LP v. Apotex, Inc.*, 633 F.3d 1042, 1056-61 (Fed. Cir. 2010) (affirming grant of preliminary injunction against generic manufacturer for inducing infringement of the patented use even though generic product was approved for the non-patented use). Thus, in considering a motion to dismiss in a pharmaceutical case like this, the court must still determine whether the complaint plausibly alleges inducement.

The inducement inquiry considers whether the complaint plausibly alleges that the generic manufacturer “offer[ed] a product with the object of promoting its use to infringe, as shown by clear expression or other affirmative steps taken to foster infringement.” *DSU Med. Corp. v. JMS Co.*, 471 F.3d 1293, 1305-06 (Fed. Cir.

2006) (en banc in relevant part). And the Supreme Court has held that advertising or instructing an infringing use is evidence of active steps that both encourage infringement and demonstrate the affirmative intent to induce infringement. *Metro-Goldwyn-Mayer Studios Inc. v. Gorkster, Ltd.*, 545 U.S. 913, 936 (2005); see *Takeda Pharms. U.S.A., Inc. v. West-Ward Pharm. Corp.*, 785 F.3d 625, 630-31 (Fed. Cir. 2015) (“Inducement can be found where there is ‘[e]vidence of active steps taken to encourage direct infringement,’ which can in turn be found in ‘advertising an infringing use or instructing how to engage in an infringing use.’”) (quoting *Gorkster*, 545 U.S. at 936). Rather than alleging a defendant’s “mere knowledge” that its product could be used to infringe, *Warner-Lambert Co. v. Apotex Corp.*, 316 F.3d 1348, 1364 (Fed. Cir. 2003), allegations should plausibly suggest “culpable conduct, directed to encouraging another’s infringement.” *DSU Med.*, 471 F.3d at 1306.

C. Taking the allegations together, Amarin’s claim against Hikma is plausible, and discovery is likely to further support Amarin’s case

Under the plausibility pleading standard, Amarin was not required to prove its inducement claim to survive the motion to dismiss. Amarin was only required to plead a plausible claim for relief. Amarin did so.

Because Hikma did not dispute either that its generic form of Vascepa[®] was being used to reduce cardiovascular risk or that such use directly infringed the asserted patents, the issue before the district court was whether Amarin had pled

facts plausibly showing Hikma acted to induce that infringement. *MEMC Elec. Materials*, 420 F.3d at 1378. Amarin pled multiple facts that plausibly demonstrated how Hikma’s cumulative actions were taken with the intent to encourage and promote infringement. *Grokster*, 545 U.S. at 936; *DSU Med.*, 471 F.3d at 1305-06.

First, on its website, Hikma described its generic as being therapeutically equivalent to Vascepa® in a broad indication category of “hypertriglyceridemia,” i.e., HTG generally. Appx532 (FAC ¶ 125) (citing Appx820 (FAC, Ex. T (Hikma’s website))). That broad category exceeds Hikma’s authorized use to treat *severe* HTG in patients with triglyceride levels above 500 mg/dL because it includes the unauthorized, patented use to reduce cardiovascular risk in patients with triglyceride levels above 150 mg/dL, i.e., patients who suffer from HTG (where the primary concern is cardiovascular risk reduction) but not *severe* HTG (where the primary concern is pancreatitis). Appx532-533 (FAC, ¶¶ 125, 126). Thus, Hikma communicated to the market that its generic was equivalent to Vascepa® for the patented use.

Second, Hikma’s marketing campaign included press releases that communicated to the market that Hikma’s generic was equivalent to Vascepa® without limitation on use. At the close of the Nevada trial over the severe HTG patents, Hikma announced that it was seeking approval for “its generic version of Vascepa®.” Appx709-710. That press release did not divulge that Hikma was

seeking approval only for a particular use of Vascepa[®]. To the contrary, Hikma identified that particular use as only among the indications for which Vascepa[®] was approved. *Id.* (describing Vascepa[®] as “a prescription medicine that is indicated, in part, ... to reduce triglyceride levels in adult patients with severe ... hypertriglyceridemia.”). When Hikma received approval, it announced that the FDA had approved “its [E-EPA] Capsules, 1 gm, *the generic equivalent to Vascepa.*” Appx613 (emphasis added). There is no equivocation in that press release, which speaks in terms of the approval of a drug, not a use, and fails to acknowledge that the FDA had approved its generic only for a single indication. And then when Hikma won on appeal following the Nevada trial, it again announced it had received approval for its generic version of Vascepa[®] without mentioning that approval was limited to the less common indication. Appx712-713. Beyond its silence about the nature of its limited approval, Hikma added that “US sales of Vascepa were approximately \$1.1 billion in the 12 months ending in July 2020.” *Id.* That figure was extraordinarily misleading in context because, as discussed below, Hikma was aware that those sales were almost all for the use of Vascepa[®] to treat cardiovascular risk in patients with HTG, a use for which Hikma could not legally compete.

Third, in making these associations, Hikma was aware that the vast majority of prescriptions for Vascepa[®] were for reducing cardiovascular risk and not for treating severe HTG—the only indication for which its generic is approved. In the

earlier Nevada trial involving the severe HTG patents, Hikma made such an admission even before the FDA had given Amarin approval to market Vascepa® for that use. *See, e.g.*, Appx846 ¶115 (Hikma’s proposed finding of fact that “the vast majority of Vascepa prescriptions—over 75%—have been for . . . treating patients with triglycerides below 500 mg/dL.”); Appx847 ¶440 (Hikma’s proposed finding of fact that “the ‘vast majority’ of Vascepa® prescriptions are off-label, to patients with triglyceride levels lower than 500 mg/dl”); Appx528-529, Appx535 (FAC, ¶¶110, 132). Following FDA approval of treating cardiovascular risk, that second indication accounted for more than 90% of uses of Vascepa®. Appx923-925 (Amarin’s letter to its payer community after Hikma’s generic launch); Appx540 (FAC, ¶152). And Hikma knew that doctors had “rapidly associated” Vascepa® with treatment of patients with HTG to reduce cardiovascular risk. Appx519 (FAC, ¶67). Thus, Hikma’s communications, which failed to mention that limited nature of the FDA’s approval, relied on that association between Vascepa® and the patented use for which Hikma’s generic equivalent was not authorized.

Fourth, while Hikma removed some instructions for the unauthorized use from its supposedly skinny label, it maintained in the clinical studies section descriptions of statin-treated patients with the same cardiovascular event history and lipid levels covered by the patented methods. Appx534-536 (FAC, ¶¶130-131, 134); Appx702 (FAC, Ex. K §14.2). And the well-known REDUCE-IT trial related to

cardiovascular risk reduction was described on the label too. Appx696 (FAC, Ex. K § 5.1).

Fifth, Hikma omitted the CV Limitation of Use from its label, a limitation that appears on the only other medication on the market that contains E-EPA and is indicated for lowering triglyceride levels, Lovaza[®]. Appx514-517 (FAC, ¶¶ 60-61); Appx527-528 (FAC, ¶108). The absence of this limitation, present on the Lovaza[®] label and understood by physicians, was another way the label communicated to the market that Hikma's generic could be used for the unauthorized, patented cardiovascular risk reduction indication. Appx535-536 (FAC, ¶133).

Sixth, Hikma's label identifies a patient population with triglyceride levels above 150 mg/dL that overlaps with patients being treated for cardiovascular risk. Appx696 (FAC, Ex. K § 1).

Far from a purely speculative or formulaic recitation of claim elements, Amarin's complaint presents a clear theory of how Hikma's communications plausibly encouraged or promoted the patented and unapproved use of Hikma's generic version of Vascepa[®] to reduce cardiovascular risk in patients with elevated triglyceride levels. As the magistrate judge recognized, those factually supported allegations cleared the low pleading bar and prevented dismissal. But rather than reviewing the pled facts and actions together against the low bar of plausibility, the district court considered each fact in isolation. Even if no pled fact by itself could

support the reasonable inference that Hikma has induced infringement, that is irrelevant. Amarin pled that Hikma's multiple communications together encouraged and promoted doctors' use of Hikma's generic for the patented and unapproved use. Those multiple pled facts cleared the low bar for plausibility.

III. The district court erred by weighing the pled facts piecemeal against the plausibility pleading standard

By considering each factual allegation in isolation, the district court transformed the inquiry from one requiring "a short and plain statement of the claim showing that the pleader is entitled to relief," Fed. R. Civ. P. 8(a)(2), into an exacting inquiry requiring a single act to sustain the plausibility of Amarin's case. That was error.

The plausibility pleading standard simply requires sufficient factual material to show the claim for relief is plausible on its face, *Iqbal*, 556 U.S. at 678, to "raise a right to relief above the speculative level," *Twombly*, 550 U.S. at 555, and to allow "the court to draw the reasonable inference that the defendant is liable for the misconduct alleged," *Iqbal*, 556 U.S. at 678. It does not require pleading a single fact that by itself establishes the plausibility of a claim because "courts must consider the complaint in its entirety." *Tellabs, Inc. v. Makor Issues & Rts., Ltd.*, 551 U.S. 308, 322 (2007). In the context of the pleading requirement for scienter under the Private Securities Litigation Reform Act, 15 U.S.C. § 78u-4(b)(2), the Supreme Court explained, "The inquiry . . . is whether all of the facts alleged, taken

collectively, give rise to a strong inference of scienter, not whether any individual allegation, scrutinized in isolation, meets that standard.” *Id.* at 322-23.

The Third Circuit applied the Supreme Court’s rule in the context of a civil rights complaint under 42 U.S.C. § 1983 when it reversed the district court’s grant of a motion to dismiss for failure to state a claim. *Kedra v. Schroeter*, 876 F.3d 424, 432 (3d Cir. 2017). In reversing, the court explained that the district court had “done the inverse of what we are required to do at the pleading stage,” i.e., “[i]nstead of considering the complaint as a whole, [the district court] consider[ed] ‘whether any individual allegation, scrutinized in isolation, meets that standard.’” *Id.* at 444 (quoting *Tellabs*, 551 U.S. at 322-23). Inducement is similarly not limited to a single act. It considers whether the defendant took “active steps . . . to encourage direct infringement.” *Grokster*, 545 U.S. at 936. *Cf. In re Bill of Lading Transmission & Processing Sys. Pat. Litig.*, 681 F.3d 1323, 1343 (Fed. Cir. 2012) (finding induced infringement plausible and that the district court erred by concluding otherwise where “the district court analyzed the individual facts in the [] Amended Complaint in isolation and without reference to the background of the invention”). Thus, establishing a plausible inducement claim does not require a single act.

But that is effectively what the district court required here. It decoupled Hikma’s actions to determine whether one portion of Hikma’s conduct in *isolation* encouraged infringement instead of considering—as Amarin pled—whether

Hikma's *cumulative* conduct encouraged infringement. *See* Appx5 ("These allegations fall into two categories . . ."); Appx6 ("The bulk of the briefing and oral argument was directed to Hikma's label, and I will address those arguments first."); Appx9 ("Amarin's complaint has failed to plead inducement based on Hikma's label or public statements . . ."). To be clear, Amarin maintains that Hikma's press releases, its website, and even its label are enough to clear the pleading standard even if considered in isolation. But even if each piece were not enough in isolation, the district court further erred by not considering them collectively.

This error effectively raised the pleading requirement by weighing the allegations separately. The district court isolated the allegations of inducement based on the label from the allegations based on the other public statements. *See* Appx7 ("Since I find that the label does not instruct CV risk reduction, the question is whether Hikma's public statements, including press releases and Hikma's website, induce infringement."). But the district court went even further and isolated the allegations of inducement for portions of Hikma's public statements from other portions. *See* Appx8 ("Amarin has pled that the category 'hypertriglyceridemia' includes infringing uses. The question is whether this is enough, *without a label or other public statements* instructing as to infringing use, to induce infringement.") (emphasis added).

In contrast, the magistrate judge’s report and recommendation fairly considered the cumulative allegations as Amarin pled. The magistrate judge noted how Amarin alleged that “portions of Hikma’s label, *taken together* with Hikma’s public statements, instruct physicians to use Hikma’s product in a way that infringes the asserted patents.” Appx1424 (R&R, 12) (emphasis added). And rejecting Hikma’s arguments about label-only ANDA case law, the magistrate judge explained, “were this an ANDA case, and were [Amarin’s] allegations based solely on the label, [Amarin’s] inducement theory might lack merit as a matter of law. But this is not an ANDA case, and *[Amarin’s] allegations are not based solely on the label.*” Appx1426 (R&R, 14) (emphasis added). Thus, the magistrate judge concluded that, when “*taken together* and viewed in the light most favorable to [Amarin],” Amarin’s allegations “plausibly suggest . . . that Hikma’s *label and public statements* could instruct and/or encourage third parties to use its product for the CV indication, which [Amarin] allege[s] is covered by the asserted patents; and [] that Hikma both knew and intended that third parties would use its product for that purpose.” Appx1425 (R&R, 13) (emphasis added).

The report and recommendation properly applied the pleading standard based on what Amarin pled, but the district court did not. This Court should reverse.

IV. The district court erred by resolving the key factual dispute of what Hikma’s conduct communicated to the market

Although “the contents of [Hikma’s] label and public statements are undisputed,” “there is a real dispute about what those contents communicate to others, and I do not think it is appropriate to resolve it on a motion to dismiss.” Appx1427 (R&R, 15). With this explanation, the magistrate judge foreshadowed another error by the district court.

Infringement is a fact question. *Grunenthal GMBH v. Alkem Labs. Ltd.*, 919 F.3d 1333, 1339 (Fed. Cir. 2019). In evaluating Amarin’s allegations of induced infringement, the key issue is whether Hikma acted to encourage or promote infringement with the intent to cause such infringement. And as the magistrate judge explained, Amarin alleged that “several . . . portions of Hikma’s label, taken together with Hikma’s public statements, *instruct physicians* to use Hikma’s product in a way that infringes the asserted patents.” Appx1424 (R&R, 12) (emphasis added). Thus, the key factual dispute was what Hikma’s conduct as described in Amarin’s factual allegations communicated to prescribing physicians. Although the magistrate judge recognized this key disputed issue—which should have precluded dismissal—the district court ignored it.

Instead of turning to discovery and expert testimony to illuminate how physicians would have understood Hikma’s collection of communications through its website, press releases, and label, the district court short-circuited the process and

implicitly decided the key factual dispute on a motion to dismiss. Appx536-537 (FAC, ¶ 135) (“For all the reasons set forth above, Hikma knows of and specifically intends for healthcare providers to administer its [E-EPA] capsules in the place of VASCEPA® . . . and its labeling and marketing materials promote, encourage, and instruct healthcare providers to practice the methods of the Asserted Patents”). The district court made that inappropriate finding without addressing the question of what was communicated to physicians. *See* Appx6 (determining that Hikma’s specific warning about side effects was not “instruction or encouragement”); Appx7 (determining that “the lack of a [cardiovascular] limitation on Hikma’s label does not plausibly teach [cardiovascular] risk reduction”); Appx8 (determining that Hikma’s broad category listing on its website “does not rise to the level of encouraging, recommending, or promoting taking Hikma’s generic for the reduction of [cardiovascular] risk”).

GlaxoSmithKline is illustrative. There, the factfinder reviewed exactly what the district court in this case prevented Amarin from presenting—discovery and expert testimony addressing what Hikma’s promotional activities and label communicated to prescribing physicians. This Court extensively discussed the expert testimony. *GlaxoSmithKline*, 7 F.4th at 1328-38. In doing so, the Court noted how “GSK’s cardiology expert, Dr. McCullough, explained that doctors, the alleged direct infringers, receive information about generic drug products from a variety of

sources, including the drug labels,” and “then walked through each element of claim 1 of the [asserted] patent and compared it to Teva’s partial label.” *Id.* at 1328.

The fact that *GlaxoSmithKline* proceeded through discovery to trial highlights the evidence-depriving error that dismissal in the face of disputed key factual questions caused in this case. And *GlaxoSmithKline* even addressed a parallel error: “Critically, the district court erred by treating this fact question—whether [a certain] indication instructs a physician to prescribe carvedilol for a claimed use—as though it were a legal one for it to decide *de novo*.” *Id.* at 1330. Although that case went to trial, the district court usurped the jury’s factfinding role on JMOL and “decided the [relevant indication] portion of Teva’s label was insufficient to find that the label instructed an infringing use.” *Id.* As explained above, the district court in this case made multiple similar *factual* determinations, assessing the weight of the evidence without explicitly considering the key factual question of what that evidence communicated to prescribing physicians.

For “a quintessential fact question,” *id.* at 1328, involving “a real dispute about what [Hikma’s label and public statements] communicate to others,” it was “not . . . appropriate to resolve [] on a motion to dismiss,” Appx1427 (R&R, 15). The district court erred when it did so.

V. The district court erred by analogizing the wrong cases

The district court also erred in its misapplication of skinny-label precedent.

A. This case involves evidence showing inducement in addition to the label—it is not a “label only” case

This case is like *GlaxoSmithKline* and *AstraZeneca* in that those cases also included extra-label evidence to show inducement. In *GlaxoSmithKline*, this Court vacated a grant of JMOL for non-infringement and reinstated a jury’s verdict finding induced infringement. 7 F.4th at 1323. As discussed above, the Court relied on detailed expert testimony from Dr. McCullough, including the assertion that doctors obtain information about generic medications from drug labels and other sources. *Id.* at 1328-38. The Court found evidence showing that “Teva intended its label to affect physician’s prescribing practices,” but that was “not the only evidence” because “GSK also presented extensive expert testimony along with Teva’s marketing efforts, catalogs, press releases, and testimony from Teva’s own witnesses, showing that Teva encouraged carvedilol sales for [congestive heart failure] despite its attempted carve-out.” *Id.* at 1335.

Similar to Hikma’s website, Appx820, Teva’s first press release described its medication as “the AB rated generic equivalent of [GSK]’s Coreg® Tablets” and as “indicated for *treatment of heart failure* and hypertension.” *Id.* at 1335-36 (internal quotations omitted) (emphasis in original). The Court noted how Teva’s press release used the term “heart failure” in a way that did “not parse between congestive heart failure,” i.e., the patented use, and the specific “post-MI LVD” indication that was not patented. *Id.* at 1336. “This [was] not an errant reference to ‘heart failure’;

it [was] Teva in a press release telling the world that its generic is a substitute for GSK's Coreg® tablets to treat congestive heart failure in the same manner as Coreg® (which is a method that infringed the '000 patent)." *Id.* And the Court noted Dr. McCullough's additional testimony that this press release "indicate[d] physicians should prescribe generic carvedilol for heart failure." *Id.* Teva's second press release "stated that it had received final approval to market its Generic version of [GSK]'s cardiovascular agent Coreg® (Carvedilol) Tablets." *Id.* (internal quotations omitted). Again, Dr. McCullough testified what this communicated to doctors by explaining how Teva's "use of 'cardiovascular agent' indicated to doctors they could use Teva's carvedilol 'for all indications,' including heart failure." *Id.* (internal quotations omitted).

Like the communications in *GlaxoSmithKline* of therapeutic equivalence in a general category of "heart failure" and as a general "cardiovascular agent" that encompassed the patented use to treat congestive heart failure, *id.* at 1335–36, Hikma communicated that its generic medication was equivalent for a general therapeutic category of "hypertriglyceridemia" and listed sales for the whole Vascepa® market to encompass the patented use to reduce cardiovascular risk in patients with HTG. *See* Appx529-533 (FAC, ¶¶ 111-116, 118-123, 125, 126); Appx612-613 (Hikma's press release on May 22, 2020); Appx708-710 (Hikma's press release on March 31, 2020); Appx711-713 (Hikma's press release on

September 3, 2020). Thus, this case is like *GlaxoSmithKline* in that inducement is shown by more than the label—except the district court erred here by preventing Amarin from developing the case and addressing the key factual dispute.

AstraZeneca is another case with parallels to the dispute here. This Court affirmed the district court’s grant of a preliminary injunction that prohibited Apotex’s launch of a generic version of AstraZeneca’s budesonide medication. 633 F.3d at 1045. The label instructed patients that it was “desirable to downward-titrate to the lowest effective dose once asthma stability is achieved.” *Id.* at 1057. AstraZeneca argued this “proposed label would induce consumers to infringe the asserted method claims because the label implicitly instructed users to administer the generic drug once daily,” which was covered by the patented method. *Id.* The district court agreed, but it also relied on a letter from the FDA discussing once-daily dosing that demonstrated how Apotex was both communicating about and aware of the infringement problem. *Id.* at 1057, 1059-60. Apotex’s awareness of the infringement problem and decision to distribute the generic with the problematic language in the label “formed the basis of the district court’s specific intent finding,” and the “district court correctly concluded” this was evidence of active steps to encourage infringement. *Id.* at 1059-60. Thus, *AstraZeneca* was another case where inducement, and specifically the necessary intent to induce, was supported by the label plus more. Like in *AstraZeneca*, Amarin alleged that the label itself

demonstrated Hikma’s intent to encourage physicians to treat patients with HTG while supporting those label allegations with additional allegations about Hikma’s conduct beyond its generic label.

Instead of those cases, the district court here relied on *Grunenthal*, while Hikma cited *HZNP Medicines LLC v. Actavis Laboratories UT, Inc.*, 940 F.3d 680 (Fed. Cir. 2019), and *Takeda*, which relied on *only* the label for inducement. As the magistrate judge explained, “unlike the allegations in this case, the evidence in those cases related solely to the effects of the generic labels.” Appx1427.

In *Grunenthal*, this Court affirmed a finding of no induced infringement following a bench trial. 919 F.3d at 1336. There, the brand medication had two indications, the off-patent treatment of “moderate to severe chronic pain in adults,” and the patented treatment of “neuropathic pain associated with diabetic peripheral neuropathy (DPN),” which was “a type of [polyneuropathic] pain.” *Id.* at 1338. Hikma and another generic manufacturer carved out treatment of DPN. *Id.* at 1339. The inducement inquiry turned on whether Hikma and the other generic had “the specific intent, based on the contents of their proposed labels, to encourage physicians to use their proposed ANDA products to treat polyneuropathic pain.” *Id.* The brand owners argued that severe chronic pain, which remained on the generic labels, was broad enough to include the specific patented treatment of polyneuropathic pain. *Id.* This Court disagreed because the off-patent treatment of

severe chronic pain more broadly included other categories: “[E]ven if severe chronic pain includes polyneuropathic pain, it also includes mononeuropathic pain and nociceptive pain.” *Id.*

Critically, the brand owners in *Grunenthal* “point[ed] only to the indications of the proposed labels as grounds for inducement.” *Id.* at 1340. And the label had effectively carved out the patented indication, which included removing reference to clinical studies for the patented indication: “[I]t is undisputed that neither of the accused ANDA labels list an indication for the management of pain associated with DPN. Nor do they mention any DPN clinical studies, which served as the basis for FDA approval of [the brand medication’s] indication for the treatment of neuropathic pain.” *Id.* at 1339-40. Thus, *Grunenthal* was a case where inducement was argued based on the label alone, and the generic labels had fully excised the patented indication. In contrast, this case involves allegations of actions beyond the label, as well as allegations that the label was not skinny enough.

In *HZNP*, this Court affirmed a grant of summary judgment finding no induced infringement for patents related to treating osteoarthritis. 940 F.3d at 683. The patented method in *HZNP* required three steps: “(1) apply the inventive formulation, (2) wait for the area to dry, and (3) apply sunscreen, insect repellent, or a second topical medication.” *Id.* at 702. But the generic label’s instructions only required the first application step with a warning that the area should be allowed to

dry before application of a sunscreen, etc. *Id.* That warning was not enough to show encouragement to infringe because the “evidence, viewed in the light most favorable to [the plaintiff], establishe[d] that some users *might* infringe.” *Id.* (emphasis added). The evidence did “not establish that ‘the proposed label instruct[ed] users to perform the patented method.’” *Id.* (quoting *AstraZeneca*, 633 F.3d at 1060). *HZNP* like *Grunenthal*, was thus a case that relied on the label alone.

In *Takeda*, Hikma sought approval for a generic medication for the prophylactic treatment of gout but not for the patented “treatment of acute gout flares.” 785 F.3d at 630. Hikma’s label stated the generic was “indicated for prophylaxis,” included a limitation of use that the “safety and effectiveness of [the generic] for acute treatment of gout flares during prophylaxis has not been studied,” and included a warning that “[i]f you have a gout flare while taking [the generic], tell your healthcare provider.” *Id.* (internal quotations omitted). Takeda argued that the warning would induce infringement because, for a patient using the generic for prophylaxis, “the physician would likely tell the patient to use the [generic] product to treat the acute flare.” *Id.* Although Takeda attempted to use evidence beyond the label to demonstrate that the warning would lead physicians to prescribe the generic for the infringing use of treating acute gout flares, the Court found none of the evidence supported inducement. *Id.* at 632.

“[E]ven if we do look outside the label, there is no evidence that the label would necessarily lead doctors who are consulted by patients taking [the generic] to prescribe an off-label use of it to treat acute gout flares.” *Id.* The additional evidence did not show additional encouragement or promotion. Instead, the evidence was meant to show how the warning would “inevitably” lead doctors to prescribe the generic for off-label infringing uses, but the evidence failed to show even that. *Id.* *Takeda* was a case that relied on the label alone to show encouragement, and at that, it relied on a warning to see a doctor as the only encouragement to induce infringement. Unlike *Takeda*, this case relies on communications beyond the label.

Like *GlaxoSmithKline* and *AstraZeneca*, this is a case where a label plus more demonstrates inducement. It is not a case like *Grunenthal*, *HZNP*, or *Takeda*, where the generic label alone was used to show an action intended to induce. The district court erred when it relied on *Grunenthal* and distinguished *GlaxoSmithKline*.

B. Previous cases consistently reached a later stage than a motion to dismiss because factual issues are central to inducement

Every leading skinny-label case discussed here and at the district court reached a more advanced posture than a motion to dismiss. *See* Appx1427 (R&R, 15) (magistrate judge noting how “none of those cases [relied upon by Hikma] was resolved at the motion to dismiss stage.”). That makes sense. Infringement is a “quintessential fact question,” *GlaxoSmithKline*, 7 F.4th at 1328, and induced infringement hinges on factual determinations.

The leading precedent from this Court highlights that the motion to dismiss was an improper vehicle for resolving the key issues in this case—none of the significant cases were appealed from a dismissal on the pleadings. *See GlaxoSmithKline*, 7 F.4th at 1323 (reversing judgment as a matter of law after a jury trial finding infringement); *AstraZeneca*, 633 F.3d at 1045 (affirming grant of preliminary injunction finding likelihood of showing induced infringement); *Grunenthal*, 919 F.3d at 1338 (affirming bench trial finding no infringement); *HZNP*, 940 F.3d at 682, 686 (affirming grant of summary judgment for non-infringement); *Takeda*, 785 F.3d at 627 (affirming denial of preliminary injunction finding no likelihood of showing induced infringement).

To be sure, cases at the preliminary injunction stage are early-stage cases. But a preliminary injunction is a discretionary grant that requires likelihood of success on the merits, *see, e.g., AstraZeneca*, 633 F.3d at 1049, whereas the grant of a motion to dismiss requires the absence of even a plausible claim for relief.

Before the district court, Hikma asserted that there were “many decisions” “even at the pleadings stage” where there was no induced infringement as a matter of law. Appx481-482 n.2 (Hikma’s brief in support of its motion to dismiss). Of the ten cases Hikma cited in support, it only described three as involving a dismissal on the pleadings. *Id.* Each of those cases is distinguishable. *Bayer Schering Pharma AG v. Lupin, Ltd.*, was not a skinny-label case—the generic and brand labels were

identical—nor did it involve inducement of an FDA approved use. 676 F.3d 1316, 1321 (Fed. Cir. 2012). *AstraZeneca Pharmaceuticals LP v. Apotex Corp.*, turned on the question of whether infringement under 35 U.S.C. § 271(e)(2)(A) could be based on an ANDA seeking to market a drug that was not patented and for a use that was not patented. 669 F.3d 1370, 1378-79 (Fed. Cir. 2012). And while *Takeda Pharmaceuticals U.S.A., Inc. v. West-Ward Pharmaceutical Corp.*, did involve a dismissal on the pleadings of an inducement claim, that was based on the district court’s conclusion that Takeda’s allegations were “too conclusory to pass muster.” 188 F. Supp. 3d 367, 377 (D. Del. 2016). Significantly, the *Takeda* district court vacated that very dismissal in light of the low pleading standard after Takeda amended its complaint with allegations about specific communications that Takeda argued amounted to active encouragement. *Takeda Pharms. U.S.C., Inc. v. West-Ward Pharm. Corp.*, 2016 WL 723054 at *2 (D. Del. Dec. 14, 2016). Amarin’s specific allegations regarding Hikma’s website and press releases position this case squarely within the analysis of that later vacatur of the *Takeda* dismissal.

The fact that all the leading skinny-label cases were more advanced than a motion to dismiss confirms that the district erred when it granted the motion to dismiss.

C. This case involves a patented use that is broader than the off-patent use—unlike in *Grunenthal*

Amarin’s patented use to reduce cardiovascular risk in patients with HTG covers a broader set of patients than the off-patent use to treat patients with severe HTG. In other words, more patients suffer from elevated triglyceride levels above 150 mg/dL, i.e., patients with HTG, than the subset with levels above 500 mg/dL, i.e., patients with *severe* HTG. The district court analogized to *Grunenthal* and got this point exactly backwards. Vascepa[®]’s additional patented indication is not simply a subset of its original indication—it is a different use of the drug to treat a broader patient population for which there was a different clinical concern.

Describing the distinct situation in *Grunenthal*, the district court here explained how “a label indicated for ‘[m]oderate to severe chronic pain,’ which included both infringing and non-infringing uses, did ‘not specifically encourage use’ of the generic for the patented treatment.” Appx9 (quoting *Grunenthal*, 919 F.3d at 1339) (modification in original). And the district court concluded, “[t]his case is more like *Grunenthal*, where the broader category simply includes both infringing and non-infringing uses, without ‘specifically encourage[ing]’ the use of the generic for the non-infringing uses.” Appx9 (quoting *Grunenthal*, 919 F.3d at 1339) (modification in original). But the “broader category” in *Grunenthal* was the off-patent treatment of “moderate to severe chronic pain in adults.” *Grunenthal*, 919 F.3d at 1338. That method was directed to a broader group of patients than those

with “neuropathic pain associated with diabetic peripheral neuropathy (DPN),” which was “a type of polyneuropathic pain.” *Id.* This Court rejected the argument that the generic label for the broader, off-patent use induced the specific, patented use because the off-patented treatment of severe chronic pain more broadly included other categories: “[E]ven if severe chronic pain includes polyneuropathic pain, it also includes mononeuropathic pain and nociceptive pain.” *Id.* In sum, *Grunenthal* was a case where encouraging treatment of patients with moderate to severe pain was not enough to induce the specific and patented treatment of polyneuropathic pain.

In contrast, this case involves a broader patented use that covers a larger patient population with HTG generally (triglycerides above 150 mg/dL) and an off-patent use that covers a narrower patient population with *severe* HTG (triglycerides above 500 mg/dL). Both *Grunenthal* and this case involve generic labels that pointed at a broader category of patients. The difference is that, in *Grunenthal*, that broader category of patients, i.e., those suffering from severe chronic pain, was tied to the off-patent indication. *Id.* Whereas here, the broader category of patients that Hikma’s press releases and website identify, i.e., those with HTG generally, is associated with the patented use.

That distinction, misapplied by the district court, changes the factual question of what Hikma’s conduct would have communicated to prescribing physicians. This

case is more like *GlaxoSmithKline*, where communications that the generic medication was therapeutically equivalent for a broad category of “heart failure” and as a general “cardiovascular agent” associated the generic with the broader patented use to treat congestive heart failure. 7 F.4th at 1335-36.

CONCLUSION

Amarin stated a claim for relief that was at least plausible. The order dismissing Amarin’s first amended complaint should be reversed, and the case should be remanded for further proceedings.

Respectfully submitted,

PERKINS COIE LLP

by /s/Nathan K. Kelley

Nathan K. Kelley

Counsel for Appellants Amarin Pharma, Inc., Amarin Pharmaceuticals Ireland Ltd., and Mochida Pharmaceutical Co., Ltd.

ADDENDA

1. Memorandum Opinion, Order, and Final Judgment (Appx1-15)
2. Magistrate's Report and Recommendation (Appx1413-1430)
3. U.S. Patent No. 9,700,537 (Appx36-47)
4. U.S. Patent No. 10,568,861 (Appx77-129)

IN THE UNITED STATES DISTRICT COURT
FOR THE DISTRICT OF DELAWARE

AMARIN PHARMA, INC., AMARIN
PHARMACEUTICALS IRELAND
LIMITED, MOCHIDA PHARMACEUTICAL
CO., LTD.

Plaintiffs;

v.

HIKMA PHARMACEUTICALS USA INC.,
HIKMA PHARMACEUTICALS PLC, AND
HEALTH NET, LLC,

Defendants.

Civil Action No. 20-1630-RGA-JLH

MEMORANDUM OPINION

Jeremy D. Anderson, FISH & RICHARDSON P.C., Wilmington, DE, Elizabeth M. Flanagan,
Michael Kane (argued), Deanna J. Reichel, FISH & RICHARDSON P.C., Minneapolis, MN;
Jonathan E. Singer, FISH & RICHARDSON P.C., San Diego, CA;

Attorneys for Plaintiffs.

Dominick T. Gattuso, HEYMAN ENERIO GATTUSO & HIRZEL LLP, Wilmington, DE;
Charles B. Klein (argued), Claire A. Fundakowski, WINSTON & STRAWN LLP, Washington,
DC; Eimeric Reig-Plessis, WINSTON & STRAWN LLP, San Francisco, CA; Alison M. King,
WINSTON & STRAWN LLP, Chicago, IL;

Attorneys for Defendants Hikma Pharmaceuticals USA Inc. and Hikma Pharmaceuticals
PLC.

John C. Phillips, Jr., David A. Bilson, PHILLIPS MCLAUGHLIN & HALL, P.A., Wilmington,
DE; Don J. Mizerk (argued), HUSCH BLACKWELL LLP, Chicago, IL; Dustin L. Taylor,
HUSCH BLACKWELL LLP, Denver, CO;

Attorneys for Defendant Health Net.

January 4, 2022


 ANDREWS, U.S. DISTRICT JUDGE:

I referred this very interesting case to a magistrate judge. (D.I. 16). She wrote a Report and Recommendation on three pending motions to dismiss. (D.I. 64). Defendants filed objections (D.I. 70, 71), to which Plaintiffs responded (D.I. 77, 78). There is even an amicus brief. (D.I. 75). I heard oral argument on October 14, 2021. For the following reasons, I will ADOPT-IN-PART the Report and Recommendation. (D.I. 64). Hikma's motion to dismiss the First Amended Complaint (D.I. 19) is GRANTED. Hikma's motion to dismiss the original complaint (D.I. 11) is DISMISSED AS MOOT. Health Net's motion to dismiss the First Amended Complaint (D.I. 30) is DENIED.

I. BACKGROUND

Plaintiffs sued Defendants for induced infringement of three patents that describe methods of using icosapent ethyl for the reduction of cardiovascular risk. (D.I. 17). Plaintiffs manufacture and sell VASCEPA, a branded version of icosapent ethyl. (*Id.* at ¶¶ 1, 57-58). Defendant Hikma is a generic manufacturer of icosapent ethyl. (*Id.* at ¶ 1). Defendant Health Net is an insurer that provides coverage for Vascepa and Hikma's generic version. (*Id.* at ¶¶ 139-40).

II. LEGAL STANDARD

A motion to dismiss for failure to state a claim upon which relief may be granted is considered a dispositive motion. D. Del. LR 72.1(a)(3). A magistrate judge's Report and Recommendation regarding a case-dispositive motion is reviewed *de novo*. Fed. R. Civ. P. 72(b)(3).

When reviewing a motion to dismiss pursuant to Federal Rule of Civil Procedure 12(b)(6), the Court must accept the complaint's factual allegations as true. *See Bell Atl. Corp. v.*

Twombly, 550 U.S. 544, 555–56 (2007). Rule 8(a) requires “a short and plain statement of the claim showing that the pleader is entitled to relief.” *Id.* at 555. The factual allegations do not have to be detailed, but they must provide more than labels, conclusions, or a “formulaic recitation” of the claim elements. *Id.* (“Factual allegations must be enough to raise a right to relief above the speculative level . . . on the assumption that all the allegations in the complaint are true (even if doubtful in fact).”). Moreover, there must be sufficient factual matter to state a facially plausible claim to relief. *Ashcroft v. Iqbal*, 556 U.S. 662, 678 (2009). The facial plausibility standard is satisfied when the complaint’s factual content “allows the court to draw the reasonable inference that the defendant is liable for the misconduct alleged.” *Id.* (“Where a complaint pleads facts that are merely consistent with a defendant’s liability, it stops short of the line between possibility and plausibility of entitlement to relief.” (internal quotation marks omitted)).

Section 271(b) provides, “whoever actively induces infringement of a patent shall be liable as an infringer.” 35. U.S.C. 271(b). To state a claim for induced infringement, the complaint must allege that there has been direct infringement, that the defendant knowingly induced infringement, and that the defendant has the intent to encourage another’s infringement. *MEMC Elec. Materials, Inc. v. Mitsubishi Materials Silicon Corp.*, 420 F.3d 1369, 1378 (Fed. Cir. 2005). A generic manufacturer can be liable for inducing infringement of a patented method even when the generic has attempted to “carve out” the patented indications. *GlaxoSmithKline LLC v. Teva Pharmaceuticals USA, Inc.*, 7 F.4th 1320, 1338 (Fed. Cir. 2021) (per curiam).

III. HIKMA'S MOTION TO DISMISS

A. BACKGROUND

Amarin sells Vascepa (icosapent ethyl) for the treatment of severe hypertriglyceridemia (the “SH indication”) and cardiovascular risk reduction (the “CV indication”). (D.I. 17 at ¶¶ 1, 56). Only the CV indication is covered by Plaintiffs’ patents. (*See* D.I. 22 at 1). Hikma received FDA approval to sell a generic version for the SH indication under the “skinny label” or “section viii carveout” regime. (D.I. 17 at ¶¶ 11, 95, 108). This regime allows a generic to sidestep the typical FDA requirement that a generic’s labeling is the same as the brand’s labeling. 21 U.S.C. §§ 355(j)(2)(A)(viii). The generic does so by removing the portions of the label associated with the patented use, resulting in a “skinny label.” Plaintiffs allege that Defendant Hikma’s label is “not-skinny-enough” and that the label, along with Hikma’s public statements, induce infringement of Plaintiffs’ patents for the CV indication. (D.I. 22 at 1).

B. DISCUSSION

1. The Federal Circuit’s *GSK* Decision

Two days after the Report issued, the Court of Appeals issued the most recent authoritative opinion concerning skinny labels, albeit after the case was fully litigated in the district court. *See GlaxoSmithKline LLC v. Teva Pharmaceuticals USA, Inc.* [hereinafter “*GSK*”], 7 F.4th 1320 (Fed. Cir. 2021). The Federal Circuit affirmed a jury’s findings that Teva’s “partial label” induced infringement of GSK’s patent, notwithstanding Teva’s attempt to exclude the patented use from its label under the skinny label regime. (*Id.* at 1338). Ultimately, the Federal Circuit concluded, “Teva’s partial label did not successfully carve out the patented use, and thus, Teva was selling its generic with a label which infringed the method claim.” *Id.* Accordingly, Teva’s label was “not a skinny label.” *Id.* at 1328.

The Federal Circuit also found that two Teva press releases supported the jury’s verdict. *Id.* at 1335-37. The first press release advertised Teva’s drug as “indicated for treatment of heart failure” and did “not parse between congestive heart failure [the patented indication] or post-MI LVD [an unpatented indication].” *Id.* at 1336. The second press release stated that Teva received approval to market “its Generic version of GlaxoSmithKline’s cardiovascular agent Coreg.” *Id.* Expert testimony established that the phrase “‘cardiovascular agent’ ‘indicated to doctors they could use Teva’s carvedilol ‘for all indications,’ including heart failure.” *Id.*

The Court held that *GSK* is a “narrow, case-specific review” and that it is still the law that “generics could *not* be held liable for merely marketing and selling under a ‘skinny’ label omitting all patented indications, or for merely noting (without mentioning any infringing uses) that FDA had rated a product as therapeutically equivalent to a brand-name drug.” *Id.* at 1326. An “AB rating,” as the complaint explains, “reflects a decision [by the FDA] that a generic drug is therapeutically equivalent to a branded drug when the generic drug is used as labeled[.]” (D.I. 17 at ¶ 98). As *GSK*’s discussion of Teva’s press releases illustrates, where a generic label does not effectively carve out a patented use, advertisement that the drug is “AB rated” can support a finding of inducement. *GSK*, 7 F.4th at 1335.

2. Amarin’s Complaint

Amarin’s complaint pleads several factual allegations in support of its claim that Hikma induces infringement. These allegations fall into two categories: Hikma’s label and Hikma’s public statements. The Magistrate Judge recommends I deny Hikma’s motion to dismiss because “several . . . portions of Hikma’s label, taken together with Hikma’s public statements, instruct physicians to use Hikma’s product in a way that infringes the asserted patents.” (D.I. 64 at 12).

The bulk of the briefing and oral argument was directed to Hikma's label, and I will address those arguments first.

As to the label, Hikma objects that Amarin's complaint fails to plead instruction as to at least two claim limitations—the requirement that icosapent ethyl be administered to reduce CV risk and the requirement to co-administer with a statin. (D.I. 71 at 7-8). Because I agree with Hikma that there has been no instruction as to CV risk reduction, I will not address Hikma's argument regarding co-administration with a statin.

Amarin contends that Hikma's label teaches CV risk reduction for two reasons. First, Hikma's label contains a notice regarding side effects for patients with CV disease. (D.I. 78 at 5-6). Second, Hikma's label does not “state that Hikma's ‘generic version’ of VASCEPA should not be used for the CV Indication or that the effect of icosapent ethyl on cardiovascular mortality and morbidity in patients with severe hypertriglyceridemia has not been determined” (the “CV limitation”). (D.I. 17 at ¶¶ 108, 121). Hikma responds that (1) the notice regarding side effects is a warning and thus not an instruction to use icosapent ethyl to reduce cardiovascular risk, and (2) the removal of the CV risk reduction limitation is mere silence and that Hikma has no duty to discourage infringing use.

Regarding the warning as to side effects, I agree with Hikma. The label states, “Icosapent ethyl may cause serious side effects, including: ... Heart rhythm problems which can be serious and cause hospitalization have happened in people who take icosapent ethyl, especially in people who have heart (cardiovascular) disease or diabetes with a risk factor for heart (cardiovascular) disease[.]” (D.I. 17, Ex. K at 12-13 of 15). This is hardly instruction or encouragement. *See, e.g., Otsuka Pharm. Co. v. Torrent Pharms. Ltd.*, 99 F. Supp. 3d 461, 490 (D.N.J. 2015) (“[A] warning is just that—a warning. It is not an instruction[.]”).

Amarin also argues that Hikma “removed”¹ the CV limitation from its label, which would be “understood in the field to teach that Hikma’s product *has* been proven to reduce CV risk and to encourage its use for that purpose” because other drugs in the same class have not been shown to reduce CV risk. (D.I. 78 at 4). This amounts to an “affirmative statement” that it can be used for cardiovascular risk reduction, according to Plaintiffs. (D.I. 85 at 62:16-62:5).

The Federal Circuit has previously rejected the argument that generic labels must contain a “clear statement” discouraging use of the patented indication. *Takeda Pharms. U.S.A., Inc. v. W.-Ward Pharm. Corp.*, 785 F.3d 625, 632 n.4 (Fed. Cir. 2015). Plaintiffs must plead that “Hikma took affirmative steps to induce, not affirmative steps to make sure others avoid infringement.” *Id.* Even if Plaintiffs are right that Hikma’s label’s silence regarding CV risk reduction communicates to the public that icosapent ethyl can be used to reduce CV risk, “merely describing an infringing mode is not the same as recommending, encouraging, or promoting an infringing use.” *Id.* at 631 (cleaned up). I therefore find that the lack of a CV limitation on Hikma’s label does not plausibly teach CV risk reduction.

Since I find that the label does not instruct CV risk reduction, the question is whether Hikma’s public statements, including press releases and Hikma’s website, induce infringement. (D.I. 17 at ¶ 127). Hikma’s press releases state that its product is the “generic equivalent to Vascepa®” and that “Vascepa is a prescription medicine that is indicated, *in part*, as an adjunct to diet to reduce triglyceride levels in adult patients with severe (≥ 500 mg/dL)

¹ Hikma contests Plaintiffs’ use of the word “removal,” noting, “*Amarin* removed the limitation of use from Vascepa’s label *before* Hikma launched its product, and Hikma was required to use ‘the same [labeling] as the labeling approved for the listed drug.’” (D.I. 71 at 7 n.2 (citing 21 U.S.C. § 355(j)(2)(A)(v))). The facts pled in the complaint state that the removal happened during the FDA approval process. (D.I. 17 at ¶ 108). At any rate, it appears that there is no allegation that Hikma’s product was ever marketed with a label containing the CV limitation.

hypertriglyceridemia. According to IQVIA, US sales of Vascepa® were approximately \$919 million in the 12 months ending February 2020.” (*Id.* at ¶ 112). The sales figures cited by Hikma include Vascepa’s sales of the patented indication. The complaint further alleges that Hikma’s website states that Hikma’s generic is “AB rated” in the “Therapeutic Category: Hypertriglyceridemia.” (*Id.* at ¶ 125).

Hikma’s press releases might be relevant to intent but they do not support actual inducement. Hikma’s advertising of icosapent ethyl as the “generic equivalent” of Vascepa does not expose Hikma to liability. *GSK*, 7 F.4th at 1335 n.7. The citation of Vascepa’s sales figures go to Hikma’s intent to induce. Intent alone is not enough; Amarin must plead an inducing act.

Amarin also alleges that Hikma’s website induces infringement by advertising its product in the therapeutic category “hypertriglyceridemia.” The complaint pleads, “hypertriglyceridemia . . . does not match and is broader than the Indications and Usage sections of Hikma’s Label, which includes only Severe Hypertriglyceridemia Indication (i.e., triglycerides ≥ 500 mg/dL).” (D.I. 17 at ¶ 126). Accepting the facts in the light most favorable to Amarin, Amarin has pled that the category “hypertriglyceridemia” includes infringing uses. The question is whether this is enough, without a label or other public statements instructing as to infringing use, to induce infringement.

I hold that it is not. This statement does not rise to the level of encouraging, recommending, or promoting taking Hikma’s generic for the reduction of CV risk.

Two recent Federal Circuit cases are instructive on this point. The *GSK* majority found that Teva’s advertising of “its Generic version of GlaxoSmithKline’s cardiovascular agent,” when “cardiovascular agent” was a category that included both infringing and non-infringing uses, supported a jury’s finding of inducement. 7 F.4th at 1336. The Court emphasized that:

Teva did not merely say its drug is a cardiovascular agent, leaving the world to wonder about its uses. It said its product is a generic equivalent of GSK's cardiovascular agent Coreg®. It was reasonable for the jury to conclude, especially in light of the prior press release that expressly mentioned heart failure, that Teva was again encouraging the substitution of its product for all of Coreg's® cardiovascular indications, including as claimed in the '000 patent.

Id. at 1337. In contrast, the Federal Circuit has found that a label indicated for “[m]oderate to severe chronic pain,” which included both infringing and non-infringing uses, did “not specifically encourage use” of the generic for the patented treatment. *Grunenthal GMBH v. Alkem Lab'ys Ltd.*, 919 F.3d 1333, 1339 (Fed. Cir. 2019) (“[E]ven if severe chronic pain includes polyneuropathic pain, it also includes mononeuropathic pain and nociceptive pain. Therefore, the proposed ANDA labels do not specifically encourage use of tapentadol hydrochloride for treatment of polyneuropathic pain.”).

Here, Hikma stated that its product was “AB Rated” in a category that includes both patented and non-patented uses. The “AB rating” points to the label, as the *GSK* court explained:

We do not hold that an AB rating in a true section viii carve-out (one in which a label was produced that had no infringing indications) would be evidence of inducement. In this case, Teva's representation of AB rating would point physicians to its partial label, which, for the reasons above, the jury was free to credit as evidence of induced infringement.

GSK, 7 F.4th at 1335 n.7. Unlike Teva's press release in *GSK*, Hikma has not pointed to Vascepa's patented uses in describing itself as Vascepa's generic equivalent. This case is more like *Grunenthal*, where the broader category simply includes both infringing and non-infringing uses, without “specifically encourage[ing]” the use of the generic for the non-infringing uses. 919 F.3d at 1339.

Since I find that Amarin's complaint has failed to plead inducement based on Hikma's label or public statements, I will grant Hikma's motion to dismiss.

IV. HEALTH NET'S MOTION TO DISMISS

A. BACKGROUND

Defendant Health Net provides insurance coverage for Plaintiffs' branded Vascepa and Defendant Hikma's generic version. According to Plaintiffs, Health Net's formulary placement induces infringement of Plaintiffs' patents by encouraging the use of Hikma's generic for the CV indication. Health Net's formulary lists Hikma's generic in a lower tier than Amarin's Vascepa, resulting in lower copays when a patient opts for Hikma's generic. (D.I. 17 at ¶ 143). Since it is common for pharmacies to automatically substitute an AB-rated generic such as Hikma's for the branded version, Plaintiffs allege that this formulary placement leads to substitution on "all VESCEPA prescriptions, not just the prescriptions directed to the" SH indication. (*Id.* at ¶ 151).

B. DISCUSSION

The Report recommends I deny Health Net's motion to dismiss because there are factual questions regarding whether Health Net has taken an affirmative act to induce infringement and whether Health Net's actions actually cause others to infringe. (D.I. 64 at 17). Health Net objects, "Plaintiffs fail to allege facts (not conclusions or speculation) supporting a plausible conclusion that Health Net was aware of the asserted patents, and once aware, took affirmative steps with the specific intent to induce another's infringement of those patents—rather than merely acting despite knowledge that others may infringe." (D.I. 70 at 2). I disagree.

I find that the complaint pleads enough facts to plausibly allege knowledge of the asserted patents. Amarin sent a pre-suit letter to its point of contact for Health Net. (D.I. 17 at ¶ 87). It is true that the pre-suit letter did not specify the patent numbers. However, the letter states that Amarin has patent exclusivity for the CV indication, and the complaint elsewhere pleads that the patents associated with the CV indication are readily available through a resource

well-known in the industry, the FDA's Orange Book. (*Id.* at ¶¶ 84, 88). Thus, I agree with the Magistrate Judge that these facts, taken together in the light most favorable to the Plaintiffs, make it plausible that Health Net had specific knowledge of the patents at issue.

Read in the light most favorable to Amarin, the complaint also plausibly alleges affirmative acts taken with a specific intent to induce another's infringement. Formulary selection and the prior authorization process, as pled, could be affirmative acts under the law of induced infringement. Health Net argues that the selection of its formulary is automatic, based on Plaintiff's own pricing as compared to the generic. (D.I. 85 at 75:5-12 (noting that "this is done by a computer program")). This may be true, but it is not a shield. Health Net added generic icosapent ethyl capsules to its formularies. (D.I. 17 at ¶¶ 140-143). It is immaterial whether the placement was done by a human or a computer.

Amarin also plausibly pleads specific intent to induce. At the very least, Health Net's prior authorization form supports an inference of specific intent because it lists the patented indication on the generic icosapent ethyl capsules form. (D.I. 17 at ¶ 159). Health Net's placement of generic icosapent ethyl on a preferred tier encourages the substitution of the generic for the branded drug, including for the patented indication. (*Id.* at ¶¶ 145, 151). Together, this is enough to plead specific intent to induce.

In its objections, Health Net argues that the "preferred" language in its formularies cannot be an active step because they are required by state law to disclose which drugs are "preferred." (*Id.* at 5). This may be true, but it is not the language of the formulary that is at issue; it is the incentives the formulary puts in place. (*See id.* at ¶¶ 145, 151).

Health Net stresses that they are just a payer, not the physician writing the prescription nor the pharmacist making the substitution. (D.I. 70 at 9). As the Report points out, "It may ...

turn out that, despite knowledge of infringement by its beneficiaries and their providers, Health Net's actions in selecting its formulary and adopting its prior authorization procedure ... do not, in fact, influence the decisions of beneficiaries, pharmacists, and medical providers to use, dispense, and prescribe Hikma's generic product in an infringing way[.]” (D.I. 64 at 17; *see Warner-Lambert Co. v. Apotex Corp.*, 316 F.3d 1348, 1364 (Fed. Cir. 2003) (“[I]f a physician, without inducement by Apotex, prescribes a use of gabapentin in an infringing manner, Apotex's knowledge is legally irrelevant. In the absence of any evidence that Apotex has or will promote or encourage doctors to infringe the neurodegenerative method patent, there has been raised no genuine issue of material fact.”)). These are factual questions that cannot be resolved on a motion to dismiss.

Ultimately, I agree with the Magistrate Judge that Plaintiffs have pled enough to proceed with their case against Health Net.

V. CONCLUSION

An appropriate order will follow.

IN THE UNITED STATES DISTRICT COURT
FOR THE DISTRICT OF DELAWARE

AMARIN PHARMA, INC., AMARIN
PHARMACEUTICALS IRELAND
LIMITED, MOCHIDA PHARMACEUTICAL
CO., LTD.

Plaintiffs;

v.

HIKMA PHARMACEUTICALS USA INC.,
HIKMA PHARMACEUTICALS PLC, AND
HEALTH NET, LLC,


Defendants.

Civil Action No. 20-1630-RGA-JLH

ORDER

For the reasons stated in the accompanying Memorandum Opinion, Hikma's motion to dismiss the First Amended Complaint (D.I. 19) is **GRANTED**. Hikma's motion to dismiss the original complaint (D.I. 11) is **DISMISSED AS MOOT**. Health Net's motion to dismiss the First Amended Complaint (D.I. 30) is **DENIED**. The first amended complaint (D.I. 17) as to the Hikma Defendants is **DISMISSED** without prejudice.¹

IT IS SO ORDERED this 4th day of January 2022.


United States District Judge

¹ Plaintiffs requested leave to amend if the first amended complaint was dismissed. (D.I. 22 at 20). Plaintiffs gave no indication as to what more they could plead, but if they have something more, they may file a motion in compliance with the Local Rules seeking leave to amend.

IN THE UNITED STATES DISTRICT COURT
FOR THE DISTRICT OF DELAWARE

AMARIN PHARMA, INC., AMARIN
PHARMACEUTICALS IRELAND
LIMITED, MOCHIDA
PHARMACEUTICAL CO., LTD.,

Plaintiffs,

v.

C.A. No. 20-1630-RGA-JLH

HIKMA PHARMACEUTICALS USA
INC., HIKMA PHARMACEUTICALS
PLC, AND HEALTH NET, LLC

Defendants.

**FINAL JUDGMENT UNDER
FEDERAL RULE OF CIVIL PROCEDURE 54(b)**

THIS MATTER having come before the Court on Defendants Hikma Pharmaceuticals USA Inc. and Hikma Pharmaceuticals PLC's (collectively, "Hikma") Motion for Entry of Final and Appealable Judgment under Federal Rule of Civil Procedure 54(b), and the Court having considered Hikma's arguments and submissions in support of the Motion;

It is hereby **ORDERED** that the Motion is **GRANTED**.


For the reasons set forth by Hikma in its moving papers, the Court finds that the Court's order granting Hikma's motion to dismiss Plaintiffs' first amended complaint (D.I. 98) is a final judgment resolving Plaintiffs' claims against Hikma, and the Court expressly determines that there is no just reason for delay (*see* Fed. R. Civ. P. 54(b)).

Considering the factors set forth in *Berkeley Inv. Grp., Ltd. v. Colkitt*, 455 F.3d 195, 203 (3d Cir. 2006), the Court finds that (1) the relationship between the adjudicated claims against Hikma and the unadjudicated claims against the remaining Defendant, Health Net, LLC, is

minimal because Plaintiff's theories of infringement against these respective defendants are materially different; (2) the only foreseeable possibility that the need for review might be mooted by future developments in this Court is the invalidation of the asserted patents, which is unlikely to occur for more than a year; (3) the possibility that the reviewing court might be obliged to consider the same issue a second time is minimal because any appeal of the Court's order granting Hikma's motion to dismiss does not relate to Plaintiffs' infringement theory against Health Net; (4) there is no claim or counterclaim which could result in a set-off against the judgment sought to be made final; and (5) no miscellaneous factors (such as delay, economic and solvency considerations, shortening the time of trial, frivolity of competing claims, expense, and the like) weigh against entering final judgment at this time.

Accordingly, final judgment is **ENTERED** in favor of Hikma and against Plaintiffs; Plaintiffs' claims against Hikma in this action are **DISMISSED WITH PREJUDICE**; and each party shall bear its own costs and fees.

IT IS SO ORDERED this 13th day of October, 2022.


Honorable Richard G. Andrews
United States District Court Judge

**IN THE UNITED STATES DISTRICT COURT
FOR THE DISTRICT OF DELAWARE**

AMARIN PHARMA, INC., AMARIN)	
PHARMACEUTICALS IRELAND)	
LIMITED, MOCHIDA)	
PHARMACEUTICAL CO., LTD.,)	
)	
Plaintiffs,)	
)	
v.)	C.A. No. 20-1630-RGA-JLH
)	
HIKMA PHARMACEUTICALS USA INC.,)	
HIKMA PHARMACEUTICALS PLC, AND)	
HEALTH NET, LLC,)	
)	
Defendants.)	
)	

REPORT AND RECOMMENDATION

Plaintiffs Amarin Pharma, Inc., Amarin Pharmaceuticals Ireland Limited (collectively, “Amarin”), and Mochida Pharmaceutical Co., Ltd. (“Mochida”) filed this suit against Defendants Hikma Pharmaceuticals USA Inc., Hikma Pharmaceuticals PLC (collectively, “Hikma”), and Health Net, LLC (“Health Net”). Plaintiffs allege that Hikma and Health Net have each induced infringement of U.S. Patent Nos. 9,700,537 (the ’537 patent), 8,642,077 (the ’077 patent), and 10,568,861 (the ’861 patent) under 35 U.S.C. § 271(b). Hikma and Health Net have separately moved to dismiss under Federal Rule of Civil Procedure 12(b)(6).

Plaintiffs’ infringement case against Hikma is what is referred to by those in the know as a “skinny label” case. Amarin developed and markets a branded prescription drug that has two FDA-approved indications. One of those indications is patented, the other is not. Hikma launched a generic version after receiving FDA approval for the non-patented indication only. Notwithstanding the limited approval, Plaintiffs allege that Hikma—through its product label,

website, and press releases—instructs and encourages physicians to use its generic version for the patented indication, making Hikma liable for inducing infringement under 35 U.S.C. § 271(b).

Plaintiffs have an entirely different (and apparently novel) theory as to Health Net. Health Net is a health insurance provider. It does not prescribe drugs, but it does pay for drugs that are prescribed to its beneficiaries by physicians. Plaintiffs allege that the way that Health Net has set up its approval and payment process for Amarin’s product and Hikma’s generic version amounts to active encouragement to use Hikma’s generic version for the patented indication, making Health Net liable for inducing infringement under 35 U.S.C. § 271(b).

This case is at the pleadings stage. I cannot make factual findings about what Hikma’s label and advertisements communicate to physicians. Nor is it appropriate at this stage to make findings about how Health Net’s prescription drug coverage operates and whether it actually has any effect on anyone’s decision to use Hikma’s product for the patented use. The only determination at this stage is whether Plaintiffs’ allegations state plausible claims for relief.

“The plausibility standard is not akin to a ‘probability requirement,’”¹ and “a well-pleaded complaint may proceed even if it strikes a savvy judge that actual proof of the facts alleged is improbable, and that a recovery is very remote and unlikely.”² I conclude that Plaintiffs’ claims satisfy the plausibility standard. Accordingly, I recommend that both motions to dismiss be DENIED.

¹ *Ashcroft v. Iqbal*, 556 U.S. 662, 678 (2009) (quoting *Bell Atl. Corp. v. Twombly*, 550 U.S. 544, 556 (2007)).

² *Twombly*, 550 U.S. at 556 (2007) (internal marks omitted).

I. BACKGROUND

The statutory scheme for obtaining FDA approval of a generic drug for only non-patented uses has been well explained in numerous cases and I could do no better here.³ Accordingly, this Report and Recommendation assumes familiarity with the key features of the Hatch-Waxman generic drug approval process as it relates to “carve out” labels (aka “skinny” labels) and associated infringement litigation.

A. Amarin’s VASCEPA®⁴

The active ingredient in Amarin’s Vascepa product is icosapent ethyl, an ethyl ester of an omega-3 fatty acid (EPA) commonly found in fish oils. (D.I. 17 ¶¶ 25, 28, 54, Ex. D.) Vascepa currently has two FDA-approved indications: (1) treatment of severe hypertriglyceridemia (the “SH indication”); and (2) cardiovascular risk reduction (the “CV indication”). (*Id.* ¶¶ 1, 56.)

Severe hypertriglyceridemia (SH) is a condition where patients have triglyceride levels greater than 500 mg/dL. (*Id.* ¶ 30, Ex. D.) Vascepa received FDA approval for the SH indication in 2012. (*Id.* ¶ 30.) At that time, and up until 2019, the Vascepa label contained the following “limitation of use” regarding the CV indication: “The effect of VASCEPA on cardiovascular mortality and morbidity in patients with severe hypertriglyceridemia has not been determined.” (*Id.* ¶ 60, Exs. E, F.)

After receiving FDA approval to market Vascepa for the SH indication, Amarin conducted further clinical studies to examine the effects of Vascepa on cardiovascular risk reduction. (*Id.*

³ See, e.g., *AstraZeneca LP v. Apotex, Inc.*, 633 F.3d 1042, 1045-46 (Fed. Cir. 2010) (describing Hatch-Waxman scheme and carve out labels); *GlaxoSmithKline LLC v. Teva Pharms. USA, Inc.*, No. 14-878-LPS-CJB, 2016 WL 3946770, at *2-3 (D. Del. July 20, 2016) (same), *report and recommendation adopted*, No. 14-878-LPS-CJB, 2017 WL 1050574 (D. Del. Mar. 20, 2017).

⁴ I assume the facts alleged in Plaintiffs’ First Amended Complaint to be true for purposes of resolving the motions to dismiss for failure to state a claim. *Iqbal*, 556 U.S. at 678.

¶¶ 31-33.) One clinical study assessed the effectiveness of Vascepa as an add-on to statin therapy to reduce major cardiovascular events in patients with persistent elevated triglycerides. (*Id.* ¶ 33.) Based on the results of the study, the FDA approved Vascepa in December 2019 for the CV indication, that is, “as an adjunct to maximally tolerated statin therapy to reduce the risk of myocardial infarction, stroke, coronary revascularization, and unstable angina requiring hospitalization in adult patients with elevated triglyceride (TG) levels (≥ 150 mg/dL) and established cardiovascular disease or diabetes mellitus and 2 or more additional risk factors for cardiovascular disease.” (*Id.* ¶ 34, Ex. D.) When the FDA approved the use of Vascepa for the CV indication, Amarin was permitted to add the CV indication to the Vascepa label and remove the CV limitation of use. (*Id.* ¶ 63; *compare id.*, Ex. D with *id.*, Exs. E, F.)

B. The asserted patents

Plaintiffs have patents covering methods of using icosapent ethyl to reduce the risk of cardiovascular events in patients. The '537 patent was issued on July 11, 2017 and is assigned to Mochida. Amarin has an exclusive license. (*Id.* ¶¶ 41-43.) Claim 1 of the '537 patent describes a method of reducing the risk of a cardiovascular event by administering icosapent ethyl with a statin to a patient with high cholesterol, elevated triglycerides, and reduced HDL-C (good cholesterol).⁵ It recites as follows:

1. A method of reducing occurrence of a cardiovascular event in a hypercholesterolemia patient consisting of:
 - identifying a patient having triglycerides (TG) of at least 150 mg/DL and HDL-C of less than 40 mg/dL in a blood sample taken from the patient as a risk factor of a cardiovascular event, wherein the patient has not previously had a cardiovascular event, and administering ethyl icosapentate in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor, wherein said 3-hydroxyl-3-methylglutaryl coenzyme A reductase inhibitor is administered to the patient at least one of before, during and after

⁵ I am attempting to describe the invention in a way that facilitates ease of understanding. In so doing, I make some generalizations about the claim elements. Nothing I say here should be taken as the Court's views on any current or future claim construction (or any other) issues.

administering the ethyl icosapentate; and
 wherein the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor is selected from the group consisting of pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, rosuvastatin, and salts thereof, and
 wherein daily dose of the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor are 5 to 60 mg for pravastatin, 2.5 to 60 mg for simvastatin, 10 to 180 mg for fluvastatin sodium, 5 to 120 mg for atorvastatin calcium hydrate, 0.5 to 12 mg for pitavastatin calcium, 1.25 to 60 mg for rosuvastatin calcium, 5 to 160 mg for lovastatin, and 0.075 to 0.9 mg for cerivastatin sodium.

(*Id.*, Ex. C ('537 Patent).)

The '077 patent was issued on February 4, 2014 and is assigned to Amarin. (*Id.* ¶¶ 46-48.)

Claim 1 describes a method of reducing triglycerides in a patient with mixed dyslipidemia (abnormal lipid levels) on statin therapy by administering icosapent ethyl. Claims 1 and 8 of the '077 patent recite as follows:

1. A method of reducing triglycerides in a subject with mixed dyslipidemia on statin therapy comprising, administering to the subject a pharmaceutical composition comprising about 2500 mg to 5000 mg per day of ethyl eicosapentaenoate and not more than about 5%, by weight of all fatty acids, docosahexaenoic acid or its esters to effect a reduction in fasting triglyceride levels in the subject.

8. The method of claim 1 wherein the subject exhibits a reduction in hs-CRP compared to placebo control.

(*Id.*, Ex. O ('077 Patent).)

The '861 patent was issued on February 25, 2020. It is also assigned to Amarin. (*Id.* ¶¶ 50-52.) Claim 1 describes a method of reducing the risk of cardiovascular death in a patient with established cardiovascular disease by administering icosapent ethyl. Dependent claim 2 specifies that the patient must have a triglyceride level “of about 135 mg/dL to about 500 mg/dL” (*i.e.*, potentially elevated but not necessarily severely high) and an LDL-C (bad cholesterol) level within a specified range. Claims 1 and 2 of the '861 patent recite as follows:

1. A method of reducing risk of cardiovascular death in a subject with established cardiovascular disease, the method comprising administering to said subject about 4 g of

ethyl icosapentate per day for a period effective to reduce risk of cardiovascular death in the subject.

2. The method of claim 1, wherein the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL and a fasting baseline LDL-C level of about 40 mg/dL to about 100 mg/dL.

(*Id.*, Ex. P ('861 Patent).)

After Amarin received FDA approval for the CV indication, it listed the '537, '077, and '861 patents (the “asserted patents”) in the Orange Book for Vascepa. (*Id.* ¶¶ 70-79.)

C. Hikma’s generic product

On November 5, 2020, Hikma launched a generic version of Vascepa after receiving FDA approval of its Abbreviated New Drug Application (ANDA). (*Id.* ¶¶ 11, 13.) Hikma’s ANDA contained a so-called “section viii carve out” regarding the asserted patents. (*Id.* ¶¶ 104, 105.) That is, Hikma represented to the FDA that it would not market its generic product for the uses covered by those patents.

When Hikma originally submitted its ANDA in 2016, it only sought approval for the SH indication, as the FDA had not yet approved Vascepa for the CV indication. (*Id.* ¶ 108.) At that time, Hikma’s proposed generic label (like the Vascepa label at that time) referred only to the SH indication and contained the CV limitation of use. (*Id.*) After Amarin received approval for the CV indication and listed the asserted patents in the Orange Book, Hikma submitted section viii statements with respect to those patents. (*Id.* ¶¶ 104, 108.) Hikma did not propose to add the CV indication to its label, but Hikma did remove the CV limitation of use from its proposed label. (*Id.* ¶ 108.)

The FDA approved Hikma’s ANDA on May 21, 2020. (*Id.* ¶ 105.) The “Indications and Usage” section of Hikma’s approved label refers only to the SH indication, but it does not contain the CV limitation of use. (*Id.*, ¶ 107 Ex. K.) By the time Hikma’s product hit the market in

November 2020, the majority of doctors who prescribed Vascepa did so for uses other than the SH indication, and Hikma was aware of that fact. (*Id.* ¶ 110.)

Hikma issued press releases in 2020 regarding its generic product. In a March 31, 2020 press release, Hikma referred to its then-unapproved product as a “generic version of Amarin Corporation’s Vascepa® 1 gm (icosapent ethyl) capsules.” (*Id.* ¶¶ 111-113, Ex. L.) The press release further stated that “Vascepa® is a prescription medicine that is indicated, *in part*, as an adjunct to diet to reduce triglyceride levels in adult patients with severe (≥ 500 mg/dL) hypertriglyceridemia” (emphasis added), and that the prior year’s “US sales of Vascepa® were approximately \$919 million.” (*Id.*) The Vascepa sales figure cited by Hikma in the press release included sales for the CV indication, and Hikma knew that. (*Id.*) Hikma issued another press release on September 3, 2020 that contained similar statements. (*Id.* ¶¶ 118-120, Ex. M.) Hikma’s March and September 2020 press releases were still accessible on Hikma’s website when Plaintiffs filed this action. (*Id.* ¶¶ 117, 124.)

Hikma’s website also advertises its generic version as being “AB” rated in the “Therapeutic Category: Hypertriglyceridemia.” (*Id.* ¶ 125-126, Ex. T.) That webpage does not refer to the fact that Hikma’s product is only FDA-approved for “severe hypertriglyceridemia.” (*Id.*)

According to the First Amended Complaint, Hikma’s label, press releases, and website “instruct, promote, and encourage” healthcare providers and patients to administer Hikma’s product in a way that infringes the asserted patents. (*Id.* ¶ 127.)

D. Health Net

Health Net is a health insurance provider. (*Id.* ¶ 137.) Vascepa is covered by Health Net’s insurance plans and appears on Health Net’s formularies as a covered drug. (*Id.* ¶ 139.) When Hikma launched its generic version, Health Net added the generic to its formularies, meaning that

it would provide insurance coverage and/or payment for Hikma's product. (*Id.* ¶ 140.) Some of Health Net's formularies currently list Hikma's generic version as a Tier 1 drug and Vascepa as a Tier 3 drug. (*Id.* ¶¶ 143, 157.) The result is that plan beneficiaries have to pay a higher copay for Vascepa than they do for Hikma's generic version. (*Id.* ¶ 145.)

At least one of Health Net's plans requires "Prior Authorization" before it will cover and pay for either Vascepa or Hikma's generic version. (*Id.* ¶¶ 153, 159.) To obtain prior authorization from the plan, the patient's medical provider must submit documentation demonstrating that the prescription is being given for either the SH or the CV indication.⁶ (*Id.* ¶¶ 153, 154, 159, 160, Exs. EE, HH.)

Plaintiffs allege that Health Net is aware that use of Hikma's generic for the CV indication infringes Plaintiffs' patents because (among other reasons) Plaintiffs sent a letter in December 2020 to Mr. Mike Flynn at Envolve Pharmacy Solutions, Inc. (*Id.* ¶ 87.) Envolve is Health Net's Pharmacy Benefit Manager, and Mr. Flynn is Amarin's point of contact for both Envolve and Health Net. (*Id.*) The letter stated that "[t]he Hikma generic does not have an FDA-approved indication for CV risk reduction." (*Id.* ¶¶ 87-90, Ex. GG.) The letter further stated that Amarin "had sued Hikma for patent infringement for encouraging use of its generic product in the CV risk reduction indication" and that "the Hikma generic should not be dispensed for this indication." (*Id.*)

⁶ For example, Health Net's Essential Drug List formulary requires a prior authorization before covering either Vascepa or Hikma's generic version. The prior authorization has criteria that (Amarin contends) map to the SH indication and the CV indication:

- (1) "Hypertriglyceridemia without ASCVD," where the patient has "[f]asting triglycerides \geq 500 mg/dL," or
- (2) "Reduction of Cardiovascular Disease Risk" with "[d]ocumentation (labs must be within 90 days) of fasting triglycerides between 150-499 mg/dL" and, "[f]or members on statin therapy," "Vascepa is prescribed in conjunction with a statin at the maximally tolerated dose."

(*Id.* ¶ 153, Ex. HH; *see also id.* ¶¶ 154, 159-60, Ex. EE.)

According to the First Amended Complaint, Health Net's implementation of the above-described formulary and prior authorization arrangement amounts to encouragement to providers and patients to administer Hikma's product for the CV indication, which, Plaintiffs allege, results in infringement of the asserted patents.

E. Procedural background

Plaintiffs filed their original Complaint on November 30, 2020. (D.I. 1.) The original Complaint only contained claims against Hikma. On January 4, 2021, Hikma filed a motion to dismiss the Complaint for failure to state a claim. (D.I. 11.)

On January 25, 2021, Plaintiffs filed a First Amended Complaint. (D.I. 17.) The First Amended Complaint added new factual allegations against Hikma and added new claims against Health Net. Counts I-III allege that Hikma induces infringement of the '537, '077, and '861 patents under 35 U.S.C. § 271(b). Counts IV-VI allege that Health Net induces infringement of the '537, '077, and '861 patents under 35 U.S.C. § 271(b).

Hikma and Health Net each filed motions to dismiss the claims against them for failure to state a claim. (D.I. 19; D.I. 30.) Health Net also moved to sever Plaintiffs' claims against Health Net from Plaintiffs' claims against Hikma. (D.I. 32.) The Court heard oral argument on all pending motions on May 26, 2021. This is my Report and Recommendation on Hikma's and Health Net's motions to dismiss.

II. LEGAL STANDARD

A defendant may move to dismiss a complaint under Federal Rule of Civil Procedure 12(b)(6) for failure to state a claim. “To survive a motion to dismiss, a complaint must contain sufficient factual matter, accepted as true, to ‘state a claim to relief that is plausible on its face.’” *Iqbal*, 556 U.S. at 678 (quoting *Twombly*, 550 U.S. at 570). A claim is plausible on its face when the complaint contains “factual content that allows the court to draw the reasonable inference that the defendant is liable for the misconduct alleged.” *Id.* (citing *Twombly*, 550 U.S. at 556). A possibility of relief is not enough. *Id.* “Where a complaint pleads facts that are ‘merely consistent with’ a defendant’s liability, it ‘stops short of the line between possibility and plausibility of entitlement to relief.’” *Id.* (quoting *Twombly*, 550 U.S. at 557). In determining the sufficiency of the complaint under the plausibility standard, all “well-pleaded facts” are assumed to be true, but legal conclusions are not. *Id.* at 679.

III. DISCUSSION

Section 271(b) of Title 35 provides that “[w]hoever actively induces infringement of a patent shall be liable as an infringer.” 35 U.S.C. § 271(b). To state a claim of induced infringement under § 271(b), the complaint must plausibly allege that (1) there has been direct infringement, (2) the defendant knowingly induced infringement, and (3) the defendant possessed the intent to encourage another’s infringement. *MEMC Elec. Materials, Inc. v. Mitsubishi Materials Silicon Corp.*, 420 F.3d 1369, 1378 (Fed. Cir. 2005); *FO2GO LLC v. KeepItSafe, Inc.*, No. 18-807-RGA, 2019 WL 1615398, at *3 (D. Del. Apr. 16, 2019).

In the pharmaceutical drug context, a generic manufacturer can be liable under § 271(b) for inducing infringement of a patented method even where the FDA has not approved the generic

product for use in accordance with the patented method.⁷ See *AstraZeneca*, 633 F.3d at 1056-61 (affirming district court’s grant of preliminary injunction against generic manufacturer for inducing infringement of patented method under § 271(b) even though generic product was not approved for patented once-daily use); *GlaxoSmithKline*, 2016 WL 3946770, at *15 (“The decision in *AstraZeneca 2010* indicates that there can, in fact, be situations where a generic manufacturer seeks and obtains a section viii carve-out for a use of a drug that is (according to the FDA) a ‘different’ use from a patented use—and yet the generic’s label could nevertheless be written in such a way that it evidences active steps to induce patent infringement.”); see also *id.*, 2017 WL 1050574, at *1-2 (denying generic defendant’s motion to dismiss inducement claim notwithstanding section viii carve out, where plaintiff alleged that defendant’s label and other conduct encouraged use of the generic product in an infringing manner).

The assessment of whether a complaint plausibly alleges inducement in a pharmaceutical case is thus no different than the analysis in any other case. The court must determine whether the complaint plausibly alleges that the generic manufacturer “offer[ed] a product with the object of promoting its use to infringe, as shown by clear expression or other affirmative steps taken to foster infringement.” *DSU Med. Corp. v. JMS Co.*, 471 F.3d 1293, 1305-06 (Fed. Cir. 2006) (en banc in relevant part). Such “affirmative steps” may include allegations that a defendant “advertis[ed] an infringing use or instruct[ed] how to engage in an infringing use.” *Takeda Pharms. U.S.A., Inc. v. West-Ward Pharm. Corp.*, 785 F.3d 625, 630-31 (Fed. Cir. 2015) (quoting *Metro-Goldwyn-Mayer Studios Inc. v. Grokster, Ltd.*, 545 U.S. 913, 935-36 (2005)).

To be clear, it is not enough to allege that a defendant had “mere knowledge” that its

⁷ In contrast, in an ANDA case, a generic manufacturer cannot be liable under 35 U.S.C. § 271(e)(2) for infringing a method patent unless its ANDA seeks FDA approval for the patented use. *Bayer Schering Pharma AG v. Lupin, Ltd.*, 676 F.3d 1316, 1321-22 (Fed. Cir. 2012); *Warner-Lambert Co. v. Apotex Corp.*, 316 F.3d 1348 (Fed. Cir. 2003).

product could be—or is being—used to infringe. *Warner-Lambert*, 316 F.3d at 1364. Rather, the allegations must plausibly suggest “culpable conduct, directed to encouraging another’s infringement.” *DSU Med.*, 471 F.3d at 1306. Moreover, a defendant who sells a product having substantial noninfringing uses has no duty to take affirmative steps to make sure that others avoid infringement. *Takeda*, 785 F.3d at 632 n.4.

A. Hikma

The First Amended Complaint alleges that Hikma’s product label, press releases, and website encourage infringement of the asserted patents. Hikma contends that the claims against it must be dismissed because the allegations fail to state a plausible claim of inducement. I disagree.

The First Amended Complaint alleges that, notwithstanding the lack of an express instruction regarding the CV indication in the “Indications and Usage” section of Hikma’s label, several other portions of Hikma’s label, taken together with Hikma’s public statements, instruct physicians to use Hikma’s product in a way that infringes the asserted patents. For example, claim 1 of the ’537 patent covers a method of treating hypercholesterolemia patients with elevated triglyceride (TG) levels of at least 150 mg/dL and HDL-C less than 40 mg/mL, and who are on a statin, in order to reduce the risk of a cardiovascular event. The “Dosage and Administration” section of Hikma’s label instructs providers to “[a]ssess lipid levels before initiating therapy.” (D.I. 17 ¶ 130, Ex. K § 2.1.) The “Indications and Usage” section instructs administration to patients with TG levels \geq 500 mg/dL, which, by definition, is at least 150 mg/dL. In addition, the “Clinical Studies” section of Hikma’s label describes treatment of patients with (1) median total cholesterol of 254 mg/dL (*i.e.*, hypercholesterolemia); (2) baseline TG levels between 500 and 2,000 mg/dL, with a median baseline of 684 mg/dL (*i.e.*, \geq 150 mg/dL); (3) a median baseline HDL-C level of 27 mg/dL; and (4) with 25% of the patients on concomitant statin therapy. (*Id.*

¶ 130, Ex. K § 14.2.) The “Patient Information” section describes that the product may be used by patients at risk of having a cardiovascular event. (*Id.* Ex. K.) And, Hikma removed the CV limitation of use from its proposed label, which, according to Plaintiffs, “communicat[es] to the market that Hikma’s generic product has been shown to reduce CV risk.” (*Id.* ¶ 133.) The First Amended Complaint contains similar allegations regarding the ’861 and ’077 patents. (*Id.* ¶¶ 131, 134, Ex. K.)

The FAC further alleges that Hikma is aware that the majority of Vascepa prescriptions are for uses other than the SH indication and that Hikma’s public statements encourage the use of its product for the same indications that Vascepa is used for. (*Id.* ¶¶ 110, 115, 122.) Plaintiffs point to Hikma’s March and September 2020 press releases, which describe its product as a generic version of Vascepa and refer to sales figures that—Hikma knew—include sales for the CV indication. (*Id.* ¶¶ 111, 113, 118, 120.) Plaintiffs also point to Hikma’s website, which advertises its generic version as “AB” rated in the “Therapeutic Category: Hypertriglyceridemia,” which is broader than the “severe hypotriglyceridemia” (SH) indication for which it has FDA approval, and which may suggest administration to patients having merely elevated triglycerides as required by certain claims of the asserted patents. (*Id.* ¶¶ 125-126, Ex. T.)

Those allegations, taken together and viewed in the light most favorable to Plaintiffs, plausibly suggest the following: (1) that Hikma’s label and public statements could instruct and/or encourage third parties to use its product for the CV indication, which Plaintiffs allege is covered by the asserted patents; and (2) that Hikma both knew and intended that third parties would use its product for that purpose. In my view, that is enough.

I am not persuaded by Hikma’s arguments to the contrary. Hikma contends that Plaintiffs have not alleged sufficient “active steps” to encourage infringement. (D.I. 20 at 13-14.) But

Hikma's decision to continue to seek FDA approval after removing the CV limitation of use from its proposed label, its decision to sell its product accompanied by the current version of its label, and its public statements all constitute actions that are alleged to encourage infringement. And, at this stage, those allegations must be viewed in the light most favorable to Plaintiffs.

Hikma also points out that mere knowledge of direct infringement is insufficient to support an inducement claim. That is true. But Plaintiffs allege more than mere knowledge.

Hikma further points out that it has no duty to discourage infringement. Also true. But it cannot present information in a way that encourages infringement. The above-described allegations make it plausible that Hikma, rather than merely failing to prevent infringement, intended to cause others to infringe and knew that their acts would infringe.⁸

To the extent Hikma is suggesting that it cannot be liable for inducement absent FDA approval to use its product for CV therapy and/or explicit instructions in the "Indications and Usage" section of its label to use its product for a CV indication, I disagree. As explained above, lack of FDA approval for an infringing use does not preclude a finding of inducement. *See AstraZeneca*, 633 F.3d at 1060; *see also GlaxoSmithKline*, 2016 WL 3946770, at *13. Many of the cases relied on by Hikma at best establish that were this an ANDA case, and were Plaintiffs' allegations based solely on the label, Plaintiffs' inducement theory might lack merit as a matter of law.⁹ But this is not an ANDA case, and Plaintiffs' allegations are not based solely on the label.

⁸ Of course, in the absence of other evidence of intent, the Court could not find that Defendants are liable for inducement based solely on their failure to take affirmative steps to prevent others' infringement. But Defendants' knowledge that others are using Hikma's product in an infringing way, combined with their failure to take steps to deter such use, could be relevant to their intent to encourage others' infringement. *Cf. Grokster, Ltd.*, 545 U.S. at 939 n.12.

⁹ *See, e.g., Bayer Schering*, 676 F.3d at 1321-24; *AstraZeneca LP v. Apotex, Inc.*, 669 F.3d 1370, 1378-1380 (Fed. Cir. 2012); *Warner-Lambert*, 316 F.3d at 1362-65; *see also GlaxoSmithKline*, 2017 WL 1050574, at *2 (acknowledging difference between claims under § 271(e)(2) and § 271(b)).

Hikma urges the Court to resolve this case at the pleadings stage, pointing out that the contents of its label and public statements are undisputed. But there is a real dispute about what those contents communicate to others, and I do not think it is appropriate to resolve it on a motion to dismiss. Stated another way, at this stage of the case, I am not prepared to say that Hikma’s label and public statements—as a matter of law—could never amount to instruction and encouragement to infringe the asserted patents.

In support of its contention that its actions cannot constitute inducement, Hikma cites the Federal Circuit’s opinions in *HZNP Medicines LLC v. Actavis Laboratories UT, Inc.*, 940 F.3d 680 (Fed. Cir. 2019), *Grunenthal GMBH v. Alkem Laboratories Ltd.*, 919 F.3d 1333 (Fed. Cir. 2019), and *Takeda*, 785 F.3d 625. But none of those cases was resolved at the motion to dismiss stage. *See HZNP*, 940 F.3d at 687-88 (bench trial); *Grunenthal*, 919 F.3d at 1338 (same); *Takeda*, 785 F.3d at 628 (preliminary injunction). And, unlike the allegations in this case, the evidence in those cases related solely to the effects of the generic labels. *See HZNP*, 940 F.3d at 702; *Grunenthal*, 919 F.3d at 1338-39 (“Here, [the plaintiffs] point only to the indications of the proposed labels as grounds for inducement”); *Takeda*, 785 F.3d at 632.¹⁰

I conclude that Plaintiffs have pleaded an inducement claim against Hikma that is at least plausible. While Hikma may be right that Plaintiffs will ultimately be unable to prove inducement, I cannot make that determination at this stage. I recommend that Hikma’s motion to dismiss be denied.

B. Health Net

According to the First Amended Complaint, Health Net’s implementation of its prior

¹⁰ Moreover, while I need not decide whether Plaintiffs’ allegations regarding the label alone state a plausible claim of inducement, I do note that the Federal Circuit in *Takeda* expressly declined to decide “whether evidence as to the invariable response of physicians could ever transform a vague label into active encouragement.” *Takeda*, 785 F.3d at 632.

authorization process for icosapent ethyl prescriptions, combined with its placement of Hikma's generic on the formulary as a tier 1 drug and Vascepa as a tier 3 drug, amounts to encouragement to providers and patients to administer Hikma's product for the unapproved CV indication, which, Plaintiffs allege, results in infringement of the asserted patents.

To my knowledge, this is a novel theory. Neither side has cited any case in which a health insurer has been found liable to a pharmaceutical company for inducing infringement of a drug method of use patent. Viewing the allegations in the light most favorable to Plaintiffs, and in the absence of precedent to the contrary, I cannot say at this stage that Plaintiffs' theory is so implausible as to require dismissal at the pleadings stage.

The thrust of the allegations against Health Net are (1) that it provides coverage and payment for Hikma's generic product even in cases where Health Net actually knows that a particular beneficiary is using the generic version for an unapproved—and allegedly infringing—CV use, and (2) that Health Net actually encourages use of Hikma's product instead of Vascepa for the CV use because Health Net requires its beneficiaries to pay a higher copay for Vascepa than for Hikma's generic version, even when Hikma's version has been prescribed for the infringing/CV use. Plaintiffs allege that Health Net knows when a particular beneficiary is using Hikma's product for the CV use because Health Net's prior authorization process requires the beneficiary's provider to submit documentation supporting the use for which it has been prescribed. Plaintiffs further allege that Health Net had knowledge that its beneficiaries' use of Hikma's product for the CV indication amounted to infringement of Plaintiffs' patents because Amarin sent a pre-suit letter to its point of contact for Health Net informing it of that fact.¹¹ Taken

¹¹ While the letter did not identify the asserted patents by number, the First Amended Complaint plausibly alleges that "[i]t is known in the field, and Health Net would have been aware, that any patents covering a branded drug, such as VASCEPA®, are listed in the Orange Book." (D.I. 17 ¶ 84.)

together, and in the light most favorable to Plaintiffs, it is at least plausible that Health Net knowingly induced infringement and that it had specific intent to do so.

I understand Health Net's position that it merely provides coverage for drugs after they have been prescribed: it neither prescribes medication nor fills the prescriptions. It may ultimately turn out, as Health Net contends, that it has not taken any affirmative acts with the intent to foster infringement. It may also turn out that, despite knowledge of infringement by its beneficiaries and their providers, Health Net's actions in selecting its formulary and adopting its prior authorization procedure for icosapent ethyl prescriptions do not, in fact, influence the decisions of beneficiaries, pharmacists, and medical providers to use, dispense, and prescribe Hikma's generic product in an infringing way or otherwise encourage infringement. It may turn out, as Health Net contends, that "it is Plaintiffs' own pricing decision that encourages use of the generic product over Plaintiffs' brand product." (D.I. 31 at 17.) But all of those are factual issues that are inappropriate for resolution on a motion to dismiss. Plaintiffs allege otherwise, and Plaintiffs' allegations must be taken as true at this stage.

Like Hikma, Health Net points out that it has no duty to discourage others' infringement. While that is true, Plaintiffs also allege that Health Net took active steps—including adopting its formulary and prior authorization procedure for icosapent ethyl prescriptions and taking coverage and payment actions—that are alleged to encourage others' infringement.¹²

I again stress that I am not concluding that this novel claim against a health insurer will or is likely to succeed on the merits. I merely conclude that Plaintiffs have stated a plausible claim and can move forward with discovery.

¹² See also n.8, *supra*.

IV. CONCLUSION

I note that the parties' extensive briefing on the pending motions contained several sub-arguments and cited to several cases not discussed above. I have reviewed those arguments and cases and conclude that they do not warrant further discussion as they do not affect the outcome of the pending motions.

For the reasons set forth above, I recommend that the pending motions to dismiss be DENIED:

1. The Court should deny Hikma's motion to dismiss the First Amended Complaint. (D.I. 19.)
2. The Court should deny Hikma's motion to dismiss the original Complaint as moot. (D.I. 11.)
3. The Court should deny Health Net's motion to dismiss the First Amended Complaint. (D.I. 30.)

This Report and Recommendation is filed pursuant to 28 U.S.C. § 636(b)(1)(B),(C), Federal Rule of Civil Procedure 72(b)(1), and District of Delaware Local Rule 72.1. Any objections to the Report and Recommendation shall be filed within fourteen days and limited to ten pages. Any response shall be filed within fourteen days thereafter and limited to ten pages. The failure of a party to object to legal conclusions may result in the loss of the right to *de novo* review in the district court.

The parties are directed to the Court's "Standing Order for Objections Filed Under Fed. R. Civ. P. 72," dated October 9, 2013, a copy of which can be found on the Court's website.

Dated: August 3, 2021



 Jennifer L. Hall
 UNITED STATES MAGISTRATE JUDGE



US009700537B2

(12) **United States Patent**
Yokoyama et al.

(10) **Patent No.:** **US 9,700,537 B2**
(45) **Date of Patent:** ***Jul. 11, 2017**

(54) **COMPOSITION FOR PREVENTING THE OCCURRENCE OF CARDIOVASCULAR EVENT IN MULTIPLE RISK PATIENT**

(71) Applicant: **MOCHIDA PHARMACEUTICAL CO., LTD.**, Tokyo (JP)

(72) Inventors: **Mitsuhiro Yokoyama**, Kobe (JP); **Hideki Origasa**, Toyama (JP); **Masunori Matsuzaki**, Ube (JP); **Yuji Matsuzawa**, Takaruzuka (JP); **Yasushi Saito**, Chiba (JP)

(73) Assignee: **MOCHIDA PHARMACEUTICAL CO., LTD.**, Tokyo (JP)

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **15/431,958**

(22) Filed: **Feb. 14, 2017**

(65) **Prior Publication Data**

US 2017/0151206 A1 Jun. 1, 2017

Related U.S. Application Data

(62) Division of application No. 14/963,291, filed on Dec. 9, 2015, which is a division of application No. 14/474,955, filed on Sep. 2, 2014, now abandoned, which is a division of application No. 12/302,790, filed as application No. PCT/JP2007/061099 on May 31, 2007, now Pat. No. 8,853,256.

(30) **Foreign Application Priority Data**

May 31, 2006 (JP) 2006-152740

(51) **Int. Cl.**

A61K 31/232 (2006.01)
A61K 31/40 (2006.01)
A61K 31/366 (2006.01)
A61K 31/225 (2006.01)
A61K 31/202 (2006.01)
A61K 31/22 (2006.01)
A61K 31/351 (2006.01)

(52) **U.S. Cl.**

CPC **A61K 31/22** (2013.01); **A61K 31/202** (2013.01); **A61K 31/225** (2013.01); **A61K 31/232** (2013.01); **A61K 31/351** (2013.01); **A61K 31/366** (2013.01); **A61K 31/40** (2013.01); **A61K 2300/00** (2013.01)

(58) **Field of Classification Search**

USPC 514/560
See application file for complete search history.

(56) **References Cited**

U.S. PATENT DOCUMENTS

5,260,305 A	11/1993	Dennick
5,656,667 A	8/1997	Breivik et al.
5,753,703 A	5/1998	Cavazza et al.
6,174,865 B1	1/2001	Klein
8,802,718 B2	8/2014	Yokoyama et al.
8,853,256 B2	10/2014	Yokoyama
2004/0009208 A1	1/2004	Edson et al.
2005/0113449 A1	5/2005	Renshaw
2007/0021504 A1	1/2007	Yokoyama et al.

FOREIGN PATENT DOCUMENTS

EP	1 157 692 A1	11/2001
EP	1 790 339 A1	5/2007
WO	WO 00/48592 A1	8/2000
WO	WO 2007/007686 A1	1/2007

OTHER PUBLICATIONS

"Aha Hatsu Sokuho JELIS Kekka Happyo," Medical Tribune, (Nov. 17, 2005), Tokubetsu Kikaku Dai 3 Bu, pp. 75 to 76 (Doitsu Naiyo no Website Kiji: URL, <http://www.medical-tribune.jp/congress/jelis/jelis.html>).

Adan et al., "Effects of docosahexaenoic and eicosapentaenoic acid on lipid metabolism, eicosanoid production, platelet aggregation and atherosclerosis in hypercholesterolemic rats", Bioscience, Biotechnology, And Biochemistry, vol. 63, No. 1, pp. 111-119, (Jan. 1999). XP002613171.

European Search Report dated Dec. 21, 2010, for European Patent Appl. No. 07744487.5.

Fruchart et al., "The Residual Risk Reduction Initiative: A Call to Action to Reduce Residual Vascular Risk in Patients with Dyslipidemia," Am. J. Cardiol. (2008), vol. 102 [Suppl]: pp. 1K-34K.

Ishikawa, "JELIS no Jisshi Keikaku to Kitai sareru Kekka," [online], JELIS Medical Asahi (Medical Asahi 1996, 12 Bassui), Mar. 29, 2007 Access], Internet, URL, <http://www.mochida.co.jp/dis/jelis/jlnwepm2.html>.

Kastelein et al., Simvastatin with or without Ezetimibe in Familial Hypercholesterolemia, N. Engl. J. Med. (2008), vol. 358, pp. 1431-1443.

Medical Drug Interview Form, EPA Preparation, EPADEL Capsules 300, Revised in Jul. 2002 and Feb. 2004, pp. 21-22, 21st Edition, issued in Dec. 2004.

Mochida Pharmaceutical Co., Ltd. New Release, "'Epadel' ga Kandomyaku Shikkan no Hassho to Saihatsu o Yokusei Daikibo Shiken 'JELIS' no Aratana Kaiseki Kekka ga Kohyo sare mashita", Nov. 15, 2006 Happyo, URL, <http://www.mochida.co.jp/news/2006/pdf/1115.pdf>.

(Continued)

Primary Examiner — Jennifer M Kim

(74) Attorney, Agent, or Firm — Birch, Stewart, Kolasch & Birch, LLP

(57) **ABSTRACT**

Disclosed is a composition which is useful for preventing the occurrence of a cardiovascular event, particularly a composition which is expected to show a prophylactic effect on a cardiovascular event occurring in a hypercholesterolemia patient despite providing the patient with a treatment with HMG-CoA RI or a cardiovascular event occurring in a multiple risk patient.

16 Claims, 1 Drawing Sheet

US 9,700,537 B2

Page 2

(56)

References Cited

OTHER PUBLICATIONS

Mochida Pharmaceutical Co., Ltd. New Release “Mochida Announces Completion of ‘JELIS’ Major Clinical Trial for ‘Epadel’ Hyperlipidemia and Artherosclerosis Obliterans Therapeutic Treatment” <http://www.mochida.co.jp/english/new/2005/0322.html>, Mar. 22, 2005.

Yokoyama et al., “Effects of Eicosapentaenoic Acid (EPA) on Major Cardiovascular Events in Hypercholesterolemic Patients,” The Japan EPA Lipid Intervention Study (JELIS), Slides 1-30 (2005).
Yokoyama et al., American Heart Journal, (2003), vol. 146, No. 4, pp. 613-620.

Yokoyama et al., Lancet (2007), vol. 369, No. 9567, pp. 1090-1098.
Yokoyama, M., “Effects of Eicosapentaenoic Acid (EPA) on Major Cardiovascular Events in Hypercholesterolemic Patients: the Japan EPA Lipid Intervention Study (JELIS),” Circulation, Late-Breaking Clinical Trial Abstracts, vol. 112, No. 21, pp. 3362-3364 (2005).
Yokoyama, M., “JELIS Revealed EPA Preparation’s Inhibitory Action on Coronary Events,” Announcement at AHA 2005, Dallas, Texas, Nov. 14, 2005.

Yorioka et al., “Lipid-lowering therapy and coagulation/fibrinolysis parameters in patients on peritoneal dialysis,” The International Journal of Artificial Organs (Jan. 2000), vol. 23, No. 1, pp. 27-32.

U.S. Patent

Jul. 11, 2017

US 9,700,537 B2

FIG. 1





FIG. 2



US 9,700,537 B2

1

COMPOSITION FOR PREVENTING THE OCCURRENCE OF CARDIOVASCULAR EVENT IN MULTIPLE RISK PATIENT

CROSS-REFERENCE TO RELATED APPLICATIONS

The present application is a 37 C.F.R. §1.53(b) divisional of U.S. application Ser. No. 14/963,291 filed Dec. 9, 2015, which is a 37 C.F.R. §1.53(b) divisional of U.S. application Ser. No. 14/474,955 filed Sep. 2, 2014 (abandoned), which is a 37 C.F.R. §1.53(b) divisional of U.S. application Ser. No. 12/302,790 filed Nov. 26, 2008 (now U.S. Pat. No. 8,853,256 B2 issued Oct. 7, 2014), which is the National Phase of PCT International Application No. PCT/JP2007/061099 filed May 31, 2007, which in turn claims priority on Japanese Patent Application No. 2006-152740 filed May 31, 2006. The entire contents of each application is hereby incorporated by reference.

TECHNICAL FIELD

This invention relates to a composition for preventing occurrence of cardiovascular events (primary prevention) in multiple risk patients, the composition containing at least ethyl icosapentate (hereinafter abbreviated as EPA-E).

BACKGROUND ART

Westernization of diet has resulted in the increase of patients suffering from lifestyle-related diseases such as diabetes, hyperlipidemia, and hypertension. Some of these diseases finally lead to arteriosclerotic diseases such as myocardial infarction, angina pectoris, and cerebral infarction. Treatment of the lifestyle-related diseases is based on the improvement of lifestyle, and more specifically, on the alimentary therapy and kinesitherapy. However, such improvement of the dietary life or the lack of exercise is often difficult in the patients suffering from the "lifestyle-related diseases," and they usually transfer to pharmacotherapy in order to prevent poor prognosis, for example, onset of myocardial infarction or cerebral infarction.

An exemplary compound having the action of improving such lifestyle-related diseases is polyunsaturated fatty acid. The polyunsaturated fatty acid is defined as a fatty acid including two or more carbon-carbon double bonds in one molecule, and the polyunsaturated fatty acids are categorized by the position of the double bond into ω 3 fatty acid, ω 6 fatty acid, and the like. The ω 3 polyunsaturated fatty acids include α -linolenic acid, icosapentaenoic acid (EPA), and docosahexaenoic acid (DHA), and the ω 6 polyunsaturated fatty acids include linoleic acid, γ -linolenic acid, and arachidonic acid. Polyunsaturated fatty acids are derived from natural products, and exhibit various actions including antiarteriosclerotic action, platelet aggregation inhibitory action, hypolipidemic action, antiinflammatory action, antitumor action, and central action, and due to the high safety, polyunsaturated fatty acids are incorporated in various kinds of food, or sold as a health food or drug.

Decrease in the death rate in the patients who have history of suffering from myocardial infarction has been reported for the administration of a mixture of ethyl ester of an ω -3 polyunsaturated fatty acid EPA (EPA-E) and ethyl ester of an ω -3 polyunsaturated fatty acid DHA (DHA-E) for 3.5 years (see Patent Document 1). However, the results disclosed in Patent Document 1 relates to the secondary prevention, that

2

is, prevention of recurrence, and the drug which is effective in the secondary prevention is not always effective in the primary prevention.

Based on the results of animal experiments and small scale clinical observations, many large scale clinical trials have been recently planned and conducted for the purpose of confirming whether various drugs which are effective in improving the lifestyle-related diseases can also prevent arteriosclerotic diseases in human. The results, however, have not necessarily been as intended, and the situation is still severe for the prevention of the occurrence of cardiovascular events in the case of patients suffering from a plurality of risk factors.

High purity EPA-E is commercially available in the trade names of Epadel™ and Epadel ST™ (manufactured by Mochida Pharmaceutical Co., Ltd.) as therapeutic drugs for hyperlipidemia. There has been reported that when such high purity EPA-E is orally administered at 600 mg per administration and 3 times a day immediately after meal (when TG is abnormal, the dose is increased to the level of 900 mg per administration and 3 times a day), serum T-Cho concentration can be reduced by 3 to 6%, and serum TG can be reduced by 14 to 20% (see Non-Patent Document 1). There has also been reported in The Heart Failure Society of America 2005 Annual Meeting that, based on such action, such high purity EPA-E was expected to have the effects of improving cardiovascular events in hyperlipidemia patients, and combined use with HMG-CoA RI was effective in inhibiting cardiac events in a large scale clinical trial. In this large scale clinical trial (DELIS, Japan EPA Lipid Intervention Study), statistically significant suppression of the cardiac events by the EPA-E was confirmed for the total of the primary prevention patients and secondary prevention patients, and for the secondary prevention patients. On the other hand, in the analysis limited to the primary prevention patients, the incidence of the events was lower in the EPA-E group (the group administered with EPA-E in combination with HMG-CoA RI) compared to the control group (the group administered with solely with HMG-CoA RI), while this difference was not statistically significant. This trial also revealed that after 5 years from the start of the trial, the LDL-cholesterol value reduced by 26% in both of the EPA-E group and control group, that no significant difference was found between these groups, and that change of the HDL-cholesterol value was slight in both groups (see Non-Patent Document 2). This trial also revealed that the total cholesterol and the LDL-cholesterol decreased by 19% and 25%, respectively, in both the EPA-E group and the control group, and that triglyceride decreased by 9% (significant) and 4% in the EPA-E group and the control group, respectively, while little change in HDL-C was noted in both the EPA-E group and the control group (see Non-Patent Document 3). There is so far no report that has analyzed prevention of the occurrence of the cardiovascular events in the case of patients having two or more risk factors.

Patent Document 1: WO 00/48592 (JP 2002-537252 A)

Non-Patent Document 1: Drug Interview Form "EPA preparation, Epadel capsule 300", revised in July, 2002, and February, 2004, version 21 issued in December, 2004; pp. 21-22.

Non-Patent Document 2: Medical Tribune, issue of Nov. 17, 2005, Feature article 3, pp. 75-76.

Non-Patent Document 3: Lancet, vol. 369, pages 1090 to 1098 (2007).

US 9,700,537 B2

3

DISCLOSURE OF THE INVENTION

Problems to be Solved by the Invention

In view of the situation that there is a serious problem that death from the cardiovascular disease is still a major cause of the death, and many cases of cardiovascular events are still impossible to prevent by the HMG-CoA RI therapy, an object of the present invention is to provide a composition for preventing onset of the cardiovascular events.

Means to Solve the Problems

In order to solve the problems as described above, the inventors of the present invention made an extensive study on a therapy of hypercholesterolemia patients and found that EPA-E has the effect of preventing occurrence of the cardiovascular events in patients suffering from multiple risk factors, and in particular, the effect of preventing occurrence of the cardiovascular events in male patients suffering from multiple risk factors. The present invention has been completed on the bases of such finding. Accordingly, the present invention is directed to the following:

(1) A composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

(2) A composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the hypercholesterolemia patient is a patient also suffering from two or more of the risk factors.

(3) A composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one of risk factors as defined by a body mass index (BMI) of at least 25 for the obesity; by a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg for the hypertension or the prehypertension; by a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5% for the diabetes, the prediabetes, or the abnormal glucose tolerance; and by triglyceride (TG) of at least 150 mg/dL and/or a HDL-C of less than 40 mg/dL for the hypertriglyceridemia and/or the low HDL cholesterol.

(4) The composition according to any one of (1) to (3) above, wherein the content of the EPA-E is at least 96.5% by weight in relation to the total content of fatty acid and derivatives thereof.

(5) The composition according to any one of (1) to (4) above, wherein the EPA-E is orally administered at a dose of 1.8 g/day to 2.7 g/day.

(6) The composition according to any one of (1) to (5) above, wherein the composition is used in combination with HMG-CoA RI.

(7) The composition according to any one of (1) to (6) above, wherein the hypercholesterolemia patient is a male patient.

4

(8) The composition according to any one of (1) to (7) above, wherein the hypercholesterolemia patient is a patient also suffering from hypertriglyceridemia and low HDL cholesterol.

(9) A method for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient by administering the patient with the composition according to any one of (1) to (8) above.

(10) Use of the composition according to any one of (1) to (8) above for the manufacture of an agent for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient.

Merits of the Invention

The above-mentioned composition of the present invention containing at least EPA-E as its effective component is effective in preventing occurrence of cardiovascular events in hypercholesterolemia patients, and in particular, in preventing occurrence of cardiovascular events in hypercholesterolemia patients who have been treated with HMG-CoA RI but still suffer from the risk of the cardiovascular events, or more particularly, in preventing occurrence of cardiovascular events in hypercholesterolemia patients also suffering from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

The effect of the composition of the present invention will be synergistically improved by combined use with the HMG-CoA RI, and such use of the composition of the present invention with the HMG-CoA RI has clinical utility since the effect of preventing the cardiovascular event occurrence is expected to be improved.

BRIEF DESCRIPTION OF THE DRAWINGS

FIG. 1 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and the time after the start of the trial in X-axis for male patients having at least 2 risk factors.

FIG. 2 is a graph prepared by plotting the incidence of the cardiovascular event in Y-axis and the time after the start of the trial in X-axis for patients having the risk factors of a triglyceride (TG) of at least 150 mg/dL and a HDL-C of less than 40 mg/dL.

BEST MODE FOR CARRYING OUT THE INVENTION

Next, the present invention is described in detail.

A first aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

US 9,700,537 B2

5

Alternatively, the first aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

The prevention of the occurrence of the cardiovascular events include all cases of primary prevention, and exemplary cases include prevention of cardiovascular death, fatal myocardial infarction, sudden cardiac death, nonfatal myocardial infarction, cardiovascular angioplasty, new occurrence of rest angina and exercise-induced angina, and destabilization of the angina. The composition of the present invention may be administered to any person who needs prevention of the occurrence of the cardiovascular events, and typical such patients are hypercholesterolemia patients.

A second aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient undergoing a HMG-CoA RI therapy, the composition containing at least EPA-E, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

Alternatively, the second aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient undergoing a HMG-CoA RI therapy, the composition containing at least EPA-E and/or DHA-E, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

While HMG-CoA RI includes all those having inhibitory action for 3-hydroxy-3-methylglutaryl coenzyme A reductase, the one used in the present invention is preferably a pharmaceutically administrable inhibitor which is preferably at least one member selected from the group consisting of pravastatin, simvastatin, lovastatin, fluvastatin, cerivastatin, atorvastatin, pitavastatin, rosuvastatin, and salts and derivatives thereof, and more preferably, pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, or rosuvastatin, and most preferably, pravastatin or simvastatin. All salts are included as long as they are pharmaceutically administrable, and preferred are sodium and calcium salts such as pravastatin sodium, fluvastatin sodium, cerivastatin sodium, atorvastatin calcium, pitavastatin calcium, and rosuvastatin calcium. In the present invention, "pravastatin," for example, also includes the pravastatin in the form of a salt unless otherwise noted.

A third aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event

6

in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol; namely; obesity, and hypertension or prehypertension; obesity, and diabetes, prediabetes, or abnormal glucose tolerance; obesity, and hypertriglyceridemia and/or low HDL cholesterol; hypertension or prehypertension, and diabetes, prediabetes or abnormal glucose tolerance; hypertension or prehypertension, and hypertriglyceridemia and/or low HDL cholesterol; diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterol; obesity, and hypertension or prehypertension, and diabetes, prediabetes, or abnormal glucose tolerance; obesity, and hypertension or prehypertension, and hypertriglyceridemia and/or low HDL cholesterol; obesity, and diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterol; hypertension or prehypertension, and diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterol; obesity, and hypertension or prehypertension, and diabetes, prediabetes, or abnormal glucose tolerance, and hypertriglyceridemia and/or low HDL cholesterol.

Alternatively, the third aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol.

A fourth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one, and more preferably, at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterol. In this case, the hypercholesterolemia patient is preferably a male patient.

Alternatively, the fourth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one, and more preferably, at least two risk factors selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,

US 9,700,537 B2

7

- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia. In this case, the hypercholesterolemia patient is preferably a male patient.

A fifth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from risk factors of hypertriglyceridemia and low HDL cholesterolemia, and more specifically, hypertriglyceridemia and low HDL cholesterolemia with a serum triglyceride (TG) concentration of at least 150 mg/dl and a serum HDL-C concentration of less than 40 mg/dl, or serum TG/HDL-C ratio of at least 3.75. In this case, the hypercholesterolemia patient is preferably a male patient. Alternatively, the fifth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event in a hypercholesterolemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from risk factors of hypertriglyceridemia and low HDL cholesterolemia, and more specifically, hypertriglyceridemia and low HDL cholesterolemia with a serum triglyceride (TG) concentration of at least 150 mg/dl and a serum HDL-C concentration of less than 40 mg/dl, or a serum TG/HDL-C ratio of at least 3.75. In this case, the hypercholesterolemia patient is preferably a male patient.

A sixth aspect of the present invention provides a composition containing at least EPA-E as its effective component, the composition exhibiting an excellent effect of preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors who has been administered with this composition for at least 2 years since the start of the administration. Alternatively, the sixth aspect of the present invention provides a composition containing at least EPA-E and/or DHA-E as its effective component, the composition exhibiting an excellent effect of preventing recurrence of a cardiovascular event in a patient suffering from multiple risk factors who has been administered with this composition for at least 2 years since the start of the administration. The hypercholesterolemia patient is preferably a male patient.

A seventh aspect of the present invention provides a method for preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors by continuously administering the patient with a composition containing at least EPA-E as its effective component for at least 2 years. Alternatively, the seventh aspect of the present invention provides a method for preventing occurrence of a cardiovascular event in a patient suffering from multiple risk factors by continuously administering the patient with a composition containing at least EPA-E and/or DHA-E as its effective component for at least 2 years. The hypercholesterolemia patient is preferably a male patient.

An eighth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a dyslipidemia patient, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

8

Alternatively, the eighth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a dyslipidemia patient, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

A ninth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient to be able to administered with HMG-CoA RI, the composition containing at least EPA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

Alternatively, the ninth aspect of the present invention provides a composition for preventing occurrence of a cardiovascular event (primary prevention) in a hypercholesterolemia patient to be able to administered with HMG-CoA RI, the composition containing at least EPA-E and/or DHA-E as its effective component, wherein the patient also suffers from at least one risk factor selected from the group consisting of

- (1) obesity,
- (2) hypertension or prehypertension,
- (3) diabetes, prediabetes, or abnormal glucose tolerance, and
- (4) hypertriglyceridemia and/or low HDL cholesterolemia.

While the EPA-E content in the total fatty acid and dosage are not particularly limited as long as intended effects of the present invention are attained, the EPA-E used is preferably the one having a high purity, for example, the one having the proportion of the EPA-E in the total fatty acid and derivatives thereof of preferably 40% by weight or higher, more preferably 90% by weight or higher, and still more preferably 96.5% by weight or higher. The daily dose in terms of EPA-E is typically 0.3 to 6 g/day, preferably 0.9 to 3.6 g/day, and still more preferably 1.8 to 2.7 g/day. Another preferable daily dose is 0.3 to 2.7 g/day, and 0.3 to 1.8 g/day. Another preferable fatty acid included is DHA-E. While the compositional ratio of EPA-E/DHA-E, content of EPA-E and DHA-E (hereinafter referred to as (EPA-E+DHA-E)) in the total fatty acid, and dosage of (EPA-E+DHA-E) are not particularly limited as long as intended effects of the present invention are attained, the composition is preferably the one having a high purity of EPA-E and DHA-E, for example, the one having a proportion of the (EPA-E+DHA-E) in the total fatty acid and derivatives thereof of preferably 40% by weight or higher, more preferably 80% by weight or higher, and still more preferably 90% by weight or higher. The daily dose in terms of EPA-E+DHA-E is typically 0.3 to 10 g/day, preferably 0.5 to 6 g/day, and still more preferably 1 to 4 g/day. Another preferable daily dose is 0.3 to 6 g/day, 0.3 to 4 g/day, and 0.3 to 1 g/day. The content of other long chain saturated fatty acids is preferably low, and among the long

US 9,700,537 B2

9

chain unsaturated fatty acids, the content of $\omega 6$ fatty acids, and in particular, the content of arachidonic acid is preferably as low as less than 2% by weight, and more preferably less than 1% by weight.

The composition of the present invention contains EPA-E and/or DHA-E, and has the effect of preventing occurrence of cardiovascular events in healthy people or those suffering from the risk factors of hyperlipidemia, diabetes, and hypertension when the composition is orally administered, and in particular, of preventing occurrence of cardiovascular events in hypercholesterolemia patients who have been treated with HMG-CoA RI but still suffering from the risk of the cardiovascular events. The composition of the present invention may also be used in combination with the HMG-CoA RI, and such combination may further prevent the occurrence of the cardiovascular events.

The composition of the present invention may be used with other drugs, for example, antiplatelet drugs such as aspirin, ticlopidine, clopidogrel, prasugrel, and cilostazol; anticoagulants such as warfarin, heparin, and ximelagatran; antihypertensive drugs such as angiotensin II receptor antagonists (candesartan, losartan, valsartan, etc.), angiotensin converting enzyme inhibitors, calcium channel antagonists (amlodipine, cilnidipine, etc.), and $\alpha 1$ blockers; diabetes drugs or abnormal glucose tolerance stimulants such as α -glucosidase inhibitors (voglibose, acarbose, etc.), biguanide drugs, thiazolidinedione drugs (pioglitazone, rosiglitazone, rivoglitazone, etc.), and prompt insulin release promoters (mitiglinide, nateglinide, etc.); antilipotropic drugs and antiarteriosclerotic drugs such as HMG-CoA RI as described above, fibrate drugs, squalene synthetase inhibitors (TAK-475, etc.), and cholesterol absorption inhibitors (ezetimibe, etc.), probucol, anion exchange resin, nicotinic acid drugs, phytosterol, elastase, dextran sulfate sodium sulfur, pantothenic acid, and polyenephosphatidylcholine.

The composition of the present invention contains smaller amounts of impurities such as saturated fatty acids and arachidonic acid which are unfavorable for cardiovascular events compared to fish oil or fish oil concentrate, and accordingly, the intended effects can be attained without causing problems like overnutrition or excessive intake of vitamin A. In addition, since the effective component of the present composition is in the form of an ester, the effective component is more stable to oxidation compared to the case of fish oil in which the effective component is in the form of a triglyceride, and a sufficiently stable composition can be produced by adding a conventional antioxidant. In other words, it is the use of the EPA-E that has for the first time enabled production of a composition for preventing onset of cardiovascular events which can be used in clinical practice.

In the present invention, the term "icosapentaenoic acid" designates all-cis-5,8,11,14,17-icosapentaenoic acid.

In the present invention, the term "hypercholesterolemia patient" means the patient with increased serum T-Chol concentration or serum LDL-Chol concentration. In a narrower sense, this term means the patient suffering from hypercholesterolemia (serum T-Chol concentration of at least about 220 mg/dl, and in more strict sense, at least 250 mg/dl) or high LDL cholesterol (serum LDL-Chol concentration of at least 140 mg/dl).

In the present invention, the term "dyslipidemia" is the condition which satisfies at least one of high LDL cholesterol (i.e. fasting serum LDL cholesterol value of at least 140 mg/dL), low HDL cholesterol (i.e. fasting serum HDL cholesterol value of less than 40 mg/dL), and hypertriglyceridemia (i.e. fasting serum triglyceride value of at least 150 mg/dL) according to the diagnostic criteria

10

described in "Guideline for Preventing Arteriosclerotic diseases, 2007" (edited and published by Japan Atherosclerosis Society).

Of the risk factors treated in the present invention, "obesity" is the state with excessive accumulation of fats in the body. For example, non-limiting examples of the obesity include a body mass index (BMI) of at least 25, a waist measurement of at least 85 cm in male and at least 90 cm in female. "Hypertension" is the state with an abnormal increase in resting arterial blood pressure of the greater circulatory system. For example, in the criteria proposed by Japanese Society of Hypertension at the time of the filing of this application, hypertension is defined as a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg. "Prehypertension" is the condition with the blood pressure between the normal blood pressure (or optimal blood pressure) and the blood pressure in the hypertension, and this condition is also referred to as "mild elevated blood pressure" or "borderline hypertension." Non-limiting exemplary criteria for such condition include a systolic blood pressure (SEP) of 120 to 139 mmHg or a diastolic blood pressure (DBP) of 80 to 89 mmHg. In the present invention, "hypertension or prehypertension" means a condition with a systolic blood pressure (SBP) of at least 120 mmHg or a diastolic blood pressure (DBP) of at least 80 mmHg, more strictly, a systolic blood pressure (SBP) of at least 135 mmHg or a diastolic blood pressure (DBP) of at least 85 mmHg, even more strictly a systolic blood pressure (SBP) of at least 140 mmHg or a diastolic blood pressure (DBP) of at least 90 mmHg. "Diabetes" is the glucose metabolism disorder caused by hyposecretion of insulin from the insulin-producing cell (β cell) in the pancreas or insufficient action of the insulin in the target cell. Exemplary non-limiting criteria proposed by Japan Diabetes Society at the time of the filing of this application is one of 1) fasting blood glucose of at least 126 mg/dL, 2) 75 g glucose tolerance test at 2 hours of at least 200 mg/dL, and 3) casual blood glucose level of at least 200 mg/dL; or a hemoglobin A1c (HbA1c) of at least 6.5%. The criteria, however, are not limited to these. "Prediabetes" is the condition in which the blood glucose level is between the normal value and the value in the diabetes. "Abnormal glucose tolerance" is the condition in which the blood glucose level in the glucose tolerance test is between the normal value and the value in the diabetes. These conditions are also referred to as the borderline diabetes, prediabetic state, and the diabetic high-risk group. For these conditions, exemplary non-limiting criteria include a fasting blood glucose of 110 to 125 mg/dL, a 75 g glucose tolerance test at 2 hours of 140 to 199 mg/dL, and a hemoglobin A1c (HbA1c) of 5.6 to 6.4%. In the present invention, "diabetes, prediabetes, or abnormal glucose tolerance" means a condition with a fasting blood glucose (FBS) of at least 110 mg/dL or a hemoglobin A1c (HbA1c) of at least 5.6%, more strictly, a fasting blood glucose (FBS) of at least 110 mg/dL or a hemoglobin A1c (HbA1c) of at least 5.9%, and even more strictly with a fasting blood glucose (FBS) of at least 126 mg/dL or a hemoglobin A1c (HbA1c) of at least 6.5%. "Hypertriglyceridemia" is the condition with an increased serum triglyceride (TG) concentration, and strictly, with the serum TG concentration of at least 150 mg/dL. "Low HDL cholesterol" is the condition with a reduced serum HDL-C concentration, and strictly, with the serum HDL-C concentration of less than 40 mg/dL. In the present invention, "hypertriglyceridemia and/or low HDL cholesterol" means the state with a serum TG concentration of at least 150 mg/dL and/or a serum HDL-C concentration of

US 9,700,537 B2

11

less than 40 mg/dL. The hypertriglyceridemia and the low HDL cholesterolemia are both diseases included in the category of dyslipidemia, and they are mutually independent risk factors. Combination of these risk factors, however, is known to result in an increased risk of the occurrence of an arteriosclerotic disease. In the present invention, “the hypertriglyceridemia and/or the low HDL cholesterolemia” is treated as a single risk factor.

In the present invention, the term “combined use of EPA-E with HMG-CoA RI” include both the embodiment in which the EPA-E and the HMG-CoA RI are simultaneously administered and the embodiment in which both agents are separately administered. When these agents are simultaneously administered, they may be formulated either as a single combined drug or separate drugs. When these agents are separately administered, EPA-E may be administered either before or after the HMG-CoA RI. The doses and ratio of the EPA-E and the HMG-CoA RI may be adequately selected.

In the present invention, the term “combined use of EPA-E and/or DHA-E with HMG-CoA RI” include both the embodiment in which the EPA-E and/or DHA-E and the HMG-CoA RI are simultaneously administered and the embodiment in which these agents are separately administered. When these agents are simultaneously administered, they may be formulated either as a single combined drug or separate drugs. When these agents are separately administered, EPA-E and/or DHA-E may be administered either before or after the HMG-CoA RI. The doses and ratio of the EPA-E and/or DHA-E and the HMG-CoA RI may be adequately selected.

The composition of the present invention has the action of preventing onset of the cardiovascular events by the sole administration of the composition, and in particular, the present composition is expected to have the effect of preventing onset of the cardiovascular events which could not be prevented by the sole administration of the HMG-CoA RI. In addition, EPA-E has not only the action of reducing the serum T-Cho concentration and the serum TG, but also the action of suppressing platelet aggregation based on inhibition of arachidonic acid cascade, which is a pharmacological action different from the HMG-CoA RI. Therefore, the action as described above can also be exerted by combined administration with the HMG-CoA RI.

Since EPA-E and DHA-E are highly unsaturated, inclusion of an effective amount of an antioxidant such as butylated hydroxytoluene, butylated hydroxyanisole, propyl gallate, gallic acid, and pharmaceutically acceptable quinine, or α -tocopherol is preferable.

The preparation may be orally administered to the patient in the dosage form of tablet, capsule, microcapsule, granules, fine granules, powder, oral liquid preparation, syrup, or jelly. Preferably, the preparation is orally administered by filling in a capsule such as soft capsule or microcapsule.

The soft capsules containing high purity EPA-E (EpadelTM and Epadel STM) are commercially available in Japan as safe therapeutic agents for arteriosclerosis obliterans and hyperlipidemia with reduced side effects, and in such products, proportion of EPA-E in total fatty acid is at least 96.5% by weight. The soft capsule (OmacorTM, Ross products, Reliant, and Pronova) containing about 46% by weight of EPA-E and about 38% by weight of DHA-E is commercially available in the U.S., Europe, and other countries as a drug applied for hypertriglyceridemia. These drugs may be purchased for use in the present invention.

The dose and administration period of the composition of the present invention for preventing the onset of the cardio-

12

vascular events is the dose and period sufficient for the expression of the intended action, and the dose and administration period may be adequately adjusted depending on the dosage form, administration route, daily frequency, severity of the symptoms, body weight, age, and the like. When orally administered, the composition may be administered at a dose in terms of EPA-E of 0.3 to 6 g/day, preferably 0.9 to 3.6 g/day, and more preferably 1.8 to 2.7 g/day, and while such dose is typically administered in 3 divided doses, if desired, such dose may be administered in a single dose or in several divided doses. The composition is preferably administered during or after the meal, and more preferably, immediately (within 30 minutes) after the meal. When such dose is orally administered, the administration period is typically at least 1 year, preferably at least 2 years, more preferably at least 3 years and still more preferably at least 5 years. The administration, however, is preferably continued as long as there is a considerable risk of onset of the cardiovascular events. If necessary, drug holidays of about 1 day to 3 months, and preferably about 1 week to 1 month may be given.

The HMG-CoA RI is preferably used according to the dosage regimen recommended for the particular drug used, and the dose may be adequately adjusted depending on the type, dosage form, administration route, daily frequency, severity of the symptoms, body weight, gender, age, and the like. When orally administered, the HMG-CoA RI is typically administered at a dose of 0.05 to 200 mg/day, and preferably 0.1 to 100 mg/day in a single dose or in two divided doses. If necessary, the total dose may be administered in several divided doses. The dose of the HMG-CoA RI may be reduced depending on the dose of the EPA-E.

It is to be noted that pravastatin sodium (MevalotinTM tablets and fine granules, Daiichi Sankyo Co., Ltd.), simvastatin (LipovasTM tablets, Banyu Pharmaceutical Co., Ltd.), fluvastatin sodium (LocholTM Tablets, Novartis Pharma K.K. and Tanabe Seiyaku Co., Ltd.), atorvastatin calcium hydrate (LipitorTM tablets, Astellas Pharma Inc. and Pfizer Inc.), pitavastatin calcium (LivaloTM, Kowa Company, Ltd. and Daiichi Sankyo Co., Ltd.), and rosuvastatin calcium (CrestorTM tablets, AstraZeneca and Shionogi & Co., Ltd.) are commercially available in Japan as drugs for treating hyperlipidemia, and lovastatin (MevacorTM tablets, Merck) is commercially available in the U.S. as a drug for treating hyperlipidemia. These drugs may be purchased and used according to the prescribed dosing schedules.

In the case of pravastatin sodium, the preferable daily dose is 5 to 60 mg, and more preferably 10 to 20 mg, and in the case of simvastatin, the preferable daily dose is 2.5 to 60 mg, and more preferably 5 to 20 mg. In the case of fluvastatin sodium, the preferable daily dose is 10 to 180 mg, and more preferably 20 to 60 mg, and in the case of atorvastatin calcium hydrate, the preferable daily dose is 5 to 120 mg, and more preferably 10 to 40 mg. In the case of pitavastatin calcium, the preferable daily dose is 0.5 to 12 mg, and more preferably 1 to 4 mg, and in the case of rosuvastatin calcium, the preferable daily dose is 1.25 to 60 mg, and more preferably 2.5 to 20 mg. In the case of lovastatin, the preferable daily dose is 5 to 160 mg, and more preferably 10 to 80 mg, and in the case of cerivastatin sodium, the preferable daily dose is 0.075 to 0.9 mg, and more preferably 0.15 to 0.3 mg. The dose, however, is not limited to those as described above.

US 9,700,537 B2

13

EXAMPLES

Next, the effects of the composition of the present invention are demonstrated by referring to Examples, which by no means limit the scope of the present invention.

Example 1: Effect of the EPA-E in Preventing Occurrence of Cardiovascular Events in Patients Having Multiple Risk Factors

Trial Procedure

This trial corresponds to a partial analysis of the results obtained in JELIS (Japan EPA Lipid Intervention Study) which is a large scale clinical trial of high purity EPA preparation which was presented in The Heart Failure Society of America 2005 Annual Meeting (for general information on JELIS, see Medical Tribune, issue of Nov. 17, 2005, Feature article 3, pp. 75-76).

More specifically, for the EPA-E group (7503 cases) and the control group (7478 cases) evaluated for the primary prevention effect in the 18,645 subject patients of the DELIS trial (EPA-E group (9,326 cases) and control group (9,319 cases)), occurrence of the cardiovascular events was observed and analysed for 5 years from the start of the administration in relation to the number of risk factors at the registration as defined by the following (1) to (4):

- (1) obesity: body mass index (BMI) of at least 25;
- (2) hypertension or prehypertension: systolic blood pressure (SBP) of at least 140 mmHg or diastolic blood pressure (DBP) of at least 90 mmHg;
- (3) diabetes, prediabetes, or abnormal glucose tolerance: fasting blood glucose (FBS) of at least 126 mg/dL or hemoglobin A1c (HbA1c) of at least 6.5%;
- (4) hypertriglyceridemia or low HDL cholesterolemia: a triglyceride (TG) of at least 150 mg/dL or a HDL-C of less than 40 mg/dL.

The EPA-E group was orally administered with Epadel (Mochida Pharmaceutical Co., Ltd.) typically at an adult dose of 600 mg per administration and 3 times a day after the meal. However, in the case of abnormal serum TG, the dose could be increased to 900 mg per administration and 3 times a day. In both groups, pravastatin sodium (Mevalotin™ tablets and fine granules, Daiichi Sankyo Co., Ltd.), simvastatin (Lipovas™ tablets, Banyu Pharmaceutical Co., Ltd.), or atorvastatin calcium hydrate (Lipitor™ tablets, Astellas Pharma Inc. and Pfizer Inc.) was used for the base drug, and these drugs were orally administered according to the prescribed dosage regimen. More specifically, pravastatin sodium was orally administered at a daily dose of 10 to 20 mg in a single dose or two divided doses; simvastatin was orally administered at a daily dose of 5 to 20 mg in a single dose; atorvastatin calcium hydrate was orally administered at a daily dose of 10 to 40 mg in a single dose.

Results

The number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group are shown in Table 1 for each number of risk factors. The rate of suppression of the incidence of the cardiovascular events was calculated by the formula: $\{(\text{incidence in the control group}) - (\text{incidence in the EPA-E group})\} / \text{incidence in the control group} \times 100$.

14

TABLE 1

Number of risk factors	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of Suppression (%)
0	14/1309 (1.1)	11/1326 (0.8)	22
1	29/2424 (1.2)	25/2468 (1.0)	15
2	46/2324 (2.0)	34/2238 (1.5)	23
3	29/1205 (2.4)	28/1229 (2.3)	5
4	9/216 (4.2)	6/242 (2.5)	40
1-2	75/4748 (1.6)	59/4706 (1.3)	21
1-3	104/5953 (1.7)	87/5935 (1.5)	18
1-4	113/6169 (1.8)	93/6177 (1.5)	18
2-3	75/3529 (2.1)	62/3467 (1.8)	16
2-4	84/3745 (2.2)	68/3709 (1.8)	18
3-4	38/1421 (2.7)	34/1471 (2.3)	14

The incidence (%) of cardiovascular events was found to increase with the increase in the number of risk factors. While the incidence was 1.1% for the risk factor number of 0 and 4.2% for the risk factor number 4 in the control group, the incidence was 0.8% in the risk factor number 0 and 2.5% for the case of risk factor number 4 in the group administered with the EPA-E. As evident from Table 1, for all cases of both groups with 1 to 4 risk factors, the cardiovascular event incidence was lower in the group administered with the EPA-E compared to the control group, and the cardiovascular events were suppressed by 5 to 40%. The effect of preventing occurrence of the cardiovascular events by the administration of the EPA-E was thereby confirmed for the hypercholesterolemia patients having the risk factors.

From the results of the trial as described above, the number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group were calculated for the male patients having at least two risk factors. The results are shown in Table 2. (The calculation was conducted by the same procedure as described above.)

TABLE 2

Number of risk factors	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of Suppression (%)
2-4	43/1053 (4.1)	19/1065 (1.8)	56

FIG. 1 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and time after the start of the trial in X-axis.

As evident from Table 2 and FIG. 1, in the case of male patients having two or more risk factors, EPA-E significantly suppressed the occurrence of cardiovascular events. It was also confirmed that decrease in the incidence of the cardiovascular events was significant after 2 years or more from

US 9,700,537 B2

15

the start of the administration. At the end of the trial, the rate of suppression of the cardiovascular event occurrence was 56% compared to the control group (the value after correcting the dispersion between groups was 55%; see FIG. 1).

From the results of the trial as described above, the number of occurrence of cardiovascular events in the observation period of 5 years, incidence (%), and rate of suppression of the incidence of the cardiovascular events in the EPA-E group with respect to the control group were calculated for the patients exhibiting a triglyceride (TG) of at least 150 mg/dL and a HDL-C of less than 40 mg/dL as the risk factors. The results are shown in Table 3. (The calculation was conducted by the same procedure as described above.)

TABLE 3

Risk factor	Incidence in the control group (cases of occurrence/ all cases, %)	Incidence in the EPA-E group (cases of occurrence/ all cases, %)	Rate of Suppression (%)
TG of at least 150 mg/dL and HDL-C of less than 40 mg/dL	21/475 (4.4)	11/482 (2.3)	48

FIG. 2 is a graph prepared by plotting the incidence of the cardiovascular events in Y-axis and time after the start of the trial in X-axis.

As evident from Table 3 and FIG. 2, EPA-E significantly suppressed occurrence of cardiovascular events in the patients having the risk factors of the triglyceride (TG) of at least 150 mg/dL and the HDL-C of less than 40 mg/dL. It was also confirmed that decrease in the incidence of the cardiovascular events was significant after 2 years or more from the start of the administration. At the end of the trial, the rate of suppression of the cardiovascular event occurrence was 48% compared to the control group (the value after correcting the dispersion between groups was 53%; see FIG. 2). This suggests that the composition containing EPA-E as its effective component effectively prevents the occurrence of the cardiovascular event in the patient having the serum TG/HDL-C ratio of at least 3.75. It is also to be noted that, while the events that occurred in the control group were fatal myocardial infarction, nonfatal myocardial infarction, new occurrence of angina and cardiovascular angioplasty, the events that occurred in the EPA-E group were either nonfatal myocardial infarction or new occurrence of angina, and occurrence of fatal events was not found in the EPA-E group.

In addition, in the group of patients having the risk factor of the triglyceride of at least 150 mg/dL, the occurrence of the cardiovascular events was suppressed by 15% in the EPA-E group compared to the control group; and in the group of patients having the risk factor of HDL-C of less than 40 mg/dL, the occurrence of the cardiovascular events was suppressed by 35% compared to the control group (both values are uncorrected values).

As described above, a significant effect of the EPA-E administration was confirmed for the prevention of the occurrence of the cardiovascular events in the hypercholesterolemia patients having the risk factors.

What is claimed:

1. A method of reducing occurrence of a cardiovascular event in a hypercholesterolemia patient consisting of: identifying a patient having triglycerides (TG) of at least 150 mg/dL and HDL-C of less than 40 mg/dL in a

16

blood sample taken from the patient as a risk factor of a cardiovascular event, wherein the patient has not previously had a cardiovascular event, and administering ethyl icosapentate in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor,

wherein said 3-hydroxyl-3-methylglutaryl coenzyme A reductase inhibitor is administered to the patient at least one of before, during and after administering the ethyl icosapentate; and

wherein the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor is selected from the group consisting of pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, rosuvastatin, and salts thereof, and

wherein daily dose of the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor are 5 to 60 mg for pravastatin, 2.5 to 60 mg for simvastatin, 10 to 180 mg for fluvastatin sodium, 5 to 120 mg for atorvastatin calcium hydrate, 0.5 to 12 mg for pitavastatin calcium, 1.25 to 60 mg for rosuvastatin calcium, 5 to 160 mg for lovastatin, and 0.075 to 0.9 mg for cerivastatin sodium.

2. The method according to claim 1, wherein the ethyl icosapentate is orally administered at a dose of 1.8 g/day to 2.7 g/day.

3. The method according to claim 1, wherein the hypercholesterolemia patient is a male patient.

4. The method according to claim 1, wherein the ethyl icosapentate is administered daily for two years or more.

5. The method according to claim 1, wherein the cardiovascular event is a fatal cardiovascular event.

6. The method according to claim 1, wherein the hypercholesterolemia patient has a serum [triglyceride (TG)/HDL-C] ratio of at least 3.75.

7. The method according to claim 1, wherein the ethyl icosapentate is orally administered at a dose of 0.3 g/day to 6 g/day.

8. The method according to claim 1, wherein the content of the ethyl icosapentate is at least 96.5% by weight in relation to the total content of fatty acid that is simultaneously administered with the ethyl icosapentate.

9. A method of reducing occurrence of a cardiovascular event in a hypercholesterolemia patient consisting of:

identifying a patient having (i) total cholesterol (TC) of at least 220 mg/dL or LDL-cholesterol (LDL-C) of at least 140 mg/dL, and (ii) triglycerides (TG) of at least 150 mg/dL and HDL-C of less than 40 mg/dL in a blood sample taken from the patient as a risk factors of a cardiovascular event, wherein the patient has not previously had a cardiovascular event, and

administering ethyl icosapentate in combination with a 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor,

wherein said 3-hydroxyl-3-methylglutaryl coenzyme A reductase inhibitor is administered to the patient at least one of before, during and after administering the ethyl icosapentate; and

wherein the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor is selected from the group consisting of pravastatin, lovastatin, simvastatin, fluvastatin, atorvastatin, pitavastatin, rosuvastatin, and salts thereof, and

wherein daily dose of the 3-hydroxy-3-methylglutaryl coenzyme A reductase inhibitor are 5 to 60 mg for pravastatin, 2.5 to 60 mg for simvastatin, 10 to 180 mg for fluvastatin sodium, 5 to 120 mg for atorvastatin calcium hydrate, 0.5 to 12 mg for pitavastatin calcium,

US 9,700,537 B2

17

18

1.25 to 60 mg for rosuvastatin calcium, 5 to 160 mg for lovastatin, and 0.075 to 0.9 mg for cerivastatin sodium.

10. The method according to claim 9, wherein the ethyl icosapentate is orally administered at a dose of 1.8 g/day to 2.7 g/day.

5

11. The method according to claim 9, wherein the hypercholesterolemia patient is a male patient.

12. The method according to claim 9, wherein the ethyl icosapentate is administered daily for two years or more.

13. The method according to claim 9, wherein the cardiovascular event is a fatal cardiovascular event.

10

14. The method according to claim 9, wherein the hypercholesterolemia patient has a serum [triglyceride (TG)/HDL-C] ratio of at least 3.75.

15. The method according to claim 9, wherein the ethyl icosapentate is orally administered at a dose of 0.3 g/day to 6 g/day.

15

16. The method according to claim 9, wherein the content of the ethyl icosapentate is at least 96.5% by weight in relation to the total content of fatty acid that is simultaneously administered with the ethyl icosapentate.

20

* * * * *

(12) **United States Patent**
Soni(10) **Patent No.:** **US 10,568,861 B1**(45) **Date of Patent:** ***Feb. 25, 2020**(54) **METHODS OF REDUCING THE RISK OF A CARDIOVASCULAR EVENT IN A SUBJECT AT RISK FOR CARDIOVASCULAR DISEASE**(71) Applicant: **Amarin Pharmaceuticals Ireland Limited, Dublin (IE)**(72) Inventor: **Paresh Soni, Mystic, CT (US)**(73) Assignee: **Amarin Pharmaceuticals Ireland Limited (IE)**

(*) Notice: Subject to any disclaimer, the term of this patent is extended or adjusted under 35 U.S.C. 154(b) by 0 days.

This patent is subject to a terminal disclaimer.

(21) Appl. No.: **16/599,374**(22) Filed: **Oct. 11, 2019****Related U.S. Application Data**

(60) Division of application No. 16/502,621, filed on Jul. 3, 2019, which is a continuation of application No. 16/287,157, filed on Feb. 27, 2019, now Pat. No. 10,383,840, which is a continuation of application No. 16/005,852, filed on Jun. 12, 2018, now Pat. No. 10,278,935, which is a continuation of application No. 15/886,422, filed on Feb. 1, 2018, now Pat. No. 10,016,386, which is a continuation of application No. 15/607,084, filed on May 26, 2017, now Pat. No. 9,918,955, which is a continuation of application No. 15/427,238, filed on Feb. 8, 2017, now Pat. No. 9,693,986, which is a continuation of application No. 15/333,991, filed on Oct. 25, 2016, now Pat. No. 9,610,272, which is a continuation of application No. 14/411,815, filed as application No. PCT/US2013/048559 on Jun. 28, 2013, now Pat. No. 9,603,826.

(60) Provisional application No. 61/666,447, filed on Jun. 29, 2012.

(51) **Int. Cl.****A61K 31/232** (2006.01)
A61K 31/397 (2006.01)
A61K 45/06 (2006.01)(52) **U.S. Cl.**CPC **A61K 31/232** (2013.01); **A61K 31/397** (2013.01); **A61K 45/06** (2013.01); **A61K 2300/00** (2013.01)(58) **Field of Classification Search**CPC **A61K 31/232**; **A61K 31/397**; **A61K 45/06**; **A61K 2300/00**
USPC **548/400**
See application file for complete search history.(56) **References Cited****U.S. PATENT DOCUMENTS**4,377,526 A 3/1983 Fujita et al.
4,526,902 A 7/1985 Rubin4,920,098 A 4/1990 Cotter et al.
4,935,243 A 6/1990 Borkan et al.
5,013,443 A 5/1991 Higashidate et al.
5,116,871 A 5/1992 Horrobin et al.
5,178,873 A 1/1993 Horrobin et al.
5,198,468 A 3/1993 Horrobin
5,215,630 A 6/1993 Hata et al.
5,252,333 A 10/1993 Horrobin
5,343,389 A 8/1994 Otvos
5,385,929 A 1/1995 Bjorge et al.
5,457,130 A 10/1995 Tisdale et al.
5,502,077 A 3/1996 Breivik et al.
5,567,730 A 10/1996 Miyashita et al.
5,589,508 A 12/1996 Schlotzer et al.
5,603,959 A 2/1997 Horrobin et al.
5,618,558 A 4/1997 Horrobin et al.
5,656,667 A 8/1997 Breivik et al.
5,698,594 A 12/1997 Breivik et al.
5,760,081 A 6/1998 Leaf et al.
5,763,496 A 6/1998 Holland
5,776,978 A 7/1998 Bruzzese
5,792,795 A 8/1998 Buser et al.
5,837,731 A 11/1998 Vaddadi
5,840,944 A 11/1998 Furihata et al.
5,886,037 A 3/1999 Klor et al.
5,888,541 A 3/1999 Horrobin et al.
5,948,818 A 9/1999 Buser et al.
6,025,008 A 2/2000 Akahoshi
6,069,168 A 5/2000 Horrobin et al.
6,193,999 B1 2/2001 Gennadios
6,284,268 B1 9/2001 Mishra et al.
6,313,330 B1 11/2001 Kiyohara et al.

(Continued)

FOREIGN PATENT DOCUMENTSCA 2628305 5/2007
CA 2653787 12/2007

(Continued)

OTHER PUBLICATIONSA study of AMR101 to evaluate its ability to reduce cardiovascular events in high risk patients with hypertriglyceridemia and on statin (REDUCE-IT). Available at: <http://clinicaltrials.gov/show/NCT01492361>. (3 pages), 2011.

(Continued)

Primary Examiner — Kristin A Vajda

(74) Attorney, Agent, or Firm — Perkins Coie LLP

(57) **ABSTRACT**

In various embodiments, the present invention provides methods of reducing the risk of a cardiovascular event in a subject on statin therapy and, in particular, a method of reducing the risk of a cardiovascular event in a subject on statin therapy having a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL, and administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of eicosapentaenoic acid ethyl ester or a derivative thereof.

7 Claims, No Drawings

US 10,568,861 B1

Page 2

(56)	References Cited				2002/0077361	A1	6/2002	Peet et al.
	U.S. PATENT DOCUMENTS				2002/0169209	A1	11/2002	Horrobin
					2002/0183389	A1	12/2002	Peet
					2002/0193439	A1	12/2002	Peet et al.
6,326,031	B1	12/2001	Hsia et al.		2002/0198177	A1	12/2002	Horrobin et al.
6,326,355	B1	12/2001	Abbruzzese et al.		2003/0100610	A1	5/2003	Shibuya
6,331,568	B1	12/2001	Horrobin		2003/0104048	A1	6/2003	Patel et al.
6,368,621	B1	4/2002	Engel et al.		2003/0161918	A1	8/2003	Kendrick et al.
6,384,077	B1	5/2002	Peet		2003/0166614	A1	9/2003	Harrison
6,479,544	B1	11/2002	Horrobin		2003/0232385	A1	12/2003	Breit et al.
6,482,421	B2	11/2002	Weidner		2004/0009208	A1	1/2004	Edson
6,531,150	B1	3/2003	Sunohara et al.		2004/0048919	A1	3/2004	Dreon et al.
6,555,700	B1	4/2003	Horrobin et al.		2004/0062847	A1	4/2004	Koiki et al.
6,596,766	B1	7/2003	Igarashi et al.		2004/0077723	A1	4/2004	Granata
6,620,821	B2	9/2003	Robl		2004/0106591	A1	6/2004	Pacioretty et al.
6,689,812	B2	2/2004	Peet		2004/0121000	A1	6/2004	Bowe et al.
6,846,942	B2	1/2005	Rubin		2004/0162348	A1	8/2004	Peet et al.
7,022,713	B2	4/2006	Aoki et al.		2004/0082402	A1	9/2004	Troup et al.
7,112,609	B2	9/2006	Hermelin et al.		2004/0204356	A1	10/2004	Guenzler-Pukall
7,119,118	B2	10/2006	Peet		2004/0258645	A1	12/2004	Trejo et al.
7,179,491	B1	2/2007	Mag		2005/0042214	A1	2/2005	Gershwin et al.
7,205,329	B2	4/2007	Chien et al.		2005/0137253	A1	6/2005	Phinney et al.
7,405,302	B2	7/2008	Hutchinson et al.		2005/0147665	A1	7/2005	Horrobin et al.
7,498,359	B2	3/2009	Yokoyama et al.		2005/0187292	A1	8/2005	Aoki et al.
7,511,131	B2	3/2009	Crooke et al.		2005/0244367	A1	11/2005	Hui et al.
7,598,227	B2	10/2009	Crooke et al.		2005/0272095	A1	12/2005	Wang
7,776,881	B2	8/2010	Aoki et al.		2006/0034815	A1	2/2006	Guzman et al.
8,188,146	B2	5/2012	Peet et al.		2006/0051418	A1	3/2006	Cowen et al.
8,293,727	B2	10/2012	Manku et al.		2006/0088502	A1	4/2006	Sata et al.
8,293,728	B2	10/2012	Manku et al.		2006/0111437	A1	5/2006	Aoki et al.
8,298,554	B2	10/2012	Manku		2006/0134178	A1	6/2006	Doisaki et al.
8,314,086	B2	11/2012	Manku et al.		2006/0134206	A1	6/2006	Iyer et al.
8,318,715	B2	11/2012	Manku et al.		2006/0135607	A1	6/2006	Kobayashi et al.
8,324,195	B2	12/2012	Manku et al.		2006/0135610	A1	6/2006	Bortz et al.
8,357,677	B1	1/2013	Manku et al.		2006/0141022	A1	6/2006	Kawamura et al.
8,367,652	B2	2/2013	Manku et al.		2006/0142390	A1	6/2006	Manku et al.
8,377,920	B2	2/2013	Manku et al.		2006/0172012	A1	8/2006	Finley et al.
8,410,086	B2	4/2013	Osterloh et al.		2006/0189682	A1	8/2006	Payne et al.
8,431,560	B1	4/2013	Manku et al.		2006/0211749	A1	9/2006	Bobotas et al.
8,440,650	B1	5/2013	Manku et al.		2006/0211761	A1	9/2006	Kumar et al.
8,455,472	B2	6/2013	Osterloh et al.		2006/0211762	A1	9/2006	Rongen
8,518,929	B2	8/2013	Manku et al.		2006/0211763	A1	9/2006	Fawzy et al.
8,524,698	B2	9/2013	Manku et al.		2006/0217356	A1	9/2006	Wright et al.
8,546,372	B2	10/2013	Manku et al.		2006/0223838	A1	10/2006	Jiang
8,551,521	B2	10/2013	Manku et al.		2006/0252833	A1	11/2006	Peet et al.
8,563,608	B2	10/2013	Manku et al.		2007/0021504	A1	1/2007	Yokoyama et al.
8,617,593	B2	12/2013	Manku et al.		2007/0060532	A1	3/2007	Junien et al.
8,617,594	B2	12/2013	Manku et al.		2007/0098787	A1	5/2007	Kakiuchi
8,618,168	B2	12/2013	Fujii et al.		2007/0104779	A1	5/2007	Rongen et al.
8,623,406	B2	1/2014	Manku et al.		2007/0105793	A1	5/2007	Hendrix
8,642,077	B2	2/2014	Manku et al.		2007/0105954	A1	5/2007	Puri
8,669,245	B2	3/2014	Osterloh et al.		2007/0141138	A1	6/2007	Feuerstein et al.
8,680,144	B2	3/2014	Osterloh et al.		2007/0167520	A1	7/2007	Bruzzese
8,691,871	B2	4/2014	Osterloh et al.		2007/0185198	A1	8/2007	Yokoyama et al.
8,703,185	B2	4/2014	Manku et al.		2007/0191467	A1	8/2007	Rongen et al.
8,709,475	B2	4/2014	Manku et al.		2007/0202159	A1	8/2007	Mathur et al.
8,906,964	B2	12/2014	Bobotas et al.		2007/0212411	A1	9/2007	Fawzy et al.
9,006,285	B2	4/2015	Ohnishi		2007/0219271	A1	9/2007	Mittmann et al.
9,060,981	B2	6/2015	Sato et al.		2007/0265340	A1	11/2007	Shalwitz et al.
9,138,415	B2	9/2015	Manku et al.		2007/0269507	A1	11/2007	Sachetto et al.
9,452,121	B2	9/2016	Manku et al.		2008/0020018	A1	1/2008	Moodley et al.
9,452,150	B2	9/2016	Ueshima et al.		2008/0057115	A1	3/2008	Okamoto
9,603,826	B2	3/2017	Soni		2008/0085911	A1	4/2008	Rongen et al.
9,610,272	B2	4/2017	Soni		2008/0089876	A1	4/2008	Cavazza
9,623,001	B2	4/2017	Soni		2008/0113046	A1	5/2008	Gardette
9,693,984	B2	7/2017	Soni		2008/0125490	A1	5/2008	Svensson et al.
9,693,985	B2	7/2017	Soni		2008/0139604	A1	6/2008	Fitzpatrick et al.
9,693,986	B2	7/2017	Soni		2008/0185198	A1	8/2008	Jones
9,855,237	B2	1/2018	Osterloh et al.		2008/0200453	A1	8/2008	Cincotta
9,918,954	B2	3/2018	Soni		2008/0200547	A1	8/2008	Peet et al.
10,058,521	B2	8/2018	Bobotas et al.		2008/0200707	A1	8/2008	Shimano et al.
10,166,209	B2	1/2019	Manku et al.		2008/0214531	A1	9/2008	Saxena
10,265,290	B2	4/2019	Manku et al.		2008/0299187	A1	12/2008	Opheim et al.
2001/0035125	A1	11/2001	Talieh et al.		2008/0306154	A1	12/2008	Svensson et al.
2002/0016312	A1	2/2002	Seed et al.		2008/0319077	A1	12/2008	Suzuki et al.
2002/0025983	A1	2/2002	Horrobin		2009/0012167	A1	1/2009	Rongen et al.
2002/0035125	A1	3/2002	Shear		2009/0018125	A1	1/2009	Mittmann et al.
2002/0055529	A1	5/2002	Bisgaier et al.		2009/0042979	A1	2/2009	Guzman et al.
2002/0055539	A1	5/2002	Bockow et al.		2009/0054329	A1	2/2009	Willemsen et al.

US 10,568,861 B1

Page 3

(56)	References Cited		2013/0079409 A1	3/2013	Manku et al.
	U.S. PATENT DOCUMENTS		2013/0090383 A1	4/2013	Manku et al.
			2013/0095178 A1	4/2013	Manku
			2013/0095179 A1	4/2013	Davidson et al.
2009/0105340 A1	4/2009	Yokoyama	2013/0096197 A1	4/2013	Manku
2009/0148543 A1	6/2009	Theoharides	2013/0102674 A1	4/2013	Manku
2009/0156675 A1	6/2009	Yokoyama et al.	2013/0115284 A1	5/2013	Fujii
2009/0182049 A1	7/2009	Opheim	2013/0131170 A1	5/2013	Manku
2009/0227602 A1	9/2009	Griffin et al.	2013/0156852 A1	6/2013	Manku et al.
2009/0233843 A1	9/2009	Marin	2013/0158120 A1	6/2013	Manku et al.
2009/0239927 A1	9/2009	Bobotas et al.	2013/0164375 A1	6/2013	Manku et al.
2009/0304784 A1	12/2009	Mane et al.	2013/0165513 A1	6/2013	Manku et al.
2009/0311322 A1	12/2009	Dlugatch et al.	2013/0171249 A1	7/2013	Manku et al.
2010/0021555 A1	1/2010	Geiringer et al.	2013/0171250 A1	7/2013	Manku et al.
2010/0063018 A1	3/2010	Pellicciari et al.	2013/0171251 A1	7/2013	Manku et al.
2010/0069492 A1	3/2010	Geiringer et al.	2013/0172413 A1	7/2013	Manku
2010/0113506 A1	5/2010	Kawano et al.	2013/0189355 A1	7/2013	Manku et al.
2010/0113811 A1	5/2010	Yadav et al.	2013/0195972 A1	8/2013	Manku et al.
2010/0119598 A1	5/2010	Yoshinari et al.	2013/0252989 A1	9/2013	Manku et al.
2010/0130608 A1	5/2010	Ryan et al.	2013/0252990 A1	9/2013	Manku et al.
2010/0160261 A1	6/2010	Fortin	2013/0253030 A1	9/2013	Osterloh et al.
2010/0233280 A1	9/2010	Driscoll	2013/0253031 A1	9/2013	Osterloh et al.
2010/0254951 A1	10/2010	Shido et al.	2013/0260403 A1	10/2013	Button et al.
2010/0278879 A1	11/2010	Manku	2013/0261180 A1	10/2013	Gillies et al.
2010/0285121 A1	11/2010	Uchiyama et al.	2013/0281534 A1	10/2013	Osterloh et al.
2010/0298379 A1	11/2010	Jacobsen	2013/0295173 A1	11/2013	Machielse et al.
2010/0305205 A1	12/2010	Yokoyama et al.	2013/0303614 A1	11/2013	Kanehiro et al.
2010/0311834 A1	12/2010	Manku et al.	2013/0324607 A1	12/2013	Mason
2011/0034555 A1	2/2011	Osterloh et al.	2013/0331447 A1	12/2013	Manku et al.
2011/0065793 A1	3/2011	Peet et al.	2014/0004183 A1	1/2014	Soni et al.
2011/0071176 A1	3/2011	Rowe	2014/0005264 A1	1/2014	Soni et al.
2011/0082119 A1	4/2011	Yano	2014/0005265 A1	1/2014	Soni et al.
2011/0092592 A1	4/2011	Yano	2014/0017306 A1	1/2014	Manku
2011/0105510 A1	5/2011	Ishikawa	2014/0057981 A1	2/2014	Fujii
2011/0130458 A1	6/2011	Breivik et al.	2014/0073692 A1	3/2014	Peet
2011/0178105 A1	7/2011	Gillies et al.	2014/0080850 A1	3/2014	Mason
2011/0195061 A1	8/2011	Minatelli	2014/0080909 A1	3/2014	Manku
2011/0218243 A1	9/2011	Rowe	2014/0088194 A1	3/2014	Manku
2011/0223158 A1	9/2011	Sacks et al.	2014/0094520 A1	4/2014	Bobotas et al.
2011/0236476 A1	9/2011	Manku	2014/0107199 A1	4/2014	Fawzy et al.
2011/0268811 A1	11/2011	Minatelli et al.	2014/0127289 A1	5/2014	Osterloh et al.
2011/0288171 A1	11/2011	Manku et al.	2014/0128453 A1	5/2014	Mullick et al.
2012/0035105 A1	2/2012	Geho et al.	2014/0128464 A1	5/2014	Rowe
2012/0035262 A1	2/2012	Osterloh et al.	2014/0154310 A1	6/2014	Osterloh et al.
2012/0039997 A1	2/2012	Manku et al.	2014/0155455 A1	6/2014	Osterloh et al.
2012/0046251 A1	2/2012	Schaefer et al.	2014/0155481 A1	6/2014	Osterloh et al.
2012/0093922 A1	4/2012	Manku et al.	2014/0186438 A1	7/2014	Manku et al.
2012/0093924 A1	4/2012	Manku et al.	2014/0187633 A1	7/2014	Manku et al.
2012/0100208 A1	4/2012	Manku	2014/0213648 A1	7/2014	Manku et al.
2012/0108659 A1	5/2012	Manku et al.	2014/0221358 A1	8/2014	Zakrzewski
2012/0108660 A1	5/2012	Manku et al.	2014/0221452 A1	8/2014	Zakrzewski
2012/0108663 A1	5/2012	Manku et al.	2014/0221486 A1	8/2014	Manku et al.
2012/0121698 A1	5/2012	Manku et al.	2014/0221676 A1	8/2014	Braeckman et al.
2012/0156285 A1	6/2012	Manku et al.	2014/0234410 A1	8/2014	Moodley et al.
2012/0157530 A1	6/2012	Manku et al.	2014/0235716 A1	8/2014	Manku et al.
2012/0157531 A1	6/2012	Osterloh et al.	2014/0243389 A1	8/2014	Zakrzewski
2012/0172432 A1	7/2012	Manku et al.	2014/0249200 A1	9/2014	Braeckman et al.
2012/0184595 A1	7/2012	Macdonald et al.	2014/0249214 A1	9/2014	Braeckman et al.
2012/0195963 A1	8/2012	Peet et al.	2014/0249220 A1	9/2014	Braeckman et al.
2012/0207800 A1	8/2012	Abu-Baker	2014/0249225 A1	9/2014	Mason
2012/0214771 A1	8/2012	Sampalis	2014/0256809 A1	9/2014	Zakrzewski
2012/0225120 A1	9/2012	Manku et al.	2014/0271841 A1	9/2014	Grandolfi
2012/0232145 A1	9/2012	Osterloh et al.	2014/0271907 A1	9/2014	Zakrzewski
2012/0237594 A1	9/2012	Manku et al.	2014/0275252 A1	9/2014	Zakrzewski
2012/0264824 A1	10/2012	Mizuguchi et al.	2014/0275253 A1	9/2014	Zakrzewski
2012/0295976 A1	11/2012	Yokoyama	2014/0357717 A1	12/2014	Braeckman et al.
2012/0302589 A1	11/2012	Manku et al.	2014/0364459 A1	12/2014	Zakrzewski
2012/0329852 A1	12/2012	Yokoyama	2015/0045431 A1	2/2015	Zakrzewski
2013/0004566 A1	1/2013	Manku et al.	2015/0051143 A1	2/2015	Harada et al.
2013/0004567 A1	1/2013	Manku et al.	2015/0051282 A1	2/2015	Zakrzewski
2013/0004568 A1	1/2013	Manku et al.	2015/0065572 A1	3/2015	Zakrzewski
2013/0004572 A1	1/2013	Manku et al.	2015/0073050 A1	3/2015	Zakrzewski
2013/0005757 A1	1/2013	Osterloh et al.	2015/0141510 A1	5/2015	Kiyohara et al.
2013/0005809 A1	1/2013	Manku et al.	2015/0157592 A1	6/2015	Soni
2013/0011471 A1	1/2013	Manku et al.	2015/0157593 A1	6/2015	Braeckman et al.
2013/0011472 A1	1/2013	Manku et al.	2015/0164850 A1	6/2015	Osterloh et al.
2013/0012580 A1	1/2013	Osterloh et al.	2015/0190361 A1	7/2015	Osterloh et al.
2013/0017256 A1	1/2013	Manku et al.	2015/0216831 A1	8/2015	Manku et al.
2013/0065956 A1	3/2013	Yokoyama	2015/0250754 A1	9/2015	Ohta

Page 4

CA	2675836	7/2008
CA	2724983	11/2009
CA	2772378	12/2010
CN	101252837	8/2008
EP	273708	7/1988
EP	277747	8/1988
EP	0302482	2/1989
EP	347509	12/1989
EP	0460917	12/1991
EP	606012	7/1994
EP	0610506	8/1994
EP	0641562	3/1995
EP	0843972	5/1998
EP	1125914	8/2001
EP	1157692	11/2001
EP	1296670	4/2003

US 10,568,861 B1

Page 5

(56) References Cited

FOREIGN PATENT DOCUMENTS

WO	WO 2010/127103	11/2010
WO	WO 2010/147994	12/2010
WO	WO2011/028689	3/2011
WO	WO 2011/038122	3/2011
WO	WO2011/085211	7/2011
WO	WO 2011/109724	9/2011
WO	WO 2012/074930	6/2012
WO	WO2012/128587	9/2012
WO	WO 2013/070735	5/2013
WO	WO2013/103958	7/2013
WO	WO2013/148136	10/2013
WO	WO2014/004861	1/2014
WO	WO2014/004993	1/2014
WO	WO2014/005013	1/2014
WO	WO 2014/057522	4/2014
WO	WO2014/074552	5/2014
WO	WO2014/130200	8/2014
WO	WO2014/134466	9/2014
WO	WO2014/143469	9/2014
WO	WO2014/143523	9/2014
WO	WO2015/021141	2/2015
WO	WO2015/066512	5/2015
WO	WO2015/195662	12/2015
WO	WO2016/140949	9/2016

OTHER PUBLICATIONS

Aarsetoey H, Gurndt H, Nygaard O. The Role of Long-Chained Marine N-3 Polyunsaturated Fatty Acids in Cardiovascular Disease. *Cardiol Res Pract.* 2012. Epub Dec. 13, 2012.

Aarsland, et al., "On the Effect of Peroxisomal beta-Oxidation and Carnitine Palmitoyltransferase Activity by Eicosapentaenoic Acid in Liver and Heart of Rats." *Lipids*, 25:546-548, (Sep. 1990).

Aas, V., et al., "Eicosapentaenoic acid (20:5 n-3) increases fatty acid and glucose uptake in cultured human skeletal muscle cells." *Journal of Lipid Research*, 47:366-374 (Feb. 2006).

Abbey, M., et al., "Effect of fish oil on lipoproteins, lecithin:cholesterol acyltransferase, and lipid transfer protein activity in humans." *Arterioscler. Thromb. Vasc. Biol.* 10:85-94 (Jan./Feb. 1990).

Abela GS, Aziz K. "Cholesterol crystals cause mechanical damage to biological membranes: a proposed mechanism of plaque rupture and erosion leading to arterial thrombosis." *Clin. Cardiol.* (Sep. 2005);28(9):413-420.

Abelo A, Andersson TB, Antonsson M, et al. "Stereoselective metabolism of omeprazole by human cytochrome P450 enzymes." *Drug Metab. Dispos.* Aug. 28, 2000 (8): 966-72.

Ackman et al., "The 'Basic' Fatty Acid Composition of Atlantic Fish Oils: Potential Similarities Useful for Enrichment of Polyunsaturated Fatty Acids by Urea Complexation," *JAOCS*, vol. 65, 1:136-138 (Jan. 1988).

Adan, Y., et al., "Effects of docosahexaenoic and eicosapentaenoic acid on lipid metabolism, eicosanoid production, platelet aggregation and atherosclerosis." *Biosci. Biotechnol. Biochem.* 63(1), 111-119 (Jan. 1999).

Adan, Y., et al., "Concentration of serum lipids and aortic lesion size in female and male apo E-deficient mice fed docosahexaenoic acid." *Biosci. Biotechnol. Biochem.* 63(2):309-313 (Feb. 1999).

Adorini et al., "Farnesoid X receptor targeting to treat nonalcoholic steatohepatitis." *Drug Discover Today*, 14(17-18):988-997 (Sep. 2012)(available online May 28, 2012).

Agren JJ, Vaisanen S, Hanninen O, et al. "Hemostatic factors and platelet aggregation after a fish-enriched diet or fish oil or docosahexaenoic acid supplementation." *Prostaglandins Leukot Essent Fatty Acids* (Oct. 1997) 57 (4-5): 419-21.

Agren, J.J., et al., "Fatty acid composition of erythrocyte, platelet, and serum lipids in strict vegans." *Lipids* 30:365-369 (Apr. 1995).

Agren, J.J., et al., "Fish diet, fish oil and docosahexaenoic acid rich oil lower fasting and postprandial plasma lipid levels." *Eur J Clin Nutr.* 50:765-771. (Nov. 1996).

Aguilar-Salinas et al., "High Prevalence of Low HDL Cholesterol Concentrations and Mixed Hyperlipidemia in a Mexican Nationwide Survey," *J Lipid Res.*, (Aug. 2001), 42:1298-1307.

Ai M, Ootokozawa S, Asztalos BF, Ito Y, Nakajima K, White CC, Cupples LA, Wilson PW, Schaefer EJ. "Small dense LDL cholesterol and coronary heart disease: results from the Framingham Offspring Study." *Clin. Chem.* (Jun. 2010);56(6):967-976.

Ait-Said, et al., "Inhibition by eicosapentaenoic acid of IL-1 β -induced PGHS-2 expression in human microvascular endothelial cells: involvement of lipoxygenase-derived metabolites and p38 MAPK pathway." *Biochimica et Biophysica Acta*, 1631:66-85 (Feb. 2003).

Albert CM, Campos H, Stampfer MJ, et al. Blood Levels of Long-Chain n-3 Fatty Acids and the Risk of Sudden Death. *N Engl J Med* 346(15):1113-1138, Apr. 2002.

Alberti K, et al. Harmonizing the Metabolic Syndrome: A Joint Interim Statement of the International Diabetes Federation Task Force on Epidemiology and Prevention; National Heart, Lung, and Blood Institute; American Heart Association; World Heart Federation; International Atherosclerosis Society; and International Association for the Study of Obesity. *Circulation.* 120:1640-1645; Oct. 20, 2009.

Alderman, J.D., et al., "Effect of a modified, well-tolerated niacin regimen on serum total cholesterol, high density lipoprotein cholesterol and the cholesterol to high density lipoprotein ratio," *Am. J. Cardio*, 64: 725-729.A (Oct. 1989).

Alessandri, J-M., et al., "Estradiol favors the formation of eicosapentaenoic acid (20:5n-3) and n-3 docosapentaenoic acid (22:5n-3) from alpha-linolenic acid (18:3n-3) in SH-SY5Y neuroblastoma cells." *Lipids* 43:19-28 (Jan. 2008).

Allard et al. "Nutritional assessment and hepatic fatty acid composition in non-alcoholic fatty liver disease (NAFLD): a cross-sectional study." *J Hepatol.* Feb. 2008;48(2):300-7.

Allred, C., et al., "PPAR γ 1 as a molecular target of eicosapentaenoic acid in human colon cancer (HT-29) cells." *J. Nutr.* 138:250-256 (Feb. 2008).

Almeida et al., "Effect of nebicapone on the pharmacokinetics and pharmacodynamics of warfarin in healthy subjects." *Eur J Clin Pharmacol.* (Oct. 2008);64(10):961-6.

Amarin Appoints Medpace as CRO for Two Phase 3 Cardiovascular Trials, published Oct. 19, 2009 (2 pages).

Amarin Corporation Announces First Patients Enrolled in Two Phase 3 Clinical Trials Assessing AMR101 for the Treatment of Cardiovascular Disease [online], Amarin Corporation, Jan. 11, 2010 [retrieved Apr. 27, 2011], Retrieved from the Internet: <<http://investor.amarincorp.com/releasedetail.cfm?ReleaseID=504380>> (2 pages).

Amarin Corporation, Annual Report, Jun. 24, 2010 (245 pages total)(submitted in three parts; Part I: Cover and pp. 1-39 (81 pages); Part II: pp. 40 through F-10 (81 pages); Part III: pp. F11-F51 (83 pages)).

Amarin Corporation, Executive Informational Overview, "Neurological Disease-Focused Biopharmaceutical Opportunity," SEC filing dated Oct. 11, 2005 (99 pages).

Amarin Corporation, Globe Newswire press release, "Reduce-It™ Cardiovascular Outcomes Study of Vascepa® (icosapent ethyl) Capsules Met Primary Endpoint," Sep. 24, 2018 (4 pages).

Amarin Corporation, press release (Jan. 18, 2008)(1 page).

Amarin Presentation "Next Generation Lipid Modification in Cardiovascular Disease," (Aug. 2011)(27 pages).

Amarin Presentation "Next Generation Lipid Modification in Cardiovascular Disease," (Mar. 2010)(25 pages).

Amarin Proceeding to Phase 3 with AMR101 for Hypertriglyceridemia, published Jul. 23, 2008 (1 page).

Amarin, Next Generation Lipid Modification in Cardiovascular Disease, Investor Meetings, Nov. 2010, (http://files.shareholder.com/downloads/AMRN/0x0x417754/AA72705F-1D67-4E1D-A989-5805E5CF0244/Investor_Presentation_2010_Nov_10.pdf, accessed Jan. 6, 2015).

Amarin's Vascepa® Briefing Document for the Endocrinologic and Metabolic Drugs Advisory Committee Meeting dated Oct. 16, 2013, (117 pages).

American Heart Association. Heart Disease and Stroke Statistics—2010 Update. Dallas, Texas: American Heart Association; 2010.

US 10,568,861 B1

Page 6

(56)

References Cited

OTHER PUBLICATIONS

Anand RG, Alkadri M, Lavie CJ, Milani RV. The Role of Fish Oil in Arrhythmia Prevention. *J Cardioplin Rehabil Preven.*, Mar./Apr. 2008; 28:92-98.

Anber V, Griffin BA, McConnell M, Packard CJ, Shepherd J. Influence of plasma lipid and LDL-subfraction profile on the interaction between low density lipoprotein with human arterial wall proteoglycans. *Atherosclerosis*. Aug. 1996;124(2):261-271.

Anderson JL, Adams CD, Antman EM, et al. ACC/AHA 2007 guidelines for the management of patients with unstable angina/non-ST-elevation myocardial infarction—executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to Revise the 2002 Guidelines for the Management of Patients With Unstable Angina/Non-ST-Elevation Myocardial Infarction) developed in Collaboration with the American College of Emergency Physicians, the Society for Cardiovascular Angiography and Interventions, and the Society of Thoracic Surgeons Endorsed by the American Association of Cardiovascular and Pulmonary Rehabilitation and the Society for Academic Emergency Medicine. *J Am Coll Cardiol* 50:652-726, Aug. 14, 2007.

Anderson TJ, Gregoire J, Hegele RA, et al. 2012 update of the Canadian Cardiovascular Society guidelines for the diagnosis and treatment of dyslipidemia for the prevention of cardiovascular disease in the adult. *Can. J. Cardiol.* Feb. 2013;29:151-167.

Anderson TJ, Meredith IT, Yeung AC, Frei B, Selwyn AP, Ganz P. The effect of cholesterol-lowering and antioxidant therapy on endothelium-dependent coronary vasomotion. *N. Engl. J. Med.* Feb. 1995;332:488-493.

Anderson, "Lipoprotein-Associated Phospholipase A2: An Independent Predictor of Coronary Artery Disease Events in Primary and Secondary Prevention," 101 *Am. J. Cardiology* 23F-33F (Jun. 2008).

Ando, M., et al., "Eicosapentaenoic acid reduces plasma levels of remnant lipoproteins and prevents in vivo peroxidation of LDL in dialysis patients." *J. Am. Soc. Nephrol.*, 10:2177-2184 (Oct. 1999).

Ando, Y., et al., "Positional distribution of highly unsaturated fatty acids in triacyl-sn-glycerols of Artemia Nauplii enriched with docosahexaenoic acid ethyl ester." *Lipids* 36:733-740 (Jul. 2001).

Andrade, SE, et al., "Discontinuation of antihyperlipidaemic drugs—do rates reported in clinical trials reflect rates in primary care settings?" *New Eng. J. Med.* 332: 1125-1131. (Apr. 1995).

Andrews HE, Bruckdorfer KR, Dunn RC, Jacobs M. Low-density lipoproteins inhibit endothelium-dependent relaxation in rabbit aorta. *Nature*. May 1987;327:237-239.

Angerer et al., "n-3 Polyunsaturated Fatty Acids and the Cardiovascular System", *Current Opinion in Lipidology*, 11(1):57-63, (Feb. 2000).

Anil, Eliz, "The Impact of EPA and DHA on Blood Lipids and Lipoprotein Metabolism: Influence of ApoE Genotype", *Proceedings of the Nutrition Society*, 66:60-68, (Feb. 2007).

Annex to Rule 161 Response dated Apr. 16, 2012 (4 pages).

Antman E, Anbe D, Armstrong P, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction—executive summary. A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (Writing Committee to revise the 1999 guidelines for the management of patients with acute myocardial infarction). *J Am Coll Cardiol* 44:671-719, Aug. 4, 2004.

Aoki T et al. "Experience of the use of ethyl eicosapentaenoic acid preparation (Epadel) in patients with arteriosclerosis obliterans complicated with diabetes mellitus. A study of the long-term effects on glycemic control and blood lipids," *Rinsho to Kenkyu*; 70:625-631. (1993) (with English translation).

Appendix A to Defendants' Invalidity Contentions, 3:14-CV-02550-MLC-DEA (D.N.J.), 478 pages (Dec. 5, 2014).

Appleton, Katherine M., et al., "Effects of n-3 long-chain polyunsaturated fatty acids on depressed mood: systematic review of published trials", *Am. J. Clin. Nutr.*, 84(6):1308-1316, (Dec. 2006).

Arca et al., "Treating statin-intolerant patients," *Diabetes, Metabolic Syndrome and Obesity: Targets and Therapy*, 4:155-156 (Apr. 28, 2011).

Armaganijan L, Lopes RD, Healey JS, Piccini JP, Nair GN, Morillo CA. Do Omega-3 fatty acids prevent atrial fibrillation after open heart surgery? A meta-analysis of randomized controlled trials. *Clinics*. 2011(accepted for publication Jul. 19, 2011); 66(11):1923-1928.

Arrol, S. et al., "The effects of fatty acids on apolipoprotein B secretion by human hepatoma cells (HEP G2)," *Atherosclerosis* 150:255-264. (Jun. 2000).

Arshad, A., et al., "Sudden cardiac death and the role of medical therapy." *Progress in Cardiovascular Diseases*, vol. 50, No. 6, 420-438, (May/Jun. 2008).

Arterburn, L., et al., "Distribution, interconversion, and dose response of n-3 fatty acids in humans." *Am J Clin Nutr.*, 83:1467S-76S (Jun. 2006).

Asahara, EPA Products What is the Clinical Significance of Epadel? *Obesity and Diabetes* 10(6):903-905 (2011) (with English translation).

Asano, M., et al., "Eicosapentaenoic acid inhibits vasopressin-activated Ca2q influx and cell proliferation in rat aortic smooth muscle cell lines." *European Journal of Pharmacology* 379:199-209 (Aug. 1999).

Asano, M., et al., "Inhibitory effects of ω-3 polyunsaturated fatty acids on receptor-mediated non-selective cation currents in rat A7r5 vascular smooth muscle cells." *British Journal of Pharmacology* 120:1367-1375, (Apr. 1997).

ASCEND Study Collaborative Group. Effects of n-3 fatty acid supplements in diabetes mellitus. *N Engl J Med*, 379(16):1540-1550 (publication date Oct. 18, 2018; epublication dated Aug. 26, 2018).

Ascenta Health "Fish Oil as Triglycerides vs. Ethyl Esters: Why this Matters." (2015)(14 pages).

Astarita et al., "Targeted lipidomics strategies for oxygenated metabolites of polyunsaturated fatty acids," *Biochim Biophys Acta*, 1851(4):456-168 (Apr. 2015).

Atorvastatin Package Leaflet, Reg. No. LSR-005205/08, Sep. 30, 2016 [retrieved Sep. 30, 2016] retrieved from the internet: academ-clinic.ru/drugs/atorvastatin (6 pages).

ATP III guidelines, NIH publication No. 01-3305 (2001).(6 pages).

Attie AD, et al., "Relationship between stearoyl-CoA desaturase activity and plasma triglycerides in human and mouse hypertriglyceridemia," *J. Lipid Res.* 2002;43:1899-907.

Ault, "Prescription omega-3 fatty acid formulation approved," *Ob.Gyn.News*, (Jan. 15, 2005).

Aung T, Halsey J, Kromhout D, et al. Associations of omega-3 fatty acid supplement use with cardiovascular disease risks: Meta-analysis of 10 trials involving 77917 individuals. *JAMA Cardiol* 3:225-34 (publication date Mar. 1, 2018; epublication date Jan. 31, 2018).

Avandia [package insert]. Research Triangle Park, NC: GlaxoSmithKline; 2011.(45 pages).

Avery et al., "Upper Gastrointestinal System," *Integrating Therapeutic and Complementary Nutrition*, Edited by Mary Marian, CRC Press (2006)(14 pages).

Aviram M, Rosenblat M, Bisgaier CL, Newton RS. Atorvastatin and gemfibrozil metabolites, but not the parent drugs, are potent antioxidants against lipoprotein oxidation. *Atherosclerosis*. Jun. 1998; 138(2):271-280.

Ayton, et al., "A pilot open case series of Ethyl-EPA supplementation in the treatment of anorexia nervosa." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 71, pp. 205-209. (Oct. 2004).

Ayton, et al., "Rapid improvement of severe anorexia nervosa during treatment with ethyl-eicosapentaenoate and micronutrients," *European Psychiatry* 19, pp. 317-319. (Aug. 2004).

Baigent, C., et al., "Efficacy and safety of cholesterol-lowering treatment: prospective meta-analysis of data from 90,056 participants in 14 randomised trials of statins." *Lancet*; 366:1267-1278. (Oct. 2005).

Baldwin RM, Ohlsson S, Pedersen RS, et al. Increased omeprazole metabolism in carriers of the CYP2C19*17 allele; a pharmacokinetic study in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2008 65 (5): 767-74.

US 10,568,861 B1

Page 7

(56)

References Cited

OTHER PUBLICATIONS

- Baldwin SJ, Clarke SE, Chenery RJ. Characterization of the cytochrome P450 enzymes involved in the in vitro metabolism of rosiglitazone. *Br. J. Clin. Pharmacol. Sep.* 1999;48:424-432.
- Balfour et al., "Rosiglitazone," *Drugs*, 57(6):921-930 (Jun. 1999).
- Balk, E.M. et al., "Effects of omega-3 fatty acids on serum markers of cardiovascular disease risk: a systematic review. *Atherosclerosis*." 189:19-30. (Nov. 2006).
- Ballantyne CM, Bays HE, Kastelein JJ, et al. Efficacy and safety of eicosapentaenoic acid ethyl ester (AMR 101) therapy in statin-treated patients with persistent high triglycerides (from the ANCHOR study). *Am J Cardiol Oct.* 2012 110 (7): 984-92.
- Ballantyne et al., "Abstract 15071: AMR101 Lowers Triglycerides, Atherogenic Lipoprotein, Phospholipase A₂, and High-sensitivity C-reactive Protein Levels in Patients with High Triglycerides and on Background Statin Therapy (the ANCHOR Study)," *Circulation, Lippincott Williams and Wilkins*, vol. 124, No. 21, Suppl., Nov. 22, 2011.
- Ballantyne et al., "Effects of icosapent ethyl on lipoprotein particle concentration and the fatty acid desaturation index in statin-treated patients with persistent high triglycerides (the ANCHOR study)." *Journ. Clin. Lipidology*, 2013, 7(3):270-271.
- Ballantyne et al., Influence of low-high density lipoprotein cholesterol and elevated triglyceride on coronary heart disease events and response to simvastatin therapy in 4S, *Circulation*, 104:3046-3051. (Dec. 2001).
- Bang HO, Dyerberg J. "Plasma lipids and Lipoproteins in Greenlandic west coast Eskimos" *Acta Med Scand*, 192:85-94. (Jul./Aug. 1972).
- Banga, A., et al., "Adiponectin translation is increased by the PPAR γ agonists pioglitazone and ω -3 fatty acids." *Am J Physiol Endocrinol Metab* 296:480-489 (Mar. 2009).
- Bangham et al., "Diffusion of univalent ions across the lamellae of swollen phospholipids." *J. Mol. Biol.* (Aug. 1965) 13(1):238-252.
- Bansal S, Buring JE, Rifai N, Mora S, Sacks FM, Ridker PM, "Fasting Compared With Nonfasting Triglycerides and Risk of Cardiovascular Events in Women," *JAMA*, 298:309-316 (Jul. 2007).
- Barter et al., "Effectiveness of Combined Statin Plus Omega-3 Fatty Acid Therapy for Mixed Dyslipidemia." *Am. J. Cardiol.* 102(8):1040-1045 (Oct. 15, 2008).
- Basu, A., et al., "Dietary Factors That Promote or Retard Inflammation." *Arterioscler. Thromb. Vasc. Biol.* 26:995-1001 (May 2006).
- Baynes JW. Role of oxidative stress in development of complications in diabetes. *Diabetes*. Apr. 1991;40(4):405-412.
- Bays HE et al. "Prescription omega 3 fatty acids and their lipid effects: physiologic mechanisms of action and clinical implications." *Expert Rev Cardiovasc Ther.*, 6:391-409. (Mar. 2008).
- Bays HE, Ballantyne CM, Braeckman RA, Stirlen WG, Soni PN. Icosapent ethyl, a pure ethyl ester of eicosapentaenoic acid: effects on circulating markers of inflammation from the MARINE and ANCHOR studies. *Am. J. Cardiovasc. Drugs*. Feb. 2013;13(1):37-46.
- Bays HE, Braeckman RA, Ballantyne CM, et al. Icosapent ethyl, a pure EPA omega-3 fatty acid: Effects on lipoprotein particle concentration and size in patients with very high triglyceride levels (the MARINE study). *J. Clin. Lipidol.* Nov./Dec. 2012;6:565-572.
- Bays HE, Safety considerations with omega-3 fatty acid therapy. *Am. J. Cardiol.* Mar. 2007 99 (6A): 35C-43C.
- Bays, H., Clinical Overview of Omacor: A Concentrated Formulation of Omega-3 Polyunsaturated Fatty Acids, *Am J Cardiol.*; 98[suppl]:711-761 (Aug. 2006).
- Bays, H., "Rationale for Prescription Omega-3-Acid Ethyl Ester Therapy for Hypertriglyceridemia: A Primer for Clinicians," *Drugs of Today*, 44(3); 205-246. (Mar. 2008).
- Bays, H.E., Eicosapentaenoic Acid Ethyl Ester (AMR101) Therapy in Patients With Very High Triglyceride Levels (from the Multi-center, pLAceto-controlled, Randomized, double-blind, 12-week study with an open-label Extension [MARINE] Trial) *Am J Cardiol*;108:682-690. (Sep. 2011).
- Bays, H.E., et al., "Long-term up to 24-month efficacy and safety of concomitant prescription omega-3-acid ethyl esters and simvastatin in hypertriglyceridemic patients." *Curr Med Res Opin.*; 26:907-915. (Apr. 2010).
- Beal, M.F., *Annals of Neurology*, vol. 38, No. 3, "Aging, Energy, and Oxidative Stress in Neurodegenerative Diseases", pp. 357-366, (Sep. 1995).
- Beaumont et al., Design of Ester Prodrugs to Enhance Oral Absorption of Poorly Permeable Compounds: Challenges to the Discovery Scientist, *Current Drug and Metabolism*. (Dec. 2003) 4:461-485.
- Becker LB, Aufderheide TP, Geocadin RG, Callaway CW, Lazar RM, Donnino MW, Nadkarni VM, Abella BS, Adrie C, Berg RA, Merchant RM, O'Connor RE, Meltzer DO, Holm MB, Longstreth WT, Halperin HR. AHA Consensus Statement: Primary Outcomes for Resuscitation Science Studies: A Consensus Statement From the American Heart Association. *Circulation* 2011; CIR. 0b013e3182340239 published online before print Oct. 3, 2011, doi:10.1161/CIR.0b013e3182340239.
- Belarbi et al., "A process for high yield and scaleable recovery of high purity eicosapentaenoic acid esters from microalgae and fish oil," *Enzyme and Microbial Technology* 26:516-529 (Apr. 2000).
- Belger et al., "Assessment of prefrontal activation by infrequent visual targets and non-target novel stimuli in schizophrenia: a function MRI study," Presented at the 9th Biennial winter workshop on schizophrenia, Davos, Switzerland, Feb. 7-13, 1998, Abstract in *Schizophrenia Research*. vol. 29. No. 1/02, Jan. 1998.
- Belikov, *Pharmaceutical Chemistry in Two Parts*, 1/General Pharmaceutical Chemistry 43-47 (1993) (with English translation)(9 pages).
- Belmaker et al., "Addition of Omega-3 Fatty Acid to Maintenance Medication Treatment for Recurrent Unipolar Depressive Disorder," *Am. J. Psychiatry*, 159:477-479 (Mar. 2002).
- Belmaker, et al., "Omega-3 Eicosapentaenoic Acid in Bipolar Depression: Report of a Small Open-Label Study," *J Clin Psychiatry*; 66:726-729. (Jun. 2005).
- Bender NK, Kraynak MA, Chiquette E, et al. Effects of marine fish oils on the anticoagulation status of patients receiving chronic warfarin therapy. *J. Thromb. Thrombolysis* Jul. 5, 1998 (3): 257-61.
- Bénistant, C., et al., "Docosapentaenoic acid (22:5, n-3): metabolism and effect on prostacyclin production in endothelial cells." *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 55(4):287-292, (Oct. 1996).
- Benn et al., Improving Prediction of Ischemic Cardiovascular Disease in the General Population Using Apolipoprotein B: The Copenhagen City Heart Study, 27 *Arteriosclerosis, Thrombosis, & Avascular Biology* 661 (Mar. 2007).
- Bennett et al., "Treatment of IgA nephropathy with eicosapentaenoic acid (EPA): a two-year prospective trial [Abstract Only]." *Clin. Nephrol.* 31(3):128-131 (Mar. 1989).
- Berge, R.K., et al., "In contrast with docosahexaenoic acid, eicosapentaenoic acid and hypolipidaemic derivatives decrease hepatic synthesis and secretion of triacylglycerol by decreased diacylglycerol acyltransferase activity and stimulation of fatty acid oxidation." *Biochem J.*; 343(Pt 1):191-197. (Oct. 1999).
- Berglund L, Brunzell JD, Goldberg AC, et al. Evaluation and treatment of hypertriglyceridemia: an endocrine society clinical practice guideline. *J. Clin. Endocrinol. Metab.* Sep. 2012 97 (9): 2969-89.
- Berliner JA, Watson AD. A role for oxidized phospholipids in atherosclerosis. *N. Engl. J. Med.* Jul. 2005;353(1):9-11.
- Bertelsen M, Anggard EE, Carrier MJ. Oxidative stress impairs insulin internalization in endothelial cells in vitro. *Diabetologia*. May 2001;44(5):605-613.
- Betteridge, D.J., "Diabetic dyslipidaemia: past, present and future." *Practical Diabetes Int.* 21(2): 78-85. (Mar. 2004).
- Bhatt DL, Eagle KA, Ohman EM, et al. Comparative determinants of 4-year cardiovascular event rates in stable outpatients at risk of or with atherothrombosis. *JAMA* 304(12):1350-7 (publication date Sep. 22, 2010; epublication date Aug. 30, 2010).
- Bhatt DL, Fox KAA, Hacke W, et al; Charisma Investigators. Clopidogrel and aspirin versus aspirin alone for the prevention of atherothrombotic events. *N Engl J Med.* 354(16):1706-1717 (publication date Apr. 20, 2006; epublication date Mar. 12, 2006).

US 10,568,861 B1

Page 8

(56)

References Cited

OTHER PUBLICATIONS

- Bhatt DL, Hulot JS, Moliterno DJ, Harrington RA. Antiplatelet and anticoagulation therapy for acute coronary syndromes. *Circ Res* 114(12):1929-43 (publication date Jun. 6, 2014).
- Bhatt DL, Steg PG, Brinton EA, et al. Rationale and design of Reduce-It: Reduction of Cardiovascular Events with Icosapent Ethyl-Intervention Trial. *Clin Cardiol* 40:138-48 (publication date Mar. 2017; epublication date Mar. 15, 2017).
- Bhatt DL, Steg PG, Ohman EM, et al; Reach Registry Investigators. International prevalence, recognition and treatment of cardiovascular risk factors in outpatients with atherothrombosis. *JAMA*. 295(2):180-189 (publication date Jan. 11, 2006).
- Bhatt et al., "Cardiovascular Risk Reduction with Icosapent Ethyl for Hypertiglyceridemia." *N. Eng. J. Med.*, Nov. 10, 2018 (epub ahead of print)(12 pages)(downloaded from nejm.org on Nov. 13, 2018 at <https://www.nejm.org/doi/full/10.1056/NEJMoa1812792>).
- Bild et al., "Multi-Ethnic Study of Atherosclerosis: objectives and design," *Am J Epidemiol* 156(9):871-81 (Nov. 1, 2002).
- Black et al., "Effect of intravenous eicosapentaenoic acid on cerebral blood flow, edema, and brain prostaglandins in ischemic gerbils", *Prostaglandins*, 28(4), pp. 545-546. (Oct. 1984).
- Blankenhorn D.H. et al., "Beneficial effects of combined colestipol-niacin therapy on coronary atherosclerosis and coronary venous bypass grafts." *JAMA* 257: 3233-3240. (Jun. 1987).
- Block, R. C., et al., "EPA and DHA in blood cell membranes from acute coronary syndrome patients and controls." *Atherosclerosis*, 197(2):821-828 (Apr. 2008).
- Blumenthal, *Advanced Studies in Medicine*, 2:148-157 (2002).
- Boden WE, Probstfield JL, Anderson T, Chaitman BR, Desvignes-Nickens P, Koprowicz K, IJ McBride R, Teo K, Weintraub W. and the Aim-High Investigators, "Niacin in patients with low hdl cholesterol levels receiving intensive statin therapy," *N. Engl. J. Med.* Dec. 2011;365:2255-2267.
- Bonaa, KH et al., Docosahexaenoic and Eicosapentaenoic acids in plasma phospholipids are divergently associated with high density lipoprotein in humans, *Arterioscler. Thromb. Vasc. Biol.*;12;675-681 (Jun. 1992).
- Bonnet et al., "Comparative Effects of 10-mg Versus 80-mg Atorvastatin on High-Sensitivity C-Reactive Protein in Patients with Stable Coronary Artery Disease: Results of the CAP(Comparative Atorvastatin Pleiotropic Effects) Study," *Clinical Therapeutics*. 30(12):2298-2313 (Dec. 2008).
- Borchman D, Lamba OP, Salmassi S, Lou M, Yapped MC. The dual effect of oxidation on lipid bilayer structure. *Lipids*. Apr. 1992;27(4):261-265.
- Bordin et al., "Effects of fish oil supplementation on apolipoprotein B100 production and lipoprotein metabolism in normolipidaemic males," *Eur. J. Clin. Nutr.* 52: 104-9 (Feb. 1998).
- Borow et al., "Biologic plausibility, cellular effects, and molecular mechanisms of eicosapentaenoic acid (EPA) in atherosclerosis," *Atherosclerosis*, 242(1):357-66 (Sep. 2015).
- Borthwick et al., "The effects of an omega-3 ethyl ester concentrate on blood lipid concentrations in pateitns with hyperlipidemia," *Clin. Drug Investig.* (1998) 15(5): 397-404.
- Bossaller C, Habib GB, Yamamoto H, Williams C, Wells S, Henry PD. Impaired muscarinic endothelium-dependent relaxation and cyclic guanosine 5'-monophosphate formation in atherosclerotic human coronary artery and rabbit aorta. *J. Clin. Invest.* Jan. 1987;79:170-174.
- Bousserouel, S., et al., "Different effects of n-6 and n-3 polyunsaturated fatty acids on the activation of rat smooth muscle cells by interleukin-1beta." *J. Lipid Res.* 44:601-611 (Mar. 2003).
- Bousserouel, S., et al., "Modulation of cyclin D1 and early growth response factor-1 gene expression in interleukin-1beta-treated rat smooth muscle cells by n-6 and n-3 golyunsaturated fatty acids." *Eur. J. Biochem.* 271:4462-4473 (Nov. 2004).
- Brady, L., et al., Increased n-6 polyunsaturated fatty acids do not attenuate the effects of long-chain n-3 polyunsaturated fatty acids on insulin sensitivity or triacylglycerol reduction in Indian Asians. *Am J Clin Nutr* 79:983-91(Jun. 2004).
- Braeckman et al., "Abstract 18549: Effects of AMR101, a Pure Eicosapentaenoic Omega-3 Fatty Acid, on the Fatty Acid Profile in Plasma and Red Blood Cells in Statin-Treated Patients with Persistent High Triglycerides—Results from the ANCHOR study," *Circulation* 126(21S):A15071 (Nov. 20, 2012)(2 pages).
- Braeckman et al., "Effect of Concomitant Icosapent Ethyl (Eicosapentaenoic Acid Ethyl Ester) on Pharmacokinetics of Atorvastatin," *Clinical Drug Investigation*. (Jan. 2015) (3)45-51.
- Braeckman RA, Manku MS, Bays HE, Stirtan WG, Soni PN. Icosapent ethyl, a pure EPA omega-3 fatty acid: effects on plasma and red blood cell fatty acids in patients with very high triglyceride levels (results from the MARINE study). *Prostaglandins Leukot Essent Fatty Acids*. Sep. 2013;89(4):195-201.
- Braeckman RA, Stirtan WG, Soni PN. Pharmacokinetics of eicosapentaenoic acid in plasma and red blood cells after multiple oral dosing with AMR101 (ethyleicosapentaenoic acid) in healthy subjects [abstract]. Presented at: Congress of the International Society for the Study of Fatty Acids and Lipids, Vancouver, Canada, May 26-30, 2012.
- Braeckman RA, Stirtan WG, Soni PN. Pharmacokinetics of eicosapentaenoic acid in plasma and red blood cells after multiple oral dosing with icosapent ethyl in healthy subjects. *Clin. Pharmacol. Drug Dev.* Mar. 2014 (epub Oct. 22, 2013); 3:101-108.
- Braunersreuther V, Steffens S, Arnaud C, Pelli G, Burger F, Proudfoot A, Mach F. A novel rantes antagonist prevents progression of established atherosclerotic lesions in mice. *Arterioscler. Thromb. Vasc. Biol.* Jun. 2008;28:1090-1096.
- Breslow, J., "n-3 Fatty acids and cardiovascular disease." *Am J Clin Nutr.* 83:1477S-82S (Jun. 2006).
- Brinton EA, Ballantyne CM, Bays HE, Kastelein JJ, Braeckman RA, Soni PN. Effects of AMR101 on lipid and inflammatory parameters in patients with diabetes mellitus-2 and residual elevated triglycerides (200-500 mg/dl) on statin therapy at LDL-C goal: the ANCHOR study.[abstract 629-P] *Diabetes*. 2012;61(suppl 1):A159-A160.
- Brinton et al., "Effects of icosapent ethyl on lipid and inflammatory parameters in patients with diabetes mellitus-2, residual elevated triglycerides (200-500 mg/dL), and on statin therapy at LDL-C goal: the ANCHOR study," *Cardiovasc. Diabetol.* Jul. 9, 2013;12:100. doi: 10.1186/1475-2840-12-100.
- Brossard, N., et al., "Retroconversion and metabolism of [13C]22:6n-3 in humans and rats after intake of a single dose of [13C]22:6n-3—3-triacylglycerols." *Am. J. Clin. Nutr.* 64:577-86 (Oct. 1996).
- Brouwer, I.A., et al., "Effect of fish oil on ventricular tachyarrhythmia and death in patients with implantable cardioverter defibrillators." *JAMA*. 295(22):2613-2619 (Jun. 2006).
- Brovkovich V, Dobrucki LW, Brovkovich S, Dobrucki I, Do Nascimento CA, Burewicz A, Malinski T. Nitric oxide release from normal and dysfunctional endothelium. *J. Physiol. Pharmacol.* Dec. 1999;50:575-586.
- Brown et al., Simvastatin and Niacin, Antioxidant Vitamins, or the Combination for the Prevention of Coronary Disease, *N Engl J Med*, vol. 345, No. 22, 1583-1592 (Nov. 29, 2001).
- Brown, A. J., et al., "Administration of n-3 Fatty Acids in the Diets of Rats or Directly to Hepatocyte Cultures Results in Different Effects on Hepatocellular ApoB Metabolism and Secretion." *Arterioscler. Thromb. Vasc. Biol.* 19:106-114 (Jan. 1999).
- Brown, A. J., et al., "Persistent changes in the fatty acid composition of erythrocyte membranes after moderate intake of n-3 polyunsaturated fatty acids: study design and implications." *Am.J. Clin. Nutri.* 54:668-73(Oct. 1991).
- Brown, G., et al., "Regression of coronary artery-disease as a result of intensive lipid-lowering therapy in men with high levels of apolipoprotein," *B., N. Engl. J. Med.* 323: 1289-1298. (Nov. 1990).
- Brownlee M. Biochemistry and molecular cell biology of diabetic complications. *Nature*. Dec. 2001; 414(6865):813-820.
- Bryhn, M., et al., "The bioavailability and pharmacodynamics of different concentrations of omega-3 acid ethyl esters." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 75:19-24 (Jul. 2006).
- Budavari, S., Editor, "The Merck Index", Merck & Co., Inc., p. 725 item 4511 and p. 279 and item 2417 (1989).

US 10,568,861 B1

Page 9

(56)

References Cited

OTHER PUBLICATIONS

Budoff, "Triglycerides and Triglyceride-Rich Lipoproteins in the Causal Pathway of Cardiovascular Disease," *Am. J. Cardiol.*, 118(1):138-45 (Jul. 1, 2016).

Bunting et al. "Depression in Parkinson's Disease". *J Neurosci Nurs.*; 23(3):158-164. (Abstract Only) (Jun. 1991).

Burdge, G.C., et al., "Eicosapentaenoic and docosapentaenoic acids are the principal products of α -linolenic acid metabolism in young men." *British Journal of Nutrition* 88:355-363 (Oct. 2002).

Burdge, G.C., et al., "Lack of effect of meal fatty acid composition on postprandial lipid, glucose and insulin responses in men and women aged 50-65 years consuming their habitual diets." *British Journal of Nutrition*, 96:489-500 (Sep. 2006).

Burdge, G.C., et al., "The effect of altering the 20:5n-3 and 22:6n-3 content of a meal on the postprandial incorporation of n-3 polyunsaturated fatty acids into plasma triacylglycerol and non-esterified fatty acids in humans." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 77:59-65 (Jul. 2007).

Burr ML, Sweetnam PM, Fehily AM. Diet and reinfarction. *Eur Heart J* 15:1152-1153, Aug. 1994.

Burr, M. L., et al., "Effects of changes in fat, fish and fibre intakes on death and myocardial reinfarction: Diet and reinfarction trial." *The Lancet*, 2(8666):757-61 (Sep. 1989).

Buse JB, Ginsberg HN, Bakris GL, et al. Primary prevention of cardiovascular diseases in people with diabetes mellitus: a scientific statement from the American Heart Association and the American Diabetes Association. *Diabetes Care*. 2007;30: 162-172.

Calabresi, L., et al., "Omacor in familial combined hyperlipidemia: effects on lipids and low density lipoprotein subclasses." *Atherosclerosis* 148:387-396 (Feb. 2000).

Calder PC. Omega-3 Fatty Acids and Inflammatory Processes. *Nutrients* 2(3):355-374, Mar. 2010 (epub Mar. 18, 2010).

Calder PC. The role of marine omega-3 (n-3) fatty acids in inflammatory processes, atherosclerosis and plaque stability. *Mol. Nutr. Food Res.* Jul. 2012;56(7):1073-1080.

Campos, H., et al., "Low-density lipoprotein size, pravastatin treatment, and coronary events." *JAMA*, 286:1468-1474 (Sep. 2001).

Canner P.L. et al., "Fifteen year mortality in Coronary Drug Project patients: long-term benefit with niacin," *J. Am. Coll. Cardiol.* 8. 1245-1255. (Dec. 1986).

Cannon CP, Blazing MA, Giugliano RP, et al; Improve-It Investigators. "Ezetimibe added to statin therapy after acute coronary syndromes." *N Engl J Med.* 372:2387-2397. (Jun. 18, 2015/epub Jun. 3, 2015).

Cannon CP, Braunwald E, McCabe CH, et al. Intensive versus moderate lipid lowering with statins after acute coronary syndromes. *N Engl J Med* 350(15):1495-1504 (publication date Apr. 8, 2004; epublication date Mar. 8, 2004).

Cao H, Wang X, Huang H, Ying SZ, Guy W, Wang T, Huang CX. Omega-3 Fatty Acids in the Prevention of Atrial Fibrillation Recurrences after Cardioversion: A Meta-analysis of Randomized Controlled Trials. *Intern. Med.* 2012 (epub Sep. 15, 2012); 51:2503-2508.

Cao, et al., "Cloning, Expression, and Chromosomal Localization . . .", *Genomics*, 49:327-331, (Apr. 15, 1998).

Cao, J., et al., "Incorporation and Clearance of Omega-3 Fatty Acids in Erythrocyte Membranes and Plasma Phospholipids." *Clinical Chemistry* 52(12):2265-2272 (Dec. 2006).

Capuzzi, DM et al., "Efficacy and safety of an extended-release niacin (Niaspan): a long-term study." *Am. J. Cardiol.* 82: 74U-81U. (Dec. 17, 1998).

Carlson, L.A. & Rosenhamer G., "Reduction of mortality in the Stockholm Ischaemic Heart Disease Secondary Prevention Study by combined treatment with clofibrate and nicotinic acid." *Acta Med. Scand.* 223, 405-418 (1988).

Carlson, L.A., "Nicotinic acid: the broad spectrum lipid drug. A 50th Anniversary review", *J. Int. Med.*, 258:94-114, (Aug. 2005).

Carrero et al., "Intake of Fish Oil, Oleic Acid, Folic Acid, and Vitamins B-6 and E for 1 Year Decreases Plasma C-Reactive Protein

and Reduces Coronary Heart Disease Risk Factors in Male Patients in a Cardiac Rehabilitation Program", pp. 384-390 (Feb. 2007).

Carrero, J.J. et al. "Efectos cardiovasculares de los acidos grasos omega-3 y alternativas para incrementar su ingesta," *Nutricion Hospitalaria*. (2005) (1) 63-69 [with English abstract].

Carroll, D. N., et al., "Evidence for the Cardioprotective Effects of Omega-3 Fatty Acids." *Ann Pharmacother.*, 36:1950-6 (Dec. 2002).

Carulli et al., "Chenodeoxycholic acid and ursodeoxycholic acid effects in endogenous hypertriglyceridemias. A controlled double-blind trial." *J. Clin. Pharmacol.*, 21(10):436-42 (Oct. 1981).

Caughey GE, Mantzioris E, Gibson RA, Cleland LG, James MJ. The effect on human tumor necrosis factor α and interleukin 1 β production of diets enriched in n-3 fatty acids from vegetable oil or fish oil. *Am J Clin Nutr.* Jan. 1996;63:116-122.

Cavender MA, Steg PG, Smith SC, et al; REACH Registry Investigators. Impact of diabetes mellitus on hospitalization for heart failure, cardiovascular events, and death: outcomes at 4 years from the reduction of atherothrombosis for continued health (REACH) registry. *Circulation*. 132(10):923-931 (publication date Sep. 8, 2015; epublication date Jul. 7, 2015).

Cawood AL, Ding R, Napper FL, et al. Eicosapentaenoic acid (EPA) from highly concentrated n-3 fatty acid ethyl esters is incorporated into advanced atherosclerotic plaques and higher plaque EPA is associated with decreased plaque inflammation and increased stability. *Atherosclerosis*. Sep. 2010 (epub May 20, 2010); 212:252-259.

Cazzola, R., et al., "Age- and dose-dependent effects of an eicosapentaenoic acid-rich oil on cardiovascular risk factors in healthy male subjects." *Atherosclerosis* 193:159-167 (Jul. 2007).

Ceci et al., "The effects of oral 5-hydroxytryptophan administration on feeding behavior in obese adult female subjects," *J Neural. Transm* (1989) 76(2):109-117.

Cefali, E.A., et al., "Aspirin reduces cutaneous flushing after administration of an optimised extended-release niacin formulation", *Int. J. Clin. Pharmacol. & Ther.*, 45(2):78-88, (Feb. 2007).

Center for Drug Evaluation and Research. Application No. 21-853, 21654s016, (Omacor). Statistical Review and Evaluation: Clinical Studies, Omacor (omega-3 acid ethyl ester) Capsules, 4 grams/day; 2007. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/nda/2007/021853s000;%20021654s016_StatR.pdf. (Accessed Jan. 26, 2012) (156 pages).

Center for Drug Evaluation and Research. Approval Package for Application No. 202057Orig1s000. Review—Vascepa (formerly AMR101), 373 pages (Jul. 26, 2012)(in two parts).

Center for Drug Evaluation and Research. Approval Package for: 21-654 (Omacor/Lovaza). Statistical Review; 2004. Available at: http://www.accessdata.fda.gov/drugsatfda_docs/nda/2004/21-654_Omacor_AdminCorres_P1.pdf. Accessed Jan. 26, 2012. (54 pages).

Ceriello A, Motz E. Is oxidative stress the pathogenic mechanism underlying insulin resistance, diabetes, and cardiovascular disease? The common soil hypothesis revisited. *Arterioscler. Thromb. Vasc. Biol.* (May 2004);24(5):816-823.

Chait A, Brazg RL, Tribble DL, Krauss RM. Susceptibility of small, dense, low-density lipoproteins to oxidative modification in subjects with the atherogenic lipoprotein phenotype, pattern B. *Am. J. Med.* (Apr. 1993);94(4):350-356.

Chan et al., "Effect of Atorvastatin and Fish Oil on Plasma High-Sensitivity C-Reactive Protein Concentrations in Individuals with Visceral Obesity", *Clin. Chem.*, vol. 48, pp. 877-883 (2002).

Chan et al., Factorial Study of the Effects of Atorvastatin and Fish Oil on Dyslipidaemia in Visceral Obesity, 32 Euro. J. Clinical Investigation. 32(6):429-36 (Jun. 2002)

Chan, D.C., et al., "Randomized controlled trial of the effect of n-3 fatty acid supplementation on the metabolism of apolipoprotein B-100 and chylomicron remnants in men with visceral obesity." *Am J Clin Nutr* 77:300-7 (2003).

Chang CL, Seo T, Du CB, Accili D, Deckelbaum RJ. n-3 Fatty Acids Decrease Arterial Low-Density Lipoprotein Cholesterol Delivery and Lipoprotein Lipase Levels in Insulin-Resistant Mice. *Arterioscler Thromb Vasc Biol.* Dec. 2010 (epub Oct. 7, 2010); 30(12):2510-2517.

US 10,568,861 B1

Page 10

(56)

References Cited

OTHER PUBLICATIONS

- Chapman, M.J., et al., "Cholesteryl ester transfer protein: at the heart of the action of lipid-modulating therapy with statins, fibrates, niacin, and cholesteryl ester transfer protein inhibitors." *Eur Heart J.*, 31(2):149-164 (Jan. 2010).
- Chatterjee SN, Agarwal S. Liposomes as membrane model for study of lipid peroxidation. *Free Radic. Biol. Med.* 1988;4(1):51-72.
- Chemical Book, Eicosapentaenoic acid ethyl ester, copyright 2010, printed Jun. 16, 2011 from www.chemicalbook.com. (2010).
- Chen, H., et al., "Eicosapentaenoic acid inhibits hypoxia-reoxygenation-induced injury by attenuating upregulation of MMP-1 in adult rat myocytes." *Cardiovascular Research* 59:7-13 (Jul. 2003).
- Chen, H., et al., "EPA and DHA attenuate ox-LDL-induced expression of adhesion molecules in human coronary artery endothelial cells via protein kinase B pathway." *Journal of Molecular and Cellular Cardiology* 35:769-775 (Jul. 2003).
- Chen, I.S., et al., "In vitro clearance of chylomicron triglycerides containing (E)-3 eicosapentaenoate." *Atherosclerosis*, 65:193-198 (1987).
- Cheng et al., "Antagonism of the prostaglandin D2 receptor 1 suppresses nicotinic acid-induced vasodilation in mice and humans," *PNAS* 103(17):6682-7 (Apr. 25, 2006).
- Childs, M.T., et al., "Divergent lipoprotein Responses to Fish Oils With Various Ratios of Eicosapentaenoic Acid and Docosahexaenoic Acid", *American Society for Clinical Nutrition*, 52:632-9, (Oct. 1990).
- Christensen, J. H., et al., "Effect of fish oil on heart rate variability in survivors of myocardial infarction: a double blind randomised controlled trial." *BMJ*, 312:677-678 (Mar. 16, 1996).
- Christensen, M.S., et al., "Intestinal absorption and lymphatic transport of eicosapentaenoic (EPA), docosahexaenoic (DHA), and decanoic acids: dependence on intramolecular triacylglycerol structure." *Am J Clin Nutr* 61:56-61 (Jan. 1995).
- Citizen Petition, Pronova BioPharma Norge AS, (Aug. 4, 2009), at ii (Appendix), available at www.regulations.gov.
- Classification of Hyperlipidaemias and Hyperlipoproteinaemias, *Bulletin of the World Health Organization*, 43(6): 891-915 (1970).
- Cleland, L.G., et al., "A Biomarker of n-3 compliance in patients taking fish oil for rheumatoid arthritis." *Lipids* 38:419-424 (Apr. 2003).
- Clinical Trial NCT01047501, Effect of AMR101 (Ethyl Eicosapentate) on Triglyceride (Tg) Levels in Patients on Statins With High Tg Levels (>200 and <500 mg/dL) (ANCHOR), ClinicalTrials.gov [database online], U.S. National Institute of Health, Jan. 2010 [retrieved Apr. 27, 2011], Retrieved from the Internet: <<http://clinicaltrials.gov/ct2/show/NCT01047501>> (3 pages).
- Cohen AW, Combs TP, Scherer PE, Lisanti MP. Role of caveolin and caveolae in insulin signaling and diabetes. *American journal of physiology. Endocrinology and metabolism*. (Dec. 2003);285(6):E1151-1160.
- Cohen, J.D., et al., "30-year trends in serum lipids among United States adults: results from the National Health and Nutrition Examination Surveys II, III, and 1999-2006." *Am J Cardiol.*, 106:969-975. (Dec. 15, 2010).
- Cole et al., "Challenges and opportunities in the encapsulation of liquid and semi-solid formulations into capsules for oral administration," *Advanced Drug Delivery Reviews*, vol. 60, No. 6, pp. 747-756. (Mar. 17, 2007).
- Colhoun, H. M., et al., "Primary prevention of cardiovascular disease with atorvastatin in type 2 diabetes in the Collaborative Atorvastatin Diabetes Study (CARDS): multicentre randomised placebo-controlled trial." *Lancet* 364: 685-9 (Aug. 21-24, 2004).
- Collins, N., et al., "Differences between Dietary Supplement and Prescription Drug Omega-3 Fatty Acid Formulations: A Legislative and Regulatory Perspective." *Journal of the American College of Nutrition*, 27 (6):659-666 (Dec. 2008).
- Committee Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages. (2013).
- Conklin, S. M., et al., "Serum (E)-3 fatty acids are associated with variation in mood, personality and behavior in hypercholesterolemic community volunteers." *Psychiatry Research* 152: 1-10 (Jul. 30, 2007).
- Connor et al., "Seminars in thrombosis and hemostasis," 14:271-284. (1988).
- Connor, W.E., "Importance of n-3 Fatty Acids in Health and Disease", *Am. J. Clin. Nutr.*, 71(1(S)):171S-175S, (Jan. 2000).
- Conquer, J.A., et al., "Effect of supplementation with different doses of DHA on the levels of circulating DHA as non-esterified fatty acid in subjects of Asian Indian background. *J Lipid Res.*" 39:286-292. (Feb. 1998).
- Conquer, J.A., et al., "Supplementation with an algae source of docosahexaenoic acid increases (n-3) fatty acid status and alters selected risk factors for heart disease in vegetarian subjects." *J Nutr.*, 126: 3032-3039. (Dec. 1996).
- Contacos et al. Effect of pravastatin and omega-3 fatty acids on plasma lipids and lipoproteins in patients with combined hyperlipidemia, pp. 1755-1762 (Dec. 1993).
- Coronary Artery Bypass Grafting, NIH, published online Feb. 23, 2012 (12 pages).
- Costanzo S, di Niro V, Castelnovo AD, et al. Prevention of postoperative atrial fibrillation in open heart surgery patients by preoperative supplementation of n-3 polyunsaturated fatty acids: An updated meta-analysis. *Periop Manga*. Apr. 12, 2013; epub.
- Coumadin [package insert], Princeton, NJ: Bristol-Myers Squibb; 2011. (10 pages).
- Cox PJ, Ryan DA, Hollis FJ, et al. Absorption, disposition, and metabolism of rosiglitazone, a potent thiazolidinedione insulin sensitizer, in humans. *Drug Metab. Dispos.* Jul. 2000;28:772-780.
- Creager MA, Gallagher SJ, Gireddi XJ, Coleman SM, Dzau VJ, Cooke JP. L-arginine improves endothelium-dependent vasodilation in hypercholesterolemic humans. *J. Clin. Invest.* Oct. 1992;90:1248-1253.
- Crevel et al., "Allergenicity of Refined Vegetable Oils," *Food and Chemical Toxicology*, 38, pp. 385-393 (Apr. 2000).
- Criqui, M., "Triglycerides and Coronary Heart Disease Revisited (Again)," vol. 147 No. 6, pp. 425-427 (Sep. 18, 2007).
- Cromwell et al., "LDL particle number and risk of future cardiovascular disease in the Framingham Offspring Study—Implications for LDL Management," *Journal of Lipidology*. (Dec. 2007) 1, 583-592.
- Crowe, F. L., et al., "Serum phospholipid n-3 long-chain polyunsaturated fatty acids and physical and mental health in a population-based survey of New Zealand adolescents and adults." *Am J Clin Nutr* 86:1278-85 (Nov. 2007).
- Cruz et al., "The metabolic syndrome in children and adolescents," *Curr. Diab. Rep.*, vol. 4(1):53-62 (Feb. 2004).
- Culhane et al., "Rosuvastatin for the treatment of hypercholesterolemia," *Pharmacotherapy*, 25(7):990-1000 (Jul. 2005).
- Daggy, B., et al., Dietary fish oil decreases VLDL production rates. *Biochimica et Biophysica Acta* 920: 293-300 (Aug. 15, 1987).
- Dall et al., "Clinical utility of low-density lipoprotein particle measurement in management of cardiovascular disease: a case report," *Research Reports in Clin. Cardiol.*, vol. 2, pp. 57-62 (2011).
- Daniel et al., "The Effect of Elevated Triglycerides on the Onset and Progression of Coronary Artery Disease: A Retrospective Chart Review," *Cholesterol*, vol. 2015 (epub Nov. 4, 2015), Article ID 292935, 5 pages.
- Das, U.N., Essential fatty acids as possible mediators of the actions of statins. *Prostaglandins, Leukotrienes and Essential Fatty Acids* 65(1):37-40, (Jul. 2001).
- Davidson MH, Ballantyne CM, Jacobson TA, et al. Clinical utility of inflammatory markers and advanced lipoprotein testing: advice from an expert panel of lipid specialists. *J. Clin. Lipidol.* Sep./Oct. 2011;5:338-367.
- Davidson MH, et al., Effects of prescription omega-3-acid ethyl esters on lipo protein particle concentrations, apolipoproteins AI and CIII, and lipoprotein-associated phospholipase A₂ mass in statin-treated subjects with hypertriglyceridemia, *J.Clin. Lipid.*, vol. 3(5), pp. 332-340 (Oct. 2009).

US 10,568,861 B1

Page 11

(56)

References Cited

OTHER PUBLICATIONS

- Davidson MH, Rosenson RS, Maki KC, Nicholls SJ, Ballantyne CM, Mazzone T, Carlson DM, Williams LA, Kelly MT, Camp HS, Lele A, Stolzembach JC. Effects of fenofibric acid on carotid intima-media thickness in patients with mixed dyslipidemia on atorvastatin therapy: Randomized, placebo-controlled study (first). *Arterioscler. Thromb. Vasc. Biol.* Jun. 2014;34:1298-1306.
- Davidson MH, Stein EA, Bays HE et al. "Efficacy and tolerability of adding prescription omega-3 fatty acids 4 g/d to simvastatin 40 mg/d in hypertriglyceridemic patients: an 8-week, randomized, double-blind, placebo-controlled study," *Clin Ther.*, 29:1354-1367. (Jul. 2007).
- Davidson MH., "Mechanisms for the hypotriglyceridemic effect of marine omega 3 fatty acids." *Am J Cardiol* 98(4A):27i-33i. (Aug. 21, 2006).
- Davidson, M.H., et al., "Effects of docosahexaenoic acid on serum lipoproteins in patients with combined hyperlipidemia: a randomized, doubleblind, placebo-controlled trial." *J Am Coll Nutr.*, 16:236-243. (Jun. 1997).
- Davies et al., "Rapid separation of LDL subclasses by iodixanol gradient ultracentrifugation," *Clin. Chem.*, 49(11):1865-72. (Nov. 2003).
- Davies-Tuck et al., "Total cholesterol and triglycerides are associated with development of new bone marrow lesions in asymptomatic middle-aged women—a prospective cohort study," *Arthritis Research & Therapy*. (published online Dec. 4, 2009) pp. 1-7
- De Caterina, R, et al., "Control of Endothelial Leukocyte Adhesion Molecules by Fatty Acids." *Lipids*, vol. 31:S57-S63 (1996).
- De Caterina, R., et al., "The Omega-3 fatty acid docosahexaenoate reduces cytokine-induced expression of proatherogenic and proinflammatory proteins in human endothelial cells." *Arterioscler. Thromb. Vasc. Biol.* 14:1829-1836 (1994).
- De Graaf J, Hak-Lemmers HL, Hectors MP, Demacker PN, Hendriks JC, Stalenhoef AF. Enhanced V susceptibility to in vitro oxidation of the dense low density lipoprotein subfraction in healthy subjects. *Arterioscler. Thromb.* 1991;11(2):298-306.
- De Moraes et al., "Evaluation of lipid extraction and fatty acid composition of human plasma," *Rev. Bras. Hematol. Hemoter.* 32(6):439-443 (2010).
- Deckelbaum, R. J., et al., "Conclusions and recommendations from the symposium, Beyond Cholesterol: Prevention and Treatment of Coronary Heart Disease with n-3 Fatty Acids." *Am J Clin Nutr* 87:2010S-12S (2008).
- Defendants' Invalidity Contentions, 3:14-CV-02550-MLC-DEA (D.N.J.), 520 pages (Dec. 5, 2014).
- Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 901 pages (Dec. 5, 2014).
- DeMets DL, Lan KK. Interim Analysis: the Alpha Spending Function Approach. *Stat Med.*, Jul. 15-30, 1994; 13(13-14):1341-52.
- Dewailly, E. et al., "n-3 Fatty acids and cardiovascular disease risk factors among the Inuit of Nunavik." *Am J Clin Nutr* 74:464-73 (2001).
- Dewey FE, Gusarova V, O'Dushlaine C, et al. Supplement to: Inactivating variants in ANGPTL4 and risk of coronary artery disease. *N Engl J Med*. DOI: 10.1056/NEJMoa1510926; Mar. 24, 2016 (epub Mar. 2, 2016).
- Di Spirito, M., Morelli, G., Doyle, R.T., Johnson, J. & McKenney, J. Effect of omega-3-acid ethyl esters on steady-state plasma pharmacokinetics of atorvastatin in healthy adults. *Expert Opin. Pharmacother.* 9, 2939-2945 (2008).
- Diagnostic and Statistical Manual of Mental Disorders, 4.Ed. Text revision, published by the American Psychiatric Assoc., pp. 154-163 and 369-381 (2000).
- Diagnostic and Statistical Manual of Mental Disorders, 4.sup.th Ed., published by the American Psychiatric Assoc., pp. 285-286, (1994).
- Dijan, P., et al., *Proc. Natl. Acad. Sci.*, vol. 93, "Codon repeats in genes associated with human diseases: Fewer repeats in the genes of nonhuman primates and nucleotide substitutions concentrated at the sites of reiteration," pp. 417-421, (Jan. 9, 1996)
- Dijk, J. M., et al., "Carotid intima-media thickness and the risk of new vascular events in patients with manifest atherosclerotic disease: the SMART study." *European Heart Journal* 27:1971-1978 (2006).
- Din et al., "Omega 3 fatty acids and cardiovascular disease—fishing for a natural treatment," *BMJ*, vol. 327, No. 7430, pp. 30-35 (Jan. 3, 2004).
- Djouisse L, Akinkuolie AO, Wu JHY, Ding EL, Gaziano JM. Fish consumption, omega-3 fatty acids and risk of heart failure: A meta-analysis. *Clin Nutr.* Dec. 2012 (epub Jun. 6, 2012); 31:846-853.
- Do R, Stitzel NO, Won HH, et. al. Exome sequencing identifies rare LDLR and APOA5 alleles conferring risk for myocardial infarction. *Nature*. Feb. 5, 2015 (epub Dec. 10, 2014); 518(7537):102-106.
- Do R, Willer CJ, Schmidt EM, et al. Common variants associated with plasma triglycerides and risk for coronary artery disease. *Nat Genet* Nov. 2013 (Oct. 6, 2013); 45(11):1345-52.
- Dodin, S., et al., "Flaxseed on cardiovascular disease markers in healthy menopausal women: a randomized, double-blind, placebo-controlled trial." *Nutrition* 24:23-30 (2008).
- Doi M, Nosaka K, Miyoshi T, et al. Early eicosapentaenoic acid treatment after percutaneous coronary intervention reduced acute inflammatory responses and ventricular arrhythmias in patients with acute myocardial infarction: A randomized controlled study. *Int J Cardiol.*, 176(3):577-82 (publication date Oct. 20, 2014; epublication date Aug. 19, 2014).
- Dolecek, "Epidemiological Evidence of Relationships Between Dietary Polyunsaturated Fatty Acids and Morality in the Multiple Risk Factor Intervention Trial", *Society of Experimental Biology and Medicine*, 200(2):177-182, (1991).
- Draft Agenda for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Draft Meeting Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Draft Questions for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 1 page.
- Drexler H, Zeiher AM, Meinzer K, Just H. Correction of endothelial dysfunction in coronary microcirculation of hypercholesterolaemic patients by L-arginine. *Lancet*. 1991;338:1546-1550.
- Dullenmeijer, C., et al., "n-3 Fatty acid proportions in plasma and cognitive performance in older adults." *Am J Clin Nutr* 86:1479-85 (2007).
- Duncan, R. E., et al., "Regulation of HMG-CoA reductase in MCF-7 cells by genistein, EPA, and DHA, alone and in combination with mevastatin." *Cancer Letters* 224:221-228 (2005).
- Durrington PN et al. "An omega 3 poly unsaturated fatty acid concentrate administered for one year decreased triglycerides in simvastatin treated patients with coronary heart disease and persistent Hypertriglyceridemia," *Heart*, 85:544-48 (2001).
- Dwyer, J. H., et al., "Arachidonate 5-Lipoxygenase Promoter Genotype, Dietary Arachidonic Acid, and Atherosclerosis." *N. Engl. J. Med.*, 350:1 (2004).
- Dyerberg, J., et al., "Marine Oils and Thrombogenesis." *Prog. Lipid Res.* 21:255-269 (1982).
- Egert, S., et al., "Dietary alpha-linolenic acid, EPA, and DHA have differential effects on LDL fatty acid composition but similar effects on serum lipid profiles in normolipidemic humans." *J Nutr.*, 139:861-868 (2009).
- Ehara S, Ueda M, Naruko T, Haze K, Itoh A, Otsuka M, Komatsu R, Matsuo T, Itabe H, Takano T, Tsukamoto Y, Yoshiyama M, Takeuchi K, Yoshikawa J, Becker AE. Elevated levels of oxidized low density lipoprotein show a positive relationship with the severity of acute coronary syndromes. *Circulation*. 2001;103(15):1955-1960.
- Eilat-Adar et al. "Association of Intentional Changes in Body Weight with Coronary Heart Disease Event Rates in Overweight Subjects who have an Additional Coronary Risk Factor," *Amer. Journ. Epidemiol.* 161(4)pp. 352-358 (Sep. 9, 2004).
- Eisenberg S, Bilheimer DW, Levy RI, Lindgren FT. "On the metabolic conversion of human plasma very low density lipoprotein to low density lipoprotein," *Biochim Biophys Acta*, 326:361-77 (1973).

US 10,568,861 B1

Page 12

(56)

References Cited

OTHER PUBLICATIONS

- Eisenberg S, Rachmilewitz D. "Metabolism of rat plasma very low density lipoprotein. I. Fate in circulation of the whole lipoprotein," *Biochim Biophys Acta*, 326:378-90 (1973).
- El-Serag HB, Graham DY, Satia JA, et al. Obesity is an independent risk factor for GERD symptoms and erosive esophagitis. *Am. J. Gastroenterol.* Jun. 2005 100 (6): 1243-50.
- Elam, M.B., et al., "Effect of niacin on lipid and lipoprotein levels and glycemic control in patients with diabetes and peripheral arterial disease study: a randomized trial", *The ADMIT [Arterial Disease Multiple Intervention Trial] JAMA*, 284:1263-1270, 2000.
- El-Saadani M, Esterbauer H, Ei-Sayed M, Gober M, Nassar AY, Jurgens G. A spectrophotometric assay for lipid peroxides in serum lipoproteins using commercially available reagent. *J. Lipid Res.* 1989;30:627-630.
- El-Soehy, A., et al., "Regulation of Mevalonate Synthesis in Low Density Lipoprotein Receptor Knockout Mice Fed n-3 or n-6 Polyunsaturated Fatty Acids." *Lipids*, 34 (10): 1037-43 (1999).
- Emsley et al., "Randomized, Placebo-Controlled Study of Ethyl-Eicosapentaenoic Acid as Supplemental Treatment in Schizophrenia," *Am. J. Psychiatry*, 159:1596-1598 (2002).
- Endo et al., "The Effects of Dietary Fatty Acids on Serum Lipids and Plasma Prostaglandin Levels in the Treatment of Obesity," *Japanese Journal of Pediatric Gastroenterology and Nutrition* 7(1):67-72 (Apr. 15, 1993) (with English translation)(22 pages).
- ENews, "Cholesterol Crystals Induce Atherosclerosis-Associated Inflammation in Mice," 1-4 (Jun. 14, 2010)(4 pages).
- Engler, et al., "Docosahexaenoic acid restores endothelial function in children with hyperlipidemia: results from the EARLY Study," *International Journal of Clinical Pharmacology and Therapeutics*, vol. 42—No. Dec. 2004 (672-679). (2004).
- Engler, M.B., et al., "Mechanisms of vasorelaxation induced by eicosapentaenoic acid (20:5n-3) in WKY rat aorta." *British Journal of Pharmacology* 131:1793-1799 (2000).
- Engler, M.M., et al., "The effects of a diet rich in docosahexaenoic acid on organ and vascular fatty acid composition in spontaneously hypertensive rats." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 61(5):289-295 (1999).
- Ennis JL, Cromwell WC. Clinical utility of low-density lipoprotein particles and apolipoprotein B in patients with cardiovascular risk. *J. Fam. Pract.* 2013;62:1-8.
- Epadel—PubChem CID 9831415, Retrieved on Apr. 9, 2014 [Retrieved from the Internet] <URL: <http://pubchem.ncbi.nlm.nih.gov/compound/9831415>> (19 pages).
- Epadel 1990 and JELIS Study (4 pages).
- Epadel Capsules 300, Japan Pharmaceutical Reference 369-371 (2nd ed.) (1991). (5 pages).
- Epadel drug information brochure (2000), certified English translation(36 pages).
- Epadel Package Insert 2007 (with Translation)(6 pages).
- Epadel Summary of Product Characteristics (SPC), Mochida Pharmaceutical Co., Ltd. Tokyo, Japan, Oct. 2013.
- Epadel® [Complete prescribing information]. Update (Version 5). Tokyo, Japan: Mochida Pharmaceutical; Jan. 2007 (9 pages).
- Epanova® (omega-3-carboxylic acids) capsules, for oral use, Prescribing information, 5 pgs., AstraZeneca Pharmaceuticals LP, (Revised: Mar. 2017)(5 pages).
- Eritsland J, Arnesen H, Gronseth K, et al. Effect of dietary supplementation with n-3 fatty acids on coronary artery bypass graft patency. *Am. J. Cardiol.* Jan. 1996 77 (1): 31-6.
- Eritsland J, Arnesen H, Seljeflot I, et al. Long-term effects of n-3 polyunsaturated fatty acids on haemostatic variables and bleeding episodes in patients with coronary artery disease. *Blood Coagul. Fibrinolysis* Feb. 6, 1995 (1): 17-22.
- Errata to the FDA Briefing Document Endocrinologic and Metabolic Drug Advisory Committee Meeting Oct. 16, 2013, 1 page.
- Esposito, "Effect of a Mediterranean-Style Diet on Endothelial Dysfunction and Markers of Vascular Inflammation in the Metabolic Syndrome: A Randomized Trial", *Journal of the American Medical Association*, 2004, 292(12), 1440-1446.
- Essentialis Inc. press release, "Essentialis Meets Primary Endpoint in Phase 2b Trial of DCCR for Treatment of Hypertriglyceridemia and is Granted Extensive Patent Coverage in the US," PR Newswire (May 17, 2009)(2 pages).
- Exhibit A to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 48 pages (Dec. 5, 2014).
- Exhibit B to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit C to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 14 pages (Dec. 5, 2014).
- Exhibit D to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 19 pages (Dec. 5, 2014).
- Exhibit E to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit F to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 10 pages (Dec. 5, 2014).
- Exhibit G to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 21 pages (Dec. 5, 2014).
- Exhibit H to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 10 pages (Dec. 5, 2014).
- Exhibit I to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 14 pages (Dec. 5, 2014).
- Exhibit J to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 13 pages (Dec. 5, 2014).
- Exhibit K to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 5 pages (Dec. 5, 2014).
- Exhibit L to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 5 pages (Dec. 5, 2014).
- Exhibit M to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 7 pages (Dec. 5, 2014).
- Exhibit N to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 15 pages (Dec. 5, 2014).
- Exhibit O to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 6 pages (Dec. 5, 2014).
- Exhibit P to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 17 pages (Dec. 5, 2014).
- Exhibit Q to Defendants' Joint Invalidity Contentions, 3:14-CV-02550-MLC-TJB (D.N.J.), 64 pages (Dec. 5, 2014).
- Faggini, E., et al., "Fish Oil Supplementation Prevents Neointima Formation in Nonhypercholesterolemic Balloon-Injured Rabbit Carotid Artery by Reducing Medial and Adventitial Cell Activation." *Arterioscler. Thromb. Vasc. Biol.*, 20:152-163 (2000).
- FDA Briefing Document, Endocrinologic and Metabolic Drugs Advisory Committee Meeting, dated Oct. 16, 2013, available publicly at least as of Oct. 16, 2013, 115 pages.
- FDA News Release, "FDA approves new orphan drug Kynamro to treat inherited cholesterol disorder," U.S. Food and Drug Administration, Protecting and Promoting Your Health (Jan. 29, 2013)(2 pages).
- Fer, M., et al., "Metabolism of eicosapentaenoic and docosahexaenoic acids by recombinant human cytochromes P450." *Archives of Biochemistry and Biophysics* 471:116-125 (2008).
- Ferns, G., et al., "Investigation and management of hypertriglyceridaemia." *J. Clin. Pathol.* 61:1174-1183 (2008).
- Feron O, Dessy C, Desager JP, Balligand JL. Hydroxy-methylglutaryl-coenzyme A reductase inhibition promotes endothelial nitric oxide synthase activation through a decrease in caveolin abundance. *Circulation.* 2001;103:113-118.
- Final Agenda for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Final Meeting Roster for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 2 pages.
- Final Questions for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 1 page.
- Finnen et al., "Purification and characterisation of phospholipase A2 from human epidermis," *Biochemical Society Trans*, 19(2):91S, 1991.
- Fischer, R., et al., "Dietary n-3 polyunsaturated fatty acids and direct renin inhibition improve electrical remodeling in a model of high human renin hypertension." *Hypertension* 51:540-546 (2008).
- Fisher et al., *Journal of Biological Chemistry* (2001) 276(3) 27855-27863.

US 10,568,861 B1

Page 13

(56)

References Cited

OTHER PUBLICATIONS

- Flaten H., et al., "Fish-oil concentrate: effects on variables related to cardiovascular disease." *Am. J. Clin. Nutr.* 52:300-306 (1990).
- Food and Drug Administration (FDA), (2005) *NLASPAN niacin extended release tablets*.
- Food and Drug Administration (FDA), (2005) *Tablets ZOCOR® (SIMVASTATIN)*.
- Ford, E.S. et al., "Hypertriglyceridemia and Its Pharmacologic Treatment Among US Adults." *Arch. Intern. Med.*, 169(6): 572-78 (2009).
- Fraker TD, Fihn SD. Writing on behalf of the 2002 Chronic Stable Angina Writing Committee. 2007 chronic angina focused update of the ACC/AHA guidelines for the management of patients with chronic stable angina. A Report of the ACC/AHA Task Force on Practice Guidelines. *Circulation* 50:2264-2274, Dec. 4, 2007.
- Frangou et al., "Efficacy of ethyl-eicosapentaenoic acid in bipolar depression: randomised double-blind placebo-controlled study," *British Journ. Psychiatry*, 188, 46-50 (2006).
- Frey R, Muck W, Kirschbaum N, et al. Riociguat (BAY 63-2521) and warfarin: a pharmacodynamic and pharmacokinetic interaction study. *J. Clin. Pharmacol.* Jul. 2011 51 (7): 1051-60.
- Frick, MH, et al., "Helsinki Heart Study. Primary prevention trial with gemfibrozil in middle-aged men with dyslipidaemia. Safety of treatment, changes in risk factors and incidence of coronary heart disease", *N. Eng. J. Med.*, 317:1237-1245, (1987).
- Friedewald, W.T., et al., "Estimation of the concentration of low-density lipoprotein cholesterol in plasma, without use of the preparative ultracentrifuge." *Clin Chem.*, 18:499-502 (1972).
- Friedman, A. N., et al., "Fish Consumption and Omega-3 Fatty Acid Status and Determinants in Long-Term Hemodialysis." *Amer. J. Kidney Diseases*, 47(6):1064-1071 (2006).
- Frøyland et al., "Chronic administration of eicosapentaenoic acid and docosahexaenoic acid as ethyl esters reduced plasma cholesterol and changed the fatty acid composition in rat blood and organs." *Lipids* 31(2):169-78 (Feb. 1996).
- Frøyland, L., et al., "Hypotriacylglycerolemic component of fish oil." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 57 (4 & 5):387-388 (1997).
- Furuta T, Shirai N, Sugimoto M, et al. Influence of CYP2C19 pharmacogenetic polymorphism on proton pump inhibitor-based therapies. *Drug Metab. Pharmacokinet* Jun. 20, 2005 (3): 153-67.
- Futata et al., "Effect of Eicosapentaenoic Acid (EPA) Formulation on Glucose Metabolism in Non-Insulin Dependent Diabetic Patients," *Journal of Clinical and Experimental Medicine* 169(8):889-890 (May 21, 1994)(English translation, 4 pages).
- Galan P, Kesse-Guyot E, Czernichow S, et al. Effects of B vitamins and omega 3 fatty acids on cardiovascular diseases: a randomised placebo controlled trial. *Br Med J. Nov. 29, 2010*;341:c6273.
- Galeano NF, Al-Haideri M, Keyserman F, Rumsey SC, Deckelbaum RJ. Small dense low density lipoprotein has increased affinity for LDL receptor-independent cell surface binding sites: a potential mechanism for increased atherogenicity. *J. Lipid Res.* 1998;39(6):1263-1273.
- Gallagher et al., "Germline BRCA Mutations Denote a Clinicopathologic Subset of Prostate Cancer," *Amer. Assoc. Cancer Res. Clin Cancer Res.*, 16(7):2115-21 (Apr. 1, 2010).
- Ganda OP, Bhatt DL, Mason RP, Miller M, Boden WE. Unmet need for adjunctive dyslipidemia therapy in hypertriglyceridemia management. *J Am Coll Cardiol* 72(3):330-43 (publication date Jul. 17, 2018).
- Garber AJ, Abrahamson MJ, Barzilay JI, et al. American Association of Clinical Endocrinologists' comprehensive diabetes management algorithm 2013 consensus statement. *Endocr. Pract.* 2013;19(suppl 2):1-48.
- Gardner CD, Fortmann SP, Krauss RM. Association of small low-density lipoprotein particles with the incidence of coronary artery disease in men and women. *JAMA.* 1996;276(11):875-881.
- Garg, R., et al., "Niacin treatment increases plasma homocyst(e)ine levels", *Am. Heart. J.*, 138:1082-1087, (1999).
- Garnett, "Interactions with Hydroxymethylglutaryl-coenzyme A reductase inhibitors," *Am J Health-Sys Pharm* vol. 52, 1639-1645, (Aug. 1, 1995).
- Geleijnse JM, Giltay EJ, Grobbee DE, Donders ART, Kok FJ. Blood pressure response to fish oil supplementation: meta-regression analysis of randomized trials. *J Hypertens.* Aug. 2002;20(8):1493-1499.
- Genest, JJ, et al., "Familial lipoprotein disorders in patients with premature coronary artery disease", 85:2025-2033, (1992).
- Geppert, et al. "Microalgal docosahexaenoic acid decreases plasma triacylglycerol in normolipidaemic vegetarians: a randomized trial." *British Journal of Nutrition*, 95, 779-786. (2006).
- Gillet L, Roger S, Bougnoux P, Le Guennec JY, Besson P. Beneficial effects of omega-3 long-chain fatty acids in breast cancer and cardiovascular diseases: voltage-gated sodium channels as a common feature? *Biochim. Jan. 2011* (epub Feb. 16, 2010); 93:4-6.
- Gillies, et al. "Effect of a Novel Eicosapentaenoic Acid-Rich Oil on Serum Cholesterol in Man," DuPont 2010.
- Ginsberg HN, Elam MB, Lovato LC, Crouse JR, 3rd, Leiter LA, Linz P, Friedewald WT, Buse JB, Gerstein HC, Probstfield J, Grimm RH, Ismail-Beigi F, Bigger JT, Goff DC, Jr., Cushman WC, Simons-Morton DG, Byington RP. Effects of combination lipid therapy in type 2 diabetes mellitus. *N. Engl. J. Med.* Apr. 29, 2010;362:1563-1574.
- Ginsberg HN, Elam MB, Lovato LC, et al, for the ACCORD Study Group. Effects of combination lipid therapy in Type 2 diabetes mellitus. *N Engl J Med* 362:1563-1574, 2010.
- Ginsberg HN. "Hypertriglyceridemia: new insights and new approaches to pharmacologic therapy," *Am J Cardiol*, 87:1174-1180 (2001).
- Girotti A W. Lipid hydroperoxide generation, turnover, and effector action in biological systems. *J. Lipid Res.* 1998;39(8):1529-1542.
- GISSI-HF Investigators. Effect of n-3 polyunsaturated fatty acids in patients with chronic heart failure (the GISSI-HF trial): a randomised, double-blind, placebo-controlled trial. *Lancet.* Oct. 4, 2008 (epub Aug. 29, 2008); 372(9645):1223-1230.
- Gissi-Prevenzione Investigators, "Dietary Supplementation with n-3 Polyunsaturated Fatty Acids and Vitamin E after Myocardial Infarction: Results of the GISSI-Prevenzione Trial", *The Lancet*, 354:447-455, (Aug. 7, 1999).
- Glod, "Recent Advances in the Pharmacotherapy of Major Depression", *Arch. Psychiatr. Nurs.*, 10(6):355-364 (Dec. 1996).
- Goff DC, Lloyd-Jones DM, Bennett G, et al. ACC/AHA Prevention Guideline: 2013 ACC/AHA Guideline on the Assessment of Cardiovascular Risk: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. *Circulation.* Jun. 24, 2014 (epub Nov. 12, 2013); 129:S74-S75.
- Goldberg, A C: "Combination therapy of dyslipidemia," *Current Treatment Options in Cardiovascular Medicine* Aug. 2007 GB, vol. 9, No. 4, pp. 249-258 (2007).
- Goodman & Gilman (Robert W. Mahley & Thomas P. Bersot) *Drug Therapy for Hypercholesterolemia and Dyslipidemia*, in Goodman & Gilman's *The Pharmacological Basis for Therapeutics* 971 (Hardman et al., eds 10th ed. 2001)(32 pages).
- Gordon, DJ. et al., High density lipoprotein cholesterol and cardiovascular disease: four prospective American studies. *Circulation.* 79: 8-15. (1989).
- Gorritz JL et al., "Rhabdomyolysis and Acute Renal Failure Associated with Gemfibrozil Therapy," *Nephron* 74(2): 437-438 (1996).
- Gorritz, JL, "Rhabdomyolysis and Acute Renal Failure Associated with Bezafibrate Treatment," *Nephrol Dial Transplant* 10(12):2371-2372 (1995).
- Gosai, P. et al. Effect of omega-3-acid ethyl esters on the steady-state plasma pharmacokinetics of rosuvastatin in healthy adults. *Expert Opin. Pharmacother.* 9, 2947-2953 (2008).
- Goto, Y. et al., "Clinical Pharmacological Trial of Ethyl Icosapentate (MND-21)—Dose Finding Study." *Journal of Clinical Therapeutic & Medicines* 8:1293-309 (1992).
- Gould, A.L., et al., "Cholesterol reduction yields clinical benefit: impact of statin trials." *Circulation*, 97:946-952 (1998).
- Greenblatt DJ, von Moltke LL. Interaction of warfarin with drugs, natural substances, and foods. *J. Clin. Pharmacol.* Feb. 2005 45 (2): 127-32.
- Grenyer, Brin F.S., et al., "Fish Oil Supplementation in the Treatment of Major Depression: A Randomised Double-Blind Placebo-

US 10,568,861 B1

Page 14

(56)

References Cited

OTHER PUBLICATIONS

- Controlled Trial", *Progress in Neuro-Psychopharmacology & Biological Psychiatry*, 31:1393-1396, (2007).
- Griffin, M.D., et al., "Effects of altering the ratio of dietary n-6 to n-3 fatty acids on insulin sensitivity, lipoprotein size, and postprandial lipemia in men and postmenopausal women aged 45-70 y: the OPTILIP Study." *Am J Clin Nutr* 84:1290-8 (2006).
- Grimsgaard et al., "Effects of Highly Purified Eicosapentaenoic Acid and Docosahexaenoic Acid on Hemodynamics in Humans" *American Society for Clinical Nutrition*, 68:52-9, (1998).
- Grimsgaard, Kaare H. Bonaa, John-Bjarne Hansen, and Arne Nordoy, "Highly purified eicosapentaenoic acid and docosahexaenoic acid in humans have similar triacylglycerol-lowering effects but divergent effects on serum fatty acids" *Am J Clin Nutr*, 66:649-659, (1997).
- Grundy S.M et al., Efficacy, safety, and tolerability of once-daily niacin for the treatment of dyslipidemia associated with type 2 diabetes: results of the Assessment of Diabetes Control and Evaluation of the Efficacy of Niaspan Trial. *Arch. Intern. Med.* 162: 1568-1576 (2002).
- Grundy SM, et al. Implications of Recent Clinical Trials for the National Cholesterol Education Program Adult Treatment Panel III Guidelines. *Circulation*. 2004; 110:227-39.
- Grundy, Scott M., "Low-Density Lipoprotein, Non-High-Density Lipoprotein, and Apolipoprotein B as Targets of Lipid-Lowering Therapy" *Circulation*. 106:2526-2529 (2002).
- Guallar, E., et al., "Omega-3 fatty acids in adipose tissue and risk of myocardial infarction—The EURAMIC study." *Arterioscler. Thromb. Vasc. Biol.*, 19:1111-1118 (1999).
- Guillot, et al., "Increasing intakes of the long-chain omega-3 docosahexaenoic acid: effects on platelet functions and redox status in healthy men," *The FASEV Journal*, vol. 23, pp. 2909-2916 (2009).
- Guizy, M., et al., "ω-3 and ω-6 Polyunsaturated fatty acids block HERG channels." *Am J Physiol Cell Physiol* 289:C1251-C1260 (2005).
- Gyarmathy, M., "Selection from the industrial manufacturing. 5th part: Gelatine capsules. 5/2 part: Soft gelatine capsules," *Gyógyszerészet*, vol. 38, No. 2, pp. 105-109 (1994) (with English summary).
- Hakonarson, H., et al., "Effects of a 5-lipoxygenase-activating protein inhibitor on biomarkers associated with risk of myocardial infarction—a randomized trial." *JAMA*, 293(8):2245-56 (May 11, 2005).
- Hall, W. L., et al., "A high-fat meal enriched with eicosapentaenoic acid reduces postprandial arterial stiffness measured by digital volume pulse analysis in healthy men." *J. Nutr.* 138: 287-291 (Feb. 2008).
- Hamazaki et al., "Docosahexaenoic Acid-Rich Fish Oil Does Not Affect Serum Lipid Concentrations of Normolipidemic Young Adults", *American Institute of Nutrition*, 126(11):2784-2789, Nov. 1996.
- Hamazaki et al., "Effects of fish oil rich in eicosapentaenoic acid on serum lipid in hyperlipidemic hemodialysis patients," *Kidney Int* 1, 26:81-84 (Jul. 1984).
- Hamazaki et al., "Effects of Orally Administered Ethyl Ester of Eicosapentaenoic Acid (EPA: C20:5, omega-3) on PG12-Like Substance Production by Rat Aorta" *Prostaglandins*, vol. 23 No. 4, pp. 557-567 (Apr. 1982).
- Hamazaki T. et al., "Reduction of microalbuminuria in diabetics by Eicosapentaenoic acid ethyl ester" *Lipids*. 25 (9):542-5 (Sep. 1990).
- Hampel H, Abraham NS, El-Se rag HB. Meta-analysis: obesity and the risk for gastroesophageal reflux disease and its complications. *Ann. Intern. Med.* Aug. 2005 143 (3): 199-211.
- Han, J. J., et al., "Enhancement of both reaction yield and rate of synthesis of structured triacylglycerol containing eicosapentaenoic acid under vacuum with water activity control." *Lipids* 34:989-995 (Sep. 1999).
- Hanasaki, K., et al., "Potent modification of low density lipoprotein by group X secretory phospholipase A2 is linked to macrophage foam cell formation." *J. Biol. Chem.* 277(32):29116-24 (Aug. 9, 2002).
- Haney, E.M., et al., "Screening for lipid disorders in children and adolescents; Systematic evidence review for the U.S. Preventive Services Task Force (evidence synthesis)." No. 47. Rockville, MD: Agency for Healthcare Research and Quality, US Department of Health and Human Services; AHRQ Publication No. 07-0598-EF-1; Jul. 2007. Available at: <http://www.uspreventiveservicestaskforce.org/uspstf07/chlipid/chlipidsyn.pdf>. (Accessed Mar. 23, 2011)(573 pages).
- Hannah, J., et al., "Effect of dietary fatty acids on LDL binding." *Ann N Y Acad Sci.*, 683:178-182 (Jun. 14, 1993).
- Hansen et al., "Comparative effects of prolonged intake of highly purified fish oils as ethyl ester or triglyceride on lipids, haemostasis and platelet function in normolipemic men." *Eur. J. Clin. Nutr.* 47(7):497-507 (Jul. 1993).
- Hansen, J.B., et al., "Effects of highly purified eicosapentaenoic acid and docosahexaenoic acid on fatty acid absorption, incorporation into serum phospholipids and postprandial triglyceridemia." *Lipids* 33:131-38 (Feb. 1998).
- Harada-Shiba et al., *Journal of Clinical and Experimental Medicine*, Jun. 30, 2007, vol. 221, No. 13, pp. 1068-1073 (with English translation).
- Harris WS. International recommendations for consumption of long-chain omega-3 fatty acids. *J Cardiovasc Med (Hagerstown)* 8(suppl 1):S50-S52, Sep. 2007.
- Harris, "n-3 Fatty acids and lipoproteins: a comparison of results from human and animal studies," *Lipids* 31, 243-252 (Mar. 1996).
- Harris, W. S. et al. "Safety and efficacy of Omacor in severe hypertriglyceridemia," *Journal of Cardiovascular Risk*, 4:385-391 (Oct.-Dec. 1997).
- Harris, W. S., "Fish oils and plasma lipid and lipoprotein metabolism in humans: a critical review." *J Lipid Res.* 30:785-807 (Jun. 1989).
- Harris, W. S., "The omega-3 index as a risk factor for coronary heart disease." *Am Clin Nutr* 87:1997S-2002S (Jun. 2008).
- Harris, W. S., et al., "n-3 Fatty acids and urinary excretion of nitric oxide metabolites in humans." *Am. J. Clin. Nutr.*, 65:459-64 (Feb. 1997).
- Harris, W. S., et al., "Influence of n-3 fatty acid supplementation on the endogenous activities of plasma lipases." *Am. J. Clin. Nutr.* 66:254-60 (Aug. 1997).
- Harris, W.S., "Expert opinion: omega-3 fatty acids and bleeding—cause for concern?" *The American Journal of Cardiology* 99(6A): 45C-46C (Mar. 19, 2007).
- Harris, W.S., "n-3 Fatty acids and human lipoprotein metabolism: an update." *Lipids* 34:S257-S258 (1999).
- Harris, W.S., "n-3 Fatty acids and serum lipoproteins: human studies." *Am J Clin Nutr* 65:1645S-54S (1997).
- Harris, W.S., "Omega-3 fatty acids in cardiac biopsies from heart transplantation patients." *Circulation* 110:1645-1649 (Sep. 21, 2004).
- Harris, W.S., et al., "Comparison of the effects of fish and fish-oil capsules on the n-3 fatty acid content of blood cells and plasma phospholipids." *Am J Clin Nutr* 86:1621-5 (Dec. 2007).
- Harris, W.S., et al., "Omega-3 fatty acids and coronary heart disease risk: Clinical and mechanistic perspectives." *Atherosclerosis* 197:12-24 (Mar. 2008)(epub Dec. 26, 2007).
- Harris, W.S., et al., "Stearidonic acid increases the red blood cell and heart eicosapentaenoic acid content in dogs." *Lipids* 42:325-333 (Apr. 2007)(epub Mar. 9, 2007).
- Harris, W.S., et al., "Tissue n-3 and n-6 fatty acids and risk for coronary heart disease events." *Atherosclerosis* 193:1-10 (Jul. 2007)(epub May 15, 2007).
- Hartweg, J., et al., "Potential impact of omega-3 treatment on cardiovascular disease in type 2 diabetes." *Curr Opin Lipidol.*, 20:30-38 (Feb. 2009).
- Hata et al, *Geriatric Medicine*, 30 (5), 799-852, 1992 (with English introduction).
- Hawthorne, et al., "High dose eicosapentaenoic acid ethyl ester: effects on lipids and neutrophil leukotriene production in normal volunteers." *Br. J. Clin. Pharmacol.*, vol. 30, 187-194 (Aug. 1990).
- Hayashi et al., Decreases in Plasma Lipid Content and Thrombotic Activity by Ethyl Icosapentate Purified from Fish Oiles, *Current Therapeutic Research*, vol. 56, No. 1, pp. 24-31 (1995).

(56)

References Cited

OTHER PUBLICATIONS

- Herbette L, Marquardt J, Scarpa A, Blasie JK. A direct analysis of lamellar x-ray diffraction from hydrated oriented multilayers of fully functional sarcoplasmic reticulum. *Biophys. J.* Nov. 1977;20(2):245-272.
- Hibbeln, J. R., et al., "Healthy intakes of n-3 and n-6 fatty acids: estimations considering worldwide diversity." *Am J Clin Nutr.* 83:1483S-93S (Jun. 2006).
- Higashihara et al. "Effects of Eicosapentaenoic Acid on Biochemical Failure after Radical Prostatectomy for Prostate Cancer," *in vivo* 24:561-566 (Jul./Aug. 2010).
- Hilpert, K.F., et al., "Postprandial effect of n-3 polyunsaturated fatty acids on apolipoprotein B—containing lipoproteins and vascular reactivity in type 2 diabetes." *Am J Clin Nutr* 85:369-76 (Feb. 2007).
- Hirafuji, M., et al., "Docosahexaenoic acid potentiates interleukin-1 beta induction of nitric oxide synthase through mechanism involving p44/42 MAPK activation in rat vascular smooth muscle cells." *British Journal of Pharmacology* 136:613-619 (Jun. 2002).
- Hirai, A., et al., "The effects of the oral administration of fish oil concentrate on the release and the metabolism of [¹⁴C] arachidonic acid and [¹⁴C] eicosapentaenoic acid by human platelets", *Thromb. Res.*, 28:285-298, (Nov. 1, 1982).
- Hirano T, Ito Y, Koba S, Toyoda M, Ikejiri A, Saegusa H, Yamazaki J, Yoshino G. Clinical significance of small dense low-density lipoprotein cholesterol levels determined by the simple precipitation method. *Arterioscler. Thromb. Vasc. Biol.* Mar. 2004;24(3):558-563.(epub Jan. 15, 2004).
- Hirano, R., et al., "Regulation by long-chain fatty acids of the expression of cholesteryl ester transfer protein in HepG2 cells." *Lipids*, 36:401-406 (Apr. 2001).
- Hofacer R, et al., Omega-3 fatty acid deficiency increases stearoyl-CoA desaturase expression and activity indices in rat liver: Positive association with non-fasting plasma triglyceride levels, Prostaglandins Leukot. Essent. Fatty Acids. Jan./Feb. 2012;86:71-7. (epub Nov. 1, 2011).
- Hoffman, "Atherosclerosis: Prevention through the Ages," WebMD, <https://www.webmd.com/heart/features/atherosclerosis-prevention-through-ages#1>, (Dec. 4, 2007).
- Hohenester, "Primary Biliary Cirrhosis," *Semin Immunopathol.* 31L:283-307, 285 (Sep. 2009)(epub Jul. 15, 2009).
- Holmeide, A. K., et al., "Oxidative degradation of eicosapentaenoic acid into polyunsaturated aldehydes." *Tetrahedron* 59:7157-7162 (2003).
- Holub, B.J., PhD, "Fish Oils and Cardiovascular Disease", Canadian Medical Association Journal, 141(10):1063 (Nov. 15, 1989).
- Holvoet P, Kritchevsky SB, Tracy RP, Mertens A, Rubin SM, Butler J, Goodpaster B, Harris TB. The metabolic syndrome, circulating oxidized LDL, and risk of myocardial infarction in wellfunctioning elderly people in the health, aging, and body composition cohort. *Diabetes.* Apr. 2004;53(4):1068-1073.
- Hom et al., "Soft Gelatin Capsules II: Oxygen Permeability Study of Capsule Shells," *J Pharm Sci.* (May 1975) 64(5):851-857.
- Hombek, M., et al., "Biosynthesis of the algal pheromone fucoseratene by the freshwater diatom *Asterionella formosa* (Bacillariophyceae)." *Tetrahedron* 54:11033-11042 (1998).
- Hong KN, Fuster V, Rosenson RS, Rosendorff C, Bhatt DL. How low to go with glucose, cholesterol, and blood pressure in primary prevention of CVD. *J Am Coll Cardiol* 70(17):2171-85 (publication date Oct. 24, 2017; epublication date Oct. 16, 2017).
- Hoogeveen EK, Geleijnse JM, Kromhout D, et al. No effect of n-3 fatty acids supplementation on NT-proBNP after myocardial infarction: the Alpha Omega Trial. *Eur J Prev Cardiol.* May 2015;22:648-55.
- Horrobin, D.F. The Phospholipid Concept of Psychiatric Disorders and its Relationship to the Neurodevelopmental Concept of Schizophrenia. In M. Peet (ed.) *Phospholipid Spectrum Disorder in Psychiatry* pp. 1-19 (1999).
- Hoskins et al., "Combination use of statins and omega-3 fatty acids: an emerging therapy for combined hyperlipidemia," *Abstract*, 1(5): 579-591(13) (2006).
- Howe, P.R.C., et al., "Equal antithrombotic and triglyceride-lowering effectiveness of eicosapentaenoic acid-rich and docosahexaenoic acid-rich fish oil supplements." *Lipids* 34:S307-S308 (1999).
- HPS2-thrive Collaborative Group, "randomized placebo-controlled trial in 25 673 high-risk patients of er niacin/laroprant: Trial design, pre-specified muscle and liver outcomes, and reasons for stopping study treatment." *Eur. Heart J.* May 2013;34:1279-1291.
- HPS2-Thrive Collaborative Group, Landray MJ, Haynes R, et al. Effects of extended-release niacin with laropirant in high-risk patients. *M Engl J Med.* Jul. 17, 2014; 371(3):203-12.
- Hruska MW, Amico JA, Langae TY, Ferrell RE, Fitzgerald SM, Frye RF. The effect of trimethoprim on CYP2C8 mediated rosiglitazone metabolism in human liver microsomes and healthy subjects. *Br. J. Clin. Pharmacol.* Jan. 2005;59:70-79.
- Hughes et al., "Fish oil produces an atherogenic lipid profile in hypertensive men," *Atherosclerosis*, 84, pp. 229-237 (Oct. 1990).
- Hulthe J, Hulthen LM, Fagerberg B. Low adipocyte-derived plasma protein adiponectin CJ concentrations are associated with the metabolic syndrome and small dense low-density lipoprotein particles: atherosclerosis and insulin resistance study. *Metab. Clin. Exp.* Dec. 2003;52(12):1612-1614.
- Huntington's Disease Drug Works—The DHA Dilemma http://hddrugworks.org/index2.php?option=com_content&task=view&id=185&pop=1&pa... Printed on Aug. 22, 2008.(2 pages).
- Ignarro LJ, Buga GM, Wood KS, Byrnes RE, Chaudhuri G. Endothelium-derived relaxing factor produced and released from artery and vein is nitric oxide. *Proc. Natl. Acad. Sci. USA.* Dec. 1987;84:9265-9269.
- Illingworth, DR, et al., "Comparative effects of lovastatin and niacin in primary hypercholesterolemia: A prospective trial", *Arch. Int. Med.*, 154:1586-1595, (Jul. 25, 1994).
- Inoue, I., et al., "Expression of peroxisome proliferator-activated receptor α (PPAR α) in primary cultures of human vascular endothelial cells." *Biochem. Biophys. Res. Comm.*, 246, 370-374 (May 19, 1998).
- Inzucchi et al., "Diagnosis of Diabetes," *New Engl. Journ of Med.*, 367(6):541-550 (Aug. 9, 2012).
- Ishida, Y., et al., "α-Lipoic Acid and Insulin Autoimmune Syndrome." *Diabetes Care*, 30(9): 2240-41 (Sep. 2007).
- Isley, et al., "Pilot study of combined therapy with ω -3 fatty acids and niacin in atherogenic dyslipidemia," *Journal of Clinical Lipidology*, 1, 211-217 (Jul. 2007).
- Itoh et al., "Increased adiponectin secretion by highly purified eicosapentaenoic acid in rodent models of obesity and human obese subjects," *Arterioscler. Thromb. Vasc. Biol.*, pp. 1918-1925 (together with online Supplements 1-15) (Sep. 2007)(epub Jun. 14, 2007).
- Jacob RF, Mason RP. Lipid peroxidation induces cholesterol domain formation in model membranes. *J. Biol. Chem.* Nov. 25, 2005;280(47):39380-39387.(epub Sep. 28, 2005).
- Jacob RF, Walter MF, Self-Medlin Y, Mason RP. Atorvastatin active metabolite inhibits oxidative modification of small dense low-density lipoprotein. *J. Cardiovasc. Pharmacol.* Aug. 2013;62(2):160-166.
- Jacobson et al. "Hypertriglyceridemia and Cardiovascular Risk Reduction", *Clinical Therapeutics*, vol. 29 pp. 763-777 (May 2007).
- Jacobson TA. Opening a new lipid "apo-theary": incorporating apolipoproteins as potential risk factors and treatment targets to reduce cardiovascular risk. *Mayo Clin. Proc.* Aug. 2011;86:762-780.
- Jacobson, T. Secondary Prevention of Coronary Artery Disease with Omega-3 Fatty Acids. *Am J Cardiol*; 98 [suppl]: 61i-70i (Aug. 21, 2006).
- Jacobson, T.A., "Role of n-3 fatty acids in the treatment of hypertriglyceridemia and cardiovascular disease." *Am J Clin Nutr* 87:1981S-90S (Jun. 2008).

US 10,568,861 B1

Page 16

(56)

References Cited

OTHER PUBLICATIONS

- Jacobson, T.A., et al., "Effects of eicosapentaenoic acid and docosahexaenoic acid on low-density lipoprotein cholesterol and other lipids: A review." *J. Clin. Lipidology*, vol. 6, pp. 5-18 (Jan./Feb. 2012).
- Jakus V, Rietbrock N. Advanced glycation end-products and the progress of diabetic vascular complications. *Physiol. Res.* 2004;53(2): 131-142.
- Jenner, "Presymptomatic Detection of Parkinson's Disease." *J. Neural Transm. Suppl.*, 40:23-36. (Abstract only) (1993).
- Jialal I, Devaraj S. Antioxidants and atherosclerosis: Don't throw out the baby with the bath water. *Circulation*. Feb. 25, 2003;107:926-928.
- Jialal, I. "Editorial: Remnant lipoproteins: measurement and clinical significance." *Clinical Chemistry* 48(2):217-219 (Feb. 2002).
- Jinno Y, Nakakuki M, Kawano H, Notsu T, Mizuguchi K, Imada K. Eicosapentaenoic acid administration attenuates the pro-inflammatory properties of VLDL by decreasing its susceptibility to lipoprotein lipase in macrophages. *Atheroscler.* Dec. 2011;219:566-572. (epub Oct. 4, 2011).
- Jong et al., "Role of ApoCs in Lipoprotein Metabolism: Function Differences Between ApoC1, ApoC2, and ApoC3." *Arteriosclerosis, Thrombosis and Vascular Biology*. (Mar. 1999) 19(3):472-484.
- Jørgensen AB, Frikke-Schmidt R, Nordestgaard BG, Tybjaerg-Hansen A. Loss-of-function mutations in APOC3 and risk of ischemic vascular disease. *N Engl J Med*. Jul. 3, 2014; 371(1):32-41 (epub Jun. 18, 2014).
- Journal of Practical Pharmacy, "Hyperlipidemia Drug," 58(4):1303-1324 (2007) (with English abstract).
- Journal of the Japan Diabetes Society, "The Relationship Between Postprandial ApoB48 Increase and Insulin Resistance in Type-2 Diabetes," 55(Suppl. 1):S310 (Apr. 2012) (with English Translation) (2 pages).
- Journal of the Japanese Diabetes Society, "A Case of Familial Combined Hyperlipidemia Associated with Obesity, Type 2 Diabetes Mellitus and Severe Hypertriglyceridemia," 51(3), pp. 233-237 (Mar. 30, 2008) (with English abstract).
- Jun M, Foote C, Lv J, et al. Effects of fibrates on cardiovascular outcomes: a systematic review and meta-analysis. *Lancet* 375 (9729):1875-1884, May 29, 2010 (epub May 10, 2010).
- Jung, U.J. et al., "n-3 Fatty acids and cardiovascular disease: mechanisms underlying beneficial effects." *Am J Clin Nutr* 87: 2003S-9S (Jun. 2008).
- Kamanna et al., "Mechanism of Action of Niacin," *The American Journal of Cardiology* (Apr. 17, 2008), 101(8), S20-S26.
- Kamido et al., Lipid Composition of Platelets from Patients with Atherosclerosis: Effect of Purified Eicosapentaenoic Acid Ethyl Ester Administration, Oct. 1988, *Lipids*, 23, pp. 917-923 [Abstract only] (7 pages).
- Kaminski WE, Jendraschak E, Kieff R, et al. Dietary omega-3 fatty acids lower levels of platelet-derived growth factor mRNA in human mononuclear cells. *Blood* Apr. 1993, 81(7): 1871-9.
- Kanayasu, T., et al., "Eicosapentaenoic acid inhibits tube formation of vascular endothelial cells in vitro." *Lipids* 26:271-276 (Apr. 1991).
- Kastelein et al., Omega-3 Free Fatty Acids for the Treatment of Severe Hypertriglyceridemia: The EpanoVa for Lowering Very High Triglycerides (EVOLVE) Trial, *J. Clin. Lipidol.* (JACL 597) Jan./Feb. 2014 (epub Oct. 14, 2013).
- Katan, M. B., et al., "Kinetics of the incorporation of dietary fatty acids into serum cholesteryl esters, erythrocyte membranes, and adipose tissue: an 18-month controlled study." *J. Lipid Res.* 38: 2012-2022 (Oct. 1997).
- Katayama et al., Effect of long-term administration of ethyl eicosapentate (EPA-E) on local cerebral blood flow and glucose utilization in stroke-prone spontaneously hypertensive rats (SHRSP), *Brain Research*, vol. 761, pp. 300-305 (Dec. 31, 1997).
- Katayama et al., "Efficacy and Safety of Ethyl Icosapentate (Epadel) Given for a Long Term Against Hyperlipidemia," *Prog. Med.*, 21:457-467 (2001) (with English translation).
- Kato, T., et al., "Palmitate impairs and eicosapentaenoate restores insulin secretion through regulation of SREBP-1c in pancreatic islets." *Diabetes*, 57(9):2382-2392 (2008) (published online May 5, 2008.).
- Kawamura et al., "Effects of 4 weeks' intake of polyunsaturated fatty acid ethylester rich in eicosapentaenoic acid (ethylester) on plasma lipids, plasma and platelet phospholipid fatty acid composition and platelet aggregation; a double blind study," *Nihon Naika Gakkai Zasshi*, 72(1):18-24 (Jan. 10, 1983).
- Kawano, H., et al., "Changes in aspects such as the collagenous fiber density and foam cell size of atherosclerotic lesions composed of foam cells, smooth muscle cells and fibrous components in rabbits caused by all-cis 5, 8, 11, 14, 17-icosapentaenoic acid," *J. Atheroscler. Thromb.*, 9:170-177, (2002).
- Kawashima, H., et al., "Oral Administration of Dihomo- γ -Linolenic Acid Prevents Development of Atopic Dermatitis in NC/Nga Mice." *Lipids* 43:37-43 (Jan. 2008) (epub Nov. 6, 2007).
- Keech A, Simes RJ, Barter P, Best J, Scott R, Taskinen MR, Forde P, Pillai A, Davis T, Glasziou P, Drury P, Kesaniemi Y A, Sullivan D, Hunt D, Colman P, d'Emden M, Whiting M, Ehnholm C, Laakso M. Effects of long-term fenofibrate therapy on cardiovascular events in 9795 people with type 2 diabetes mellitus (the FIELD study): Randomised controlled trial. *Lancet*. Nov. 26, 2005;366:1849-1861.
- Kelley, D. S., et al., "Docosahexaenoic Acid Supplementation Decreases Remnant-Like Particle-Cholesterol and Increases the (n-3) Index in Hypertriglyceridemic Men." *J. Nutr.* 138: 30-35 (Jan. 2008).
- Kelley, et al., "Docosahexaenoic acid supplementation improves fasting and postprandial lip profiles in hypertriglyceridemic men." *The American Journal of Clinical Nutrition*, 86: 324-333 (Aug. 2007).
- Kellner-Weibel G, Yancey PG, Jerome WG, Walser T, Mason RP, Phillips MC, Rothblat GH. Crystallization of free cholesterol in model macrophage foam cells. *Arterioscler. Thromb. Vasc. Biol.* Aug. 1999;19(8):1891-1898.
- Kendall BJ, Macdonald GA, Hayward NK, et al. The risk of Barrett's esophagus associated with abdominal obesity in males and females. *Int. J. Cancer* May 2013 132 (9): 2192-9.
- Kerr, S., Brosnan MJ, McIntyre M, Reid JL, Dominiczak AF, Hamilton CA. Superoxide anion production is increased in a model of genetic hypertension role of the endothelium. *Hypertension*. Jun. 1999;33:1353-1358.
- Kew, S., et al., "Effects of oils rich in eicosapentaenoic and docosahexaenoic acids on immune cell composition and function in healthy humans." *Am J Clin Nutr* 79:674-81 (Apr. 2004).
- Kholodov et al., "Clinical Pharmacokinetics," *M. Medicine*. (1985) pp. 89-98, 134-138, 160, 378-380 [with English Summary] (27 pages).
- Khoueiry G, Rafeh NA, Sullivan E, et al. Do omega-3 polyunsaturated fatty acids reduce risk of sudden cardiac death and ventricular arrhythmias? A meta-analysis of randomized trials. *Heart and Lung*. Jul./Aug. 2013;42:251-256. (epub May 25, 2013).
- Kim F, Tysseling KA, Rice J, Gallis B, Haji L, Giachelli CM, Raines EW, Corson MA, Schwartz MW. Activation of IKK β by glucose is necessary and sufficient to impair insulin signaling and nitric oxide production in endothelial cells. *J. Mol. Cell. Cardiol.* Aug. 2005;39(2):327-334.
- Kim KA, Park PW, Kim HK, Ha JM, Park JY. Effect of quercetin on the pharmacokinetics of rosiglitazone, a CYP2C8 substrate, in healthy subjects. *J. Clin. Pharmacol.* Aug. 2005;45:941-946.
- Kimura, F., et al., "Long-term supplementation of docosahexaenoic acid-rich, eicosapentaenoic acid-free microalgal oil in n-3 fatty acid-deficient rat pups." *Biosci. Biotechnol. Biochem.*, 72(2):608-610 (Feb. 2008).
- Kinoshita, "Anti-hyperlipidemic agents," *Nihon Rinsho*, 60(5):968-74 (May 2002) (with English Abstract) (11 pages).
- Kinsella, J.E., et al., "Dietary n-3 polyunsaturated fatty acids and amelioration of cardiovascular disease: possible mechanisms." *Am J Clin Nutr* 52:1-28 (Jul. 1990).
- Kitada, 9th Diabetes Drug and Drug Related Seminar Diabetes Q&A, Kanazawa Medical University, Diabetes and Endocrine Internal Medicine (<http://plaza.umin.ac.jp/iby/etcdata/yakuyaku110410.pdf>) (Apr. 10, 2011) (with English translation) (105 pages).

US 10,568,861 B1

Page 17

(56)

References Cited

OTHER PUBLICATIONS

- Klempfner R, Erez A, Sagit BZ, et al. Elevated triglyceride level is independently associated with increased all-cause mortality in patients with established coronary heart disease: Twenty-two-year follow-up of the Bezafibrate Infarction Prevention Study and Registry. *Circ Cardiovasc Qual Outcomes* 9(2):100-8 (publication date Mar. 8, 2016).
- Knapp HR. Dietary fatty acids in human thrombosis and hemostasis. *Am. J. Clin. Nutr.* May 1997 65 (5 Suppl): 1687S-98S.
- Knopp, R.H., et al., "Contrasting effects of unmodified and time-release forms of niacin on lipoproteins in hyperlipidemic subjects: clues to mechanism of action of niacin", *Metabolism*, 34:642-650, (Jul. 1985).
- Koba S, Hirano T, Ito Y, Tsunoda F, Yokota Y, Ban Y, Iso Y, Suzuki H, Katagiri T. Significance of small dense low-density lipoprotein-cholesterol concentrations in relation to the severity of coronary heart diseases. *Atherosclerosis*. Nov. 2006;189(1):206-214. (epub Jan. 18, 2006).
- Kohn, M., et al., "Inhibition by Eicosapentaenoic Acid of Oxidized-LDL- and Lysophosphatidylcholine-Induced Human Coronary Artery Smooth Muscle Cell Production of Endothelin." *J. Vasc. Res.* 38:379-388 (Jul./Aug. 2001).
- Kojda G, Harrison DG. Interactions between no and reactive oxygen species: Pathophysiological importance in atherosclerosis, hypertension, diabetes and heart failure. *Cardiovasc. Res.* Aug. 15, 1999;43:562-571.
- Kojima, T, et al., "Long-term administration of highly purified eicosapentaenoic acid provides improvement of psoriasis." *Dermatologica*, 182:225-230 (1991).
- Koroshetz, W.J. Huntington's Disease. In Samuels, M. (ed.) *Office Practice of Neurology*, pp. 654-661 (1996).
- Kosonen, O., et al., "Inhibition by nitric oxide-releasing compounds of E-selectin expression in and neutrophil adhesion to human endothelial cells." *European Journal of Pharmacology* 394:149-156 (Apr. 7, 2000).
- Koyama et al., Plaque Reduction and Stabilization Observed in Borderline Diabetes Using Coronary CT Angiogram During Administration of Purified Eicosapentaenoic Acid (EPA) *Ther. Res.* 31(2):219-225 (Feb. 2010) (with English translation)(20 pages).
- Krauss RM. Heterogeneity of plasma low-density lipoproteins and atherosclerosis risk. *Curr. Opin. Lipidol.* Oct. 1994;5(5):339-349.
- Kris-Etherton, et al., "Fish Consumption, Fish Oil, Omega-3 Fatty Acids, and Cardiovascular Disease" *Circulation*, 106:2747-2757 (Nov. 19, 2002)(epub Jan. 28, 2003).
- Kris-Etherton, P. M., et al., "Omega-3 Fatty Acids and Cardiovascular Disease—New Recommendations From the American Heart Association." *Arterioscler Thromb Vasc Biol.* 23:151-152 (Feb. 1, 2003).
- Krzynowek et al., "Purification of Omega-3 Fatty Acids from Fish Oils Using HPLC: An Overview," *National Marine Fisheries—Proceedings of the first joint conference of the Tropical and Subtropical Fisheries Technological Society of the Americas with the Atlantic Fisheries Technological Society*, pp. 74-77 (1988).
- Ku, K., et al., "Beneficial Effects of to-3 Fatty Acid Treatment on the Recovery of Cardiac Function After Cold Storage of Hyperlipidemic Rats." *Metabolism*, 48(10):123-1209 (Oct. 1999).
- Kunimoto M, Inoue K, Nojima S. Effect of ferrous ion and ascorbate-induced lipid peroxidation on liposomal membranes. *Biochem. Biophys. Acta.* Aug. 6, 1981;646(1):169-178.
- Kurabayashi, T., et al., "Eicosapentaenoic acid effect on hyperlipidemia in menopausal Japanese women. The Niigata Epadel Study Group" *Obstet Gynecol* 96:521-8 (Oct. 2000).
- Labor Diagnostik Karlsruhe, "Target Values of Lipid Metabolism [Recommendation for lipid plasma levels in Germany]," (exact publication date unknown; circa 2006) (with English abstract)(4 pages).
- Lada et al., "Associations of Low Density Lipoprotein Particle Compositions with Atherogenicity," *Curr. Opin. Lipidol.* (Feb. 2004) 15(1):19-24.
- Lai, E., et al., "Suppression of niacin-induced vasodilation with an antagonist to prostaglandin D2 receptor subtype 1", *Clin. Pharm. & Ther.*, 81:849-857, (Jun. 2007/epub Mar. 28, 2007).
- Laidlaw, M., et al., "Effects of supplementation with fish oil-derived n-3 fatty acids and γ -linolenic acid on circulating plasma lipids and fatty acid profiles in women." *Am J Clin Nutr* 77:37-42 (Jan. 2003).
- Laird et al., "Relationship of early hyperglycemia to mortality in trauma patients," *J. Trauma*, 56(5):1058-1062 (May 2004).
- Lamb RE, Goldstein BJ. Modulating an Oxidative-Inflammatory Cascade: Potential New Treatment Strategy for Improving Glucose Metabolism, Insulin Resistance, and Vascular Function. *Int. J. Clin. Pract.* Jul. 2008(epub May 16, 2008); 62(7): 1087-1095.
- Lamharzi N, Renard CB, Kramer F, Pennathur S, Heinecke JW, Chait A, Bomfeldt KE. Hyperlipidemia in concert with hyperglycemia stimulates the proliferation of macrophages in atherosclerotic lesions: potential role of glucose-oxidized LDL. *Diabetes*. Dec. 2004;53(12):3217-3225.
- Landmesser U, Dikalov S, Price SR, McCann L, Fukai T, Holland SM, Mitch WE, Harrison DG. Oxidation of tetrahydrobiopterin leads to uncoupling of endothelial cell nitric oxide synthase in hypertension. *J. Clin. Invest.*, Apr. 2003;111:1201-1209.
- LaRosa JC. Understanding risk in hypercholesterolemia. *Clin Cardiol* 26(Suppl 1):3-6, Jan. 2003.
- Larsen, L.N., et al., "Heneicosapentaenoate (21:5n-3): Its incorporation into lipids and its effects on arachidonic acid and eicosanoid Synthesis." *Lipids* 32:707-714 (Jul. 1997).
- Laufs et al., "Upregulation of endothelial nitric oxide synthase by hmg coa reductase inhibitors," *Circulation* (Mar. 31, 1998) 97:1129-1135.
- Law TK, Yan AT, Gupta A, et al. Primary prevention of cardiovascular disease: global cardiovascular risk assessment and management in clinical practice. *Eur Heart J Qual Care Clin Outcomes*. 1(1):31-36 (publication date Jul. 2, 2015; epublication date Jul. 1, 2015).
- Law, M.R., et al., "Quantifying effect of statins on low density lipoprotein cholesterol, ischaemic heart disease, and stroke: systematic review and meta-analysis." *Br Med J.*, 326:1423-1427 (Jun. 28, 2003).
- Lawson et al., "Human absorption of fish oil fatty acids as triacylglycerols, free acids or ethyl esters," *Biochemical and Biophysical Research Communications* 152(1):328-335 (Apr. 15, 1988).
- Leaf A, Albert CM, Josephson M, et al. For the Fatty Acid Antiarrhythmia Trial Investigators. Prevention of Fatal Arrhythmias in High-Risk Subjects by Fish Oil n-3 Fatty Acid Intake. *Circ.* Nov. 1, 2005;112:2762-2768.
- Leaf A, Kang JX. Prevention of cardiac sudden death by N-3 fatty acids: a review of the evidence. *J Intern Med* 240:5-12, Jul. 1996.
- Leaf, "Hypertriglyceridemia: A Guide to Assessment and Treatment," *Hospital Physician* 17-23 (Sep. 2008).
- Leaf, A., "Historical overview of n3 fatty acids and coronary heart disease." *Am J Clin Nutr* 87:1978S-80S. (Jun. 2008).
- Lee and G.Y.H. Lip, "The Role of Omega-3 Fatty Acids in the Secondary Prevention of Cardiovascular Disease", *Q J Med*, 96:465-480, (Jul. 2003).
- Lee C, Sigari F, Segrado T, Horkko S, Hama S, Subbiah PV, Miwa M, Navab M, Witztum JL, Reaven PD. All ApoB-containing lipoproteins induce monocyte chemotaxis and adhesion when minimally modified. Modulation of lipoprotein bioactivity by platelet-activating factor acetylhydrolase. *Arterioscler. Thromb. Vasc. Biol.*, Jun. 1999; 19(6): 1437-1446.
- Lee, J.H., et al., "Omega-3 fatty acids for cardioprotection." *Mayo Clin Proc.*, 83(3):324-332 (Mar. 2008).
- Leigh-Firbank et al., "Eicosapentaenoic acid and docosahexanoic acid from fish oils: differential associations with lipid responses," *Br. J. Nutr.* 87:435-445 (May 2002).
- Lemaitre, R.N., et al., "n-3 Polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study." *Am J Clin Nutr* 77:319-25 (Feb. 2003).
- Leonard, Brian E., "Neurological Aspects", *Fundamentals of Psychopharmacology*, 186-187, (1997).
- Leucht, S., et al., *Schizophrenia Research*, vol. 35, "Efficacy and extrapyramidal side-effects of the new antipsychotics olanzapine,

US 10,568,861 B1

Page 18

(56)

References Cited

OTHER PUBLICATIONS

quetiapine, risperidone, and sertindole compared to conventional antipsychotics and placebo. A meta-analysis of randomized controlled trials", pp. 51-68, (Jan. 4, 1999).

Levey A, et al. A New Equation to Estimate Glomerular Filtration Rate. *Ann Intern Med.* 150:604-612; May 5, 2009.

Li, D. et al., "Effect of dietary α -linolenic acid on thrombotic risk factors in vegetarian men." *Am J Clin Nutr* 69:872-82 (May 1999).

Li, H., et al., "EPA and DHA reduce LPS-induced inflammation responses in HK-2 cells: Evidence for a PPAR- γ -dependent mechanism." *Kidney Int* 1. 67:867-74 (Mar. 2005).

Libby P. Triglycerides on the rise: should we swap seats on the seesaw? *Eur Heart J.* 36(13):774-776 (publication date Apr. 1, 2015; epublication date Dec. 29, 2014).

Libby, "Inflammation and atherosclerosis," *Nature* (Dec. 2002) 420(6917):868-874.

Lichtman et al., "Depression and Coronary Heart Disease, Recommendations for Screening, Referral and Treatment," *AHA Science Advisory, Circulation* 118:1768-1775 (Sep. 29, 2008).

Lien, E.L., "Toxicology and safety of DHA." *Prostaglandins Leukot Essent Fatty Acids.*, 81:125-132 (2009).

Lin, Pao-Yen, M.D., et al., "A Meta-Analytic Review of Double-Blind, Placebo-Controlled Trials of Antidepressant Efficacy of Omega-3 Fatty Acids", *Psychiatry*, 1056-1061 (Jul. 2007).

Lin, Y., et al., "Differential effects of eicosapentaenoic acid on glycerolipid and apolipoprotein B metabolism in primary human hepatocytes compared to HepG2 cells and primary rat hepatocytes." *Biochimica et Biophysica Acta* 1256:88-96 (Apr. 28, 1995).

Lindsey, S., et al., "Low density lipoprotein from humans supplemented with n-3 fatty acids depresses both LDL receptor activity and LDL mRNA abundance in HepG2 cells." *J Lipid Res.*, 33:647-658 (Mar. 1992).

Lipitor [package insert]. New York, NY: Parke-Davis (2012). (22 pages).

Lipitor [product information] Dublin, Ireland: Pfizer Inc. (2007). (18 pages).

Liu et al., "Effects of stable fish oil and simvastatin on plasma lipoproteins in patients with hyperlipidemia," *Nutrition Res.*, vol. 23, pp. 1027-1034 (Aug. 2003).

Liu X, et al., Stearoyl CoA Desaturase 1: Role in Cellular Inflammation and Stress, *Adv. Nutr.* Jan. 2011 (Jan. 10, 2011); 2:15-22.

Lohmussaar, E., et al., "ALOX5AP Gene and the PDE4D Gene in a Central European Population of Stroke Patients." *Stroke*, 36:731-736 (Apr. 2005) (epub Feb. 24, 2005).

Lovaza (omega-3-acid ethyl esters) Capsules, Prescribing information, GlaxoSmithKline (Nov. 2008).(9 pages).

Lovaza [package insert]. Research Triangle Park, NC: GlaxoSmithKline (2012). (14 pages).

Lovaza Side Effects, web archived webpage, archived from Drugs.com website on (Jul. 31, 2010), Retrieved from URL <<https://web.archive.org/web/20100731021902/https://www.drugs.com/sfx/lovaza-side-effects.html>> (4 pages)(Jul. 2010).

Lovaza TM (omega-3-acid ethyl esters) Capsules, Aug. 2007 (Aug. 1, 2007)m oaget 1-2, XP055589332.

Lovaza United States Prescribing Information, GlaxoSmithKline. Research Triangle Park, USA, May 2014.

Lovaza, (omega-3-acid ethyl esters) Capsules, Prescribing information Smith Kline Beechum (Jul. 2009).(17 pages).

Lovaza, GlaxoSmithKline, Lovaza Prescribing Information, Jun. 2008 [retrieved from the internet Jun. 6, 2012 <https://web.archive.org/web/20090206170311/http://us.gsk.com/products/assets/us_lovaza.pdf>]; Table 3, p. 1, section entitled 'Description'; p. 3, section entitled 'Very High Triglycerides: Monotherapy'; p. 4 section entitled 'Indications and Usage' and 'Information for Patients.' (12 pages).

Lovaza® (omega-3-acid ethyl esters) Capsules, Prescribing Information, GlaxoSmithKline, (Dec. 2010)(12 pages).

Lovaza®, Physicians' Desk Reference 2699-2701 (62d ed., 2008). (4 pages).

Lovegrove et al., "Moderate fish-oil supplementation reverses low-platelet, long chain n-3 polyunsaturated fatty acid status and reduced plasma triacylglycerol concentrations in British Indo-Asians," *Am. J. Clin. Nutr.*, 79:974-982 (Jun. 2004).

Lu, G., et al., "Omega-3 fatty acids alter lipoprotein subfraction distributions and the in vitro conversion of very low density lipoproteins to low-density lipoproteins." *J Nutr Biochem.*, 10:151-158 (Mar. 1999).

Lucas, M., et al., "Ethyl-eicosapentaenoic acid for the treatment of psychological distress and depressive symptoms in middle-aged women: a double-blind, placebo-controlled, randomized clinical trial." *Am J Clin Nutr* 89:641-51 (Feb. 2009)(epub Dec. 30, 2008).

Luria, MH, "Effect of low-dose niacin on high-density lipoprotein cholesterol and total cholesterol/high density lipoprotein cholesterol ratio", *Arch. Int. Med.*, 148:2493-2495, (Nov. 1998).

Lvovich V, Scheeline A. Amperometric sensors for simultaneous superoxide and hydrogen peroxide detection. *Anal. Chem.* Feb. 1, 1997;69:454-462.

Madhavi et al., "Effect of n-6 and n-3 fatty acids on the survival of vincristine sensitive and resistant human cervical carcinoma cells in vitro", *Cancer Letters*, vol. 84. No. 1, pp. 31-41 (Aug. 29, 1994).

Madsen, L., et al., "Eicosapentaenoic and Docosahexaenoic Acid Affect Mitochondrial and Peroxisomal Fatty Acid Oxidation in Relation to Substrate Preference." *Lipids* 34:951-963 (Sep. 1999).

Mak IT, Weglicki WB. Antioxidant properties of calcium channel blocking drugs. *Methods Enzymol.* 1994;234:620-630.

Maki et al., "Effects of Adding Prescription Omega-3 Acid Ethyl Esters to Simvastatin (20 mg/day) on Lipids and Lipoprotein Particles in Men and Women with Mixed Dyslipidemia," *Am. J. Cardiol.*, 102:429-433 (Aug. 15, 2008)(Epub May 22, 2008).

Maki, K.C., et al., "Baseline lipoprotein lipids and low-density lipoprotein cholesterol response to prescription omega-3 acid ethyl ester added to simvastatin therapy." *Am J Cardiol.*, 105:1409-1412 (May 15, 2010)(epub Mar. 30, 2010).

Maki, PhD, et al., "Lipid Responses to a Dietary Docosahexaenoic Acid Supplement in Men and Women with Below Average Levels of High Density Lipoprotein Cholesterol." *Journal of the American College of Nutrition*, vol. 24, No. 3, 189-199 (Jun. 2005).

Malinowski et al., "Elevation of Low-Density Lipoprotein Cholesterol Concentration with Over-the-Counter Fish Oil Supplementation." *Annals of Pharmacotherapy* 41:1296-1300 (Jul./Aug. 2007).

Malinski T, Taha Z. Nitric oxide release from a single cell measured in situ by a porphyrinic-based microsensor. *Nature.* Aug. 20, 1992;358:676-678.

Mallat, Z., et al., "Apoptosis in the vasculature: mechanisms and functional importance." *British Journal of Pharmacology* 130:947-962 (Jul. 2000).

Mallat, Z., et al., "Protective role of interleukin-10 in atherosclerosis." *Circ. Res.* 85:e17-e24 (Oct. 15, 1999).

Manninen V, Tenkanen L, Koskinen P, et al. Joint effects of serum triglyceride and LDL cholesterol and HDL cholesterol concentrations on coronary heart disease risk in the Helsinki Heart Study. Implications for treatment. *Circulation* 85:37-45, Jan. 1992.

Marangell, Lauren B., M.D., et al., "A Double-Blind, Placebo-Controlled Study of the Omega-3 Fatty Acid Docosahexaenoic Acid in the Treatment of Major Depression", *Am. J. Psychiatry*, 160(5):996-998, (May 2003).

Marchioli R, Barzi F, Bomba E, et al, GISSI-Prevenzione Investigators. Early protection against sudden death by n-3 polyunsaturated fatty acids after myocardial infarction: time-course analysis of the results of the Gruppo Italiano per lo Studio della Sopravvivenza nell'Infarto Miocardico (GISSI)-Prevenzione. *Circulation.* 105(16):1897-1903, Apr. 23, 2002.

Marckmann, P., "Fishing for heart protection." *Am J Clin Nutr*, 78:1-2 (Jul. 2003).

Marcoux et al., "Plasma remnant-like particle lipid and apolipoprotein levels in normolipidemic and hyperlipidemic subjects," *Atherosclerosis*, vol. 139, pp. 161-171 (Jul. 1998).

Marder, "An Approach to Treatment Resistance in Schizophrenia," *British Journ. Psychiatry*, 37:19-22 (1999).

Margolis, Simeon "What is Hyperlipidemia?" (<http://www.healthcommunities.com/highcholesterol/whatishyperlipidemia.shtml>, accessed Oct. 20, 2015, published Aug. 25, 2011)(4 pages).

US 10,568,861 B1

Page 19

(56)

References Cited

OTHER PUBLICATIONS

- Martin SS, Blaha MJ, Elshazly MB, et al. Comparison of a novel method vs the Friedewald equation for estimating low-density lipoprotein cholesterol levels from the standard lipid profile. *JAMA*. Nov. 20, 2013;310:2061-8.
- Martinez-Gonzalez J, Raposo B, Rodriguez C, Badimon L. 3-hydroxy-3-methylglutaryl coenzyme a reductase inhibition prevents endothelial no synthase downregulation by atherogenic levels of native ldl: Balance between transcriptional and posttranscriptional regulation. *Arterioscler. Thromb. Vasc. Biol.* May 2001;21:804-809.
- Martinez-Gonzalez, Jose et al., "Estatinas y acidos grasos omega-3. Disminucion de la mortalidad cardiovascular dependiente e independiente de la reduccion de la colesterolemia," (2006) *Rev Esp Cardiol Suppl.*, 6(D):20D-30D [with English abstract].
- Martin-Jadraque, R. et al., Effectiveness of low dose crystalline nicotinic acid in men with low density lipoprotein cholesterol levels. *Arch. Int. Med.* 156: 1081-1088. (May 27, 1996).
- Martz, "Moving Upstream in Huntington's," *Science-Business eXchange*, 2 pgs., Oct. 2008.
- Mason et al., "Comparative lipid antioxidant effects of omega-3 fatty acids in combination with HMG-CoA reductase inhibitors," *Journ. Clin. Lipidology* (May/Jun. 2011) 5(3):201.
- Mason et al., "Direct evidence for cholesterol crystalline domains in biological membranes: role in human pathobiology," *Biochimica et Biophysica Acta* 198-207 (Mar. 10, 2003).
- Mason et al., "Eicosapentaenoic Acid (EPA) inhibits the formation of membrane cholesterol crystalline domains by a potent antioxidant mechanism," *Journ. Clin. Lipid.*, 7(3): 272-273 (May/Jun. 2013) [Abstract only].
- Mason et al., "Eicosapentaenoic acid inhibits glucose-induced membrane cholesterol crystalline domain formation through a potent antioxidant mechanism," *Biochim. Biophys. Acta.*, 1848(2):502-9, (Feb. 2015).
- Mason et al., "Eicosapentaenoic Acid Inhibits Oxidation of ApoB-containing Lipoprotein Particles of Different Size In Vitro When Administered Alone or in Combination With Atorvastatin Active Metabolite Compared With Other Triglyceride-lowering Agents," *J. Cardiovasc. Pharmacol.*, 68(1):33-40 (Jul. 2016).
- Mason et al., "Eicosapentaenoic acid reduces membrane fluidity, inhibits cholesterol domain formation, and normalizes bilayer width in atherosclerotic-like model membranes," *Biochim. Biophys. Acta.*, 1858(12):3131-3140 (Dec. 2016).
- Mason RP, Gonye GE, Chester DW, Herbert LG. Partitioning and location of Bay K 8644, 1,4-dihydropyridine calcium channel agonist, in model and biological membranes. *Biophys. J. Apr. 1989;55(4):769-778*.
- Mason RP, Jacob RF, Kubant R, Walter MF, Bellamine A, Jacoby A, Mizuno Y, Malinski T. Effect of enhanced glycemic control with saxagliptin on endothelial nitric oxide release and CD40 levels in obese rats. *J. Atheroscler. Thromb. Epub Jun. 13, 2011;18:774-783*.
- Mason RP, Jacob RF. Membrane microdomains and vascular biology: Emerging role in atherogenesis. *Circulation*. May 6, 2003; 107:2270-2273.
- Mason RP, Kalinowski L, Jacob RF, Jacoby AM, Malinski T. Nebivolol reduces nitroxidative stress and restores nitric oxide bioavailability in endothelium of black americans. *Circulation*. Dec. 13, 2005 (epub Dec. 5, 2005); 112:3795-3801.
- Mason RP, Kubant R, Heeba G, Jacob RF, Day CA, Medlin YS, Funovics P, Malinski T. Synergistic effect of amlodipine and atorvastatin in reversing ldl-induced endothelial dysfunction. *Pharm. Res.* Aug. 2008 (epub 2007 Dec. 2018); 25:1798-1806.
- Mason RP, Walter MF, Day CA, Jacob RF. Active metabolite of atorvastatin inhibits membrane cholesterol domain formation by an antioxidant mechanism. *J. Biol. Chem.* Apr. 7, 2006 (epub Feb. 7, 2006); 281(14):9337-9345.
- Mason RP, Walter MF, Day CA, Jacob RF. Intermolecular differences for HMG-CoA reductase inhibitors contribute to distinct pharmacologic and pleiotropic actions. *Am. J. Cardiol.* Sep. 5, 2005;96(5A):11F-23F.
- Mason RP, Walter MF, Jacob RF. Effects of hmg-coa reductase inhibitors on endothelial function: Role of microdomains and oxidative stress. *Circulation*. Jun. 1, 2004;109:II34-II41.
- Mason RP, Walter MF, Mason PE. Effect of oxidative stress on membrane structure: Small angle x-ray diffraction analysis. *Free Radic. Biol. Med.* 1997;23(3):419-425.
- Mason RP. Molecular basis of differences among statins and a comparison with antioxidant vitamins. *Am. J. Cardiol.* Dec. 4, 2006 (epub Oct. 10, 2006); 98:34P-41P.
- Mataki et al., "Effect of Eicosapentaenoic Acid in Combination with HMG-CoA Reductase Inhibitor on Lipid Metabolism," *Int. Med. J.* 5(1):35-36 (Mar. 1998).
- Mater, M.K., et al., "Arachidonic acid inhibits lipogenic gene expression in 3T3-L1 adipocytes through a prostanoid pathway." *J. Lipid Res.* 39:1327-1334 (Jul. 1998).
- Matsumoto, M., et al., "Orally administered eicosapentaenoic acid reduces and stabilizes atherosclerotic lesions in ApoE-deficient mice." *Atherosclerosis*, 197(2):524-533 (Apr. 2008/epub Sep. 4, 2007).
- Matsuzaki et al., "Incremental Effects of Eicosapentaenoic Acid on Cardiovascular Events in Statin-Treated Patients with Coronary Artery Disease," *Circ. J.* 73:1283-1290 (Jul. 2009/epub May 8, 2009).
- Matsuzawa, Y., et al., "Effect of Long-Term Administration of Ethyl Icosapentate (MND-21) in Hyperlipaemic Patients," *J. Clin Therapeutic & Medicines*, 7: 1801-16 (1991).
- Mattson MP. Modification of ion homeostasis by lipid peroxidation: roles in neuronal degeneration and adaptive plasticity. *Trends Neurosci.* Feb. 1998;21(2):53-57.
- Mayatepek, E., et al., The Lancet, vol. 352, Leukotriene C4-synthesis deficiency: a new inborn error of metabolism linked to a fatal developmental syndrome, pp. 1514-1517 (Nov. 7, 1998).
- Mayo Clinic at <http://www.mayoclinic.org/diseases-conditions/high-blood-cholesterol/in-depth/cholesterol> (2014)(5 pages).
- Mayo Clinic, Diabetes Diagnosis and Treatment, 1998, <http://www.mayoclinic.org/diseases-conditions/diabetes/diagnosis-treatment/drc-20371451> (1998-2018).
- McElroy, S.L., et al., "Clozapine in the Treatment of Psychotic Mood Disorders, Schizoaffective Disorder, and Schizophrenia", *Journal of Clinical Psychiatry*, vol. 52, No. 10, pp. 411-414 (Oct. 1991).
- McIntyre M, Hamilton CA, Rees DD, Reid JL, Dominiczak AF. Sex differences in the abundance of endothelial nitric oxide in a model of genetic hypertension. *Hypertension*. Dec. 1997;30:1517-1524.
- McKenney et al., "Prescription omega-3 fatty acids for the treatment of hypertriglyceridemia," *Am. J. Health Syst. Pharm.*, 64(6):595-605 (Mar. 15, 2007).
- McKenney et al., CMRO, "Comparison of the efficacy of rosuvastatin versus atorvastatin, simvastatin and pravastatin in achieving lipid goals: results from the STELLAR trial", 689-98 (2003).
- McKenney, J., "Niacin for dyslipidemia: considerations in product selection", *Am. J. Health Syst. Pharm.*, 60:995-1005, (May 15, 2003).
- McKenney, J.M. et al. Study of the pharmacokinetic interaction between simvastatin and prescription omega-3-acid ethyl esters. *J. Clin. Pharmacol.* 46, 785-791 (Jul. 2006).
- McKenney, James et al., "Role of prescription omega-3 fatty acids in the treatment of Hypertriglyceridemia," *Pharmacotherapy, LNKD—Pubmed: 17461707*, vol. 27, No. 5, pp. 715-728 (May 2007).
- McKeone et al., "Alterations in serum phosphatidylcholine fatty acyl species by eicosapentaenoic and docosahexaenoic ethyl esters in patients with severe hypertriglyceridemia." *J. Lipid Res.* 38:429-436 (Mar. 1997).
- McMurichie, E.J., et al., "Incorporation and effects of dietary eicosapentaenoate (20 : 5(n-3)) on plasma and erythrocyte lipids of the marmoset following dietary supplementation with differing levels of linoleic acid." *Biochimica et Biophysica Acta*, 1045:164-173 (Jul. 16, 1990).
- McNamara JR, et al., Remnant-like particle (RLP) Cholesterol is an independent cardiovascular disease risk factor in women: results from the Framingham Heart Study, *Atherosclerosis*, vol. 154(1), pp. 229-236 (Jan. 2001).

US 10,568,861 B1

Page 20

(56)

References Cited

OTHER PUBLICATIONS

- MedlinePlus. "Coronary heart disease," Available at: <https://medlineplus.gov/ency/article/007115.htm> (review date Jul. 14, 2015)(accessed Sep. 2, 2016)(5 pages).
- Menuet, R. et al., "Importance and management of dyslipidemia in the metabolic syndrome," *American Journal of the Medical Sciences* Dec. 2005 US, vol. 33, No. 6, pp. 295-302 (2005).
- Merched, A.J., et al., "Atherosclerosis: evidence for impairment of resolution of vascular inflammation governed by specific lipid mediators," *FASEB J.* 22:3595-3606 (Oct. 2008/epub Jun. 17, 2008).
- Merkel et al., "Antisense Oligonucleotide Directed to Human Apolipoprotein B-100 Reduces Lipoprotein(a) Levels and Oxidized Phospholipids on Human Apolipoprotein B-100 Particles in Lipoprotein(a) Transgenic Mice," *Circulation*, vol. 118, pp. 743-753 (epub Jul. 28, 2008).
- Mesa, M., "Effects of oils rich in Eicosapentaenoic and docosahexaenoic acids on the oxidizability and thrombogenicity of low-density lipoprotein," *Artherosclerosis* 175, pp. 333-343 (Aug. 2004).
- Metcalf, R.G., et al., "Effect of dietary n-3 polyunsaturated fatty acids on the inducibility of ventricular tachycardia in patients with ischemic cardiomyopathy," *Am J Cardiol* 101:758-761 (Mar. 15, 2008/epub Jan. 14, 2008).
- Metcalf, R.G., et al., "Effects of fish-oil supplementation on myocardial fatty acids in humans," *Am J Clin Nutr* 85:1222-28 (May 2007).
- Meyer et al., "Comparison of Seal Oil to Tuna Oil on Plasma Lipid Levels and Blood Pressure in Hypertiglyceridaemic Subjects," *Lipids*, 44:827-835 (Sep. 2009).
- Meyer, et al., "Dose-Dependent Effects of Docosahexaenoic Acid Supplementation on Blood Lipids in Statin-Treated Hyperlipidaemic Subjects," *Lipids*, 42:109-115 (Mar. 2007/epub Feb. 8, 2007).
- Meyers, et al., "Nicotinic acid induces secretion of prostaglandin D2 in human macrophages: An in vitro model of the niacin-flush", *Atherosclerosis*, 192:253-258, (Jun. 2007/ epub Sep. 1, 2006).
- Micheletta F, Natoli S, Misuraca M, Sbarigia E, Diczfalussy U, Iuliano L. Vitamin E supplementation in patients with carotid atherosclerosis: Reversal of altered oxidative stress in plasma but not in plaque. *Arterioscler. Thromb. Vasc. Biol.* Jan. 2004 (epub Dec. 16, 2006); 24:136-140.
- Michos et al., "Niacin and Statin Combination Therapy for Atherosclerosis Regression and Prevention of Cardiovascular Disease Events," *Journ. Amer. Coll. Cardiol.*, vol. 59, No. 23:2058-2064 (Jun. 5, 2012)(epub Apr. 18, 2012).
- Mii, S., et al., "Perioperative use of eicosapentaenoic acid and patency of infrainguinal vein bypass: a retrospective chart review," *Curr Ther Res Clin Exp.* 68:161-174 (May 2007).
- Miles, et al., "Effect of orlistat in overweight and obese patients with type 2 diabetes treated with metformin," *Diabetes Care*, 25(7):1123-1128 (2002).
- Miller AK, DiCicco RA, Freed MI. The effect of ranitidine on the pharmacokinetics of rosiglitazone in healthy adult male volunteers. *Clin. Ther.* Jul. 2002;24:1062-1071.
- Miller AK, Inglis AM, Culkun KT, Jorkasky DK, Freed MI. The effect of acarbose on the pharmacokinetics of rosiglitazone. *Eur. J. Clin. Pharmacol.* May 2001;57:105-109.
- Miller M, Cannon CP, Murphy SA, et al. Impact of triglyceride levels beyond low-density lipoprotein cholesterol after acute coronary syndrome in the PROVE IT-TIMI 22 trial. *J Am Coll Cardiol* 51:724-730, Feb. 19, 2008.
- Miller M, Stone NJ, Ballantyne C, et al. Triglycerides and cardiovascular disease: a scientific statement from the American Heart Association. *Circulation*. May 24, 2011 (epub Apr. 18, 2011); 123:2292-2333.
- Miller M. Current perspectives on the management of hypertriglyceridemia. *Am Heart J* 140:232-40, 2000.
- Miller, M., et al., "Impact of lowering triglycerides on raising HDL-C in hypertriglyceridemic and non-hypertriglyceridemic subjects," *International Journal of Cardiology* 119:192-195 (Jul. 10, 2007)(epub Oct. 18, 2006).
- Minihane, A.M., et al., "ApoE polymorphism and fish oil supplementation in subjects with an atherogenic lipoprotein phenotype." *Arterioscler. Thromb. Vasc. Biol.* 20:1990-1997 (Aug. 2000).
- Mishra, A., et al., "Oxidized omega-3 fatty acids inhibit NF-kB activation via a PPAR α -Dependent Pathway," *Arterioscler Thromb Vasc Biol.* 24:1621-1627 (Sep. 2004)(epub Jul. 1, 2004).
- Missouri DUReport, Statin Therapy (Oct./Nov. 2003) Drug Use Review Newsletter 8(6):1-9.
- Mita, T. et al., Eicosapentaenoic acid reduces the progression of carotid intima-media thickness in patients with type 2 diabetes, *Atherosclerosis* 191:162-167 (Mar. 2007)(epub Apr. 17, 2006).
- Mizota M, Katsuki Y, Mizuguchi K, Endo S, Miyata H, Kojima M, Kanehiro H et al. "Pharmacological studies of eicosapentaenoic acid ethylester (EPA E) on high cholesterol diet-fed rabbits," *Nippon Yakurigaku Zasshi*, 91:255-66 (Apr. 1988) (with English abstract).
- Mizota M, Katsuki Y, Mizuguchi K, Endo S, Miyata H, Kojima M, Kanehiro H et al. "The effects of eicosapentaenoic acid ethylester (EPA E) on arterial thrombosis in rabbits and vascular lesions in rats," *Nippon Yakurigaku Zasshi*, 91:81-9 (Feb. 1988)(with English abstract).
- Mizuguchi K, Yano T, Kojima M, Tanaka Y, Ishibashi M, Masada A, Sato M et al. "Hypolipidemic effect of ethyl all-cis-5,8,11,14,17-eicosapentaenoate (EPA-E) in rats," *Jpn J Pharmacol.*, 59(3):307-12 (Jul. 1992).
- Mizuguchi, K., et al., "Ethyl all-cis-5,8,11,14,17-icosapentaenoate modifies the biochemical properties of rat very low-density lipoprotein." *European Journal of Pharmacology*, 231:221-227 (Apr. 28, 1993).
- Mizuguchi, K., et al., "Mechanism of the lipid-lowering effect of ethyl all-cis-5,8,11,14,17-icosapentaenoate." *European Journal of Pharmacology*, 231:121-127 (Jan. 1993).
- Mochida Press Release, Pharmaceutical Col., Ltd.: Conclusion of Distributorship Agreement Concerning Switch-OTC Drug for Hyperlipidemia Treatment, Epadel, (Apr. 30, 2009)(1 page).
- Mochida, Announcement, "Mochida Announces Completion of "JELIS" Major Clinical Trial for Epadel," Mar. 22, 2005 (2 pages).
- Mochida's Epadel Reduces Risk of Stroke Recurrence—New Results of JELIS Major Clinical Trial, JCNNetwork Newswire Nov. 13, 2006 (2 pages).
- Mora, S., et al., "LDL particle subclasses, LDL particle size, and carotid atherosclerosis in the Multi-Ethnic Study of Atherosclerosis (MESA)." *Atherosclerosis*;192:211-217 (May 2007).
- Mori et al., "Differential Effects of Eicosapentaenoic Acid and Docosahexaenoic Acid on Vascular Reactivity of the Forearm Microcirculation in Hyperlipidemic, Overweight Men," *Circulation*, 102:1264-1269 (Sep. 12, 2000).
- Mori TA, Woodman RJ. "The independent effects of eicosapentaenoic acid and docosahexaenoic acid on cardiovascular risk factors in humans," *Curr Opin Clin Nutr Metab Care* 2006; 9:95-104 (Mar. 2006).
- Mori TA. Omega-3 fatty acids and blood pressure. *Cell Mol Biol.* Feb. 25, 2010;56(1):83-92.
- Mori, et al., "Purified Eicosapentaenoic and docosahexaenoic acids have differential effects on serum lipids and lipoproteins, LDL particle size, glucose, and insulin in mildly hyperlipidemic men," *Am J Clin Nutr* 71:1085-1094 (May 2000).
- Mori, T. et al., Effect of Eicosapentaenoic acid and docosahexaenoic acid on oxidative stress and inflammatory markers in treated-hypertensive type 2 diabetic subjects, *Free Radical Biology & Medicine*, vol. 35, No. 7, pp. 772-781 (Oct. 1, 2003).
- Mori, Trevor A., et al., "Docosahexaenoic Acid but Not Eicosapentaenoic Acid Lowers Ambulatory Blood Pressure and Heart Rate in Humans", *Hypertension*, 34(2):253-60 (Aug. 1999).
- Morita, I., et al., "Effects of purified eicosapentaenoic acid on arachidonic acid metabolism in cultured murine aortic smooth muscle cells, vessel walls and platelets." *Lipids* 18:42-490 (Jan. 1983).
- Morris M, Sacks F, Rosner B. Does fish oil lower blood pressure? A meta-analysis of controlled trials. *Circ.*, Aug. 1993;88:523-533.
- Morrow, JD, "Release of markedly increased quantities of prostaglandin D2 in vivo in humans following the administration of nicotinic acid", *Prostaglandins*, 38:263-274, (Aug. 1989).

US 10,568,861 B1

Page 21

(56)

References Cited

OTHER PUBLICATIONS

- Morton, R.E., "Specificity of lipid transfer protein for molecular species of cholesteryl ester." *J Lipid Res.*, 27:523-529 (May 1986).
- Mosher I.R. et al., "Nicotinic Acid Side Effects and Toxicity: A review," *Am J Psychiat.*, 126: 1290-1296 (Mar. 1970).
- Mostad et al., "Effects of Marine N-3 Fatty Acid Supplementation on Lipoprotein Subclasses Measured by Nuclear Magnetic Resonance in Subjects with Type II Diabetes," *European Journ. Clin. Nutr.*, 62(3):419-429 (Mar. 2008/epub Feb. 27, 2007).
- Mostad, I.L. et al., "Effects of n-3 fatty acids in subjects with type 2 diabetes: reduction of insulin sensitivity and time-dependent alteration from carbohydrate to fat oxidation." *Am J Clin Nutr* 84:540-50 (Sep. 2006).
- Mozaffarian D, Benjamin EJ, Go AS, Arnett DK, Blaha MJ, et al.; on behalf of the American Heart Association Statistics Committee and Stroke Statistics Subcommittee. Heart disease and stroke statistics—2016 update: a report from the American Heart Association [published online ahead of print Dec. 16, 2015]. *Circulation*. doi: 10.1161/CIR.0000000000000350.
- Mozaffarian D, Geelen A, Brouwer I, Geleijnse J, Zock P, Katan M. Effect of Fish Oil on Heart Rate in Humans A Meta-Analysis of Randomized Controlled Trials. *Circ.Sep.* 27, 2005/ epub Sep. 19, 2005; 112:1945-1952.
- Mozaffarian D, Marchioli R, Macchia A, et al. Fish Oil and Postoperative Atrial Fibrillation the Omega-3 Fatty Acids for Prevention of Post-operative Atrial Fibrillation (OPERA) Randomized Trial. *JAMA*. Nov. 21, 2012;308(19):2001-11.
- Mozaffarian D, Psaty B, Rimm E, Lemaitre R, Burke G, Lyles M, Lefkowitz D, Siscovick D. Fish Intake and Risk of Incident Atrial Fibrillation. *Circ.*, Jul. 27, 2004/epub Jul. 19, 2004; 110:368-373.
- Mozaffarian et al., "Omega-3 fatty acids and cardiovascular disease: effects on risk factors, molecular pathways and clinical events," *J. Am. Coll. Cardiol.* (Nov. 8, 2011) 58(2):2047-2067.
- Mozaffarian, "JELIS, fish oil, and cardiac events," *www.thelancet.com* vol. 369, pp. 1062-1063 (Mar. 31, 2007).
- Mozaffarian, D., "Fish and n-3 fatty acids for the prevention of fatal coronary heart disease and sudden cardiac death." *Am J Clin Nutr*, 87:1991S-6S (Jun. 2008).
- Mozaffarian, D., et al., "Dietary fish and ω -3 fatty acid consumption and heart rate variability in US adults." *Circulation*, 117:1130-1137 (Mar. 4, 2008/ epub Feb. 19, 2008).
- Murck et al., "Ethyl-EPA in Huntington disease—Potentially relevant mechanism of action," *Brain Research Bulletin*, 72:159-164 (2007) (available online Nov. 15, 2006).
- Murphy SA, Cannon CP, Blazing MA, et al. Reduction in total cardiovascular events with ezetimibe/simvastatin post-acute coronary syndrome. *J Am Coll Cardiol.* 67(4):353-361 (publication date Feb. 2, 2016; epublication date Jan. 25, 2016).
- Naba, H., et al., "Improving effect of ethyl eicosapentanoate on statin-induced rhabdomyolysis in Eisai hyperbilirubinemic rats." *Biochemical and Biophysical Research Communications*, 340:215-220 (Feb. 2006/epub Dec. 9, 2005).
- Nagakawa et al., Effect of [EPA] on the Platelet Aggregation and Composition of Fatty Acid in Man: A Double Blind Study, *Atherosclerosis* 47(1):71-75 (Apr. 1983).
- Naik H, Wu JT, Palmer R, McLean L. The effects of febuxostat on the pharmacokinetic parameters of rosiglitazone, a CYP2C8 substrate. *Br. J. Clin. Pharmacol.* Jan. 13, 2012;74:327-335.
- Nakamura et al., Remnant lipoproteinemia is a risk factor for endothelial vasomotor dysfunction and coronary artery disease in metabolic syndrome, *Atherosclerosis*, vol. 181(2), pp. 321-327 (Aug. 2005/epub Feb. 16, 2005).
- Nakamura, et al., "Effects of Eicosapentaenoic Acids on Remnant-like Particles, Cholesterol Concentrations and Plasma Fatty Acid Composition in Patients with Diabetes Mellitus." *in vivo* 12: 311-314 (May/Jun. 1998).
- Nakamura, H., et al., "Evaluation of ethyl icosapentate in the treatment of hypercholesterolemia in kidney transplant recipients." *Transplantation Proceedings*, 30:3047-3048 (Nov. 1998).
- Nakamura, N. et al., "Joint effects of HMG-CoA reductase inhibitors and eicosapentaenoic acids on serum lipid profile and plasma fatty acid concentrations in patients with hyperlipidemia," *International Journal of Clinical and Laboratory Research*, Springer, Berlin, DE LNKD-DOI: 10.1007/S005990050057, vol. 29, No. 1, pp. 22-25 (1999).
- Nambi V, Bhatt DL. Primary prevention of atherosclerosis: Time to take a selfie? *J Am Coll Cardiol* 2017;70(24):2992-4 (publication date Dec. 19, 2017; epublication date Dec. 11, 2017).
- Nambi, V. et al., "Combination therapy with statins and omega-3 fatty acids." *Am J Cardiol* 98:34i-38i (Aug. 21, 2006/epub May 30, 2006).
- Nasa, et al., "Long-Term Supplementation With Eicosapentaenoic Acid Salvages Cardiomyocytes From Hypoxia/Reoxygenation-Induced Injury in Rats Fed With Fish-Oil-Deprived Diet," *Jpn. J. Pharmacol.* 77, 137-146 (Jun. 1998).
- National Kidney Foundation, "Glomerular Filtration Rate (GFR)," Jan. 30, 2017 (Jan. 30, 2017), retrieved on Jul. 30, 2018 from <https://web.archive.org/web/20170130183218/https://www.kidney.org/atoz/content/gfr>; entire document, especially p. 1 paragraph 1 and p. 3, paragraph 2.
- National Kidney Foundation, "The Heart and Kidney Connection," Apr. 17, 2017 (Apr. 17, 2017), retrieved on Jul. 30, 2018 from <https://web.archive.org/web/2017041700416/https://www.kidney.org/atoz/content/heart-and-kidney-connection>; entire document, especially p. 2, paragraph 1.
- Natsuno et al., "Clinical Effects of Eicosapentaenoic Acid on Type-2 Diabetes Effects on Serum Lipids, Pulse Wave Speed, and Ankle-Brachial Blood Pressure Index," *Diagnosis and Treatment* 93(12):133-137 (2005)(16 pages).
- Nattel, S. et al., "Atrial remodeling and atrial fibrillation: Mechanisms and implications." *Circ Arrhythmia Electrophysiol*, 1:62-73 (Apr. 2008).
- Needleman P, Raz A, Minkes MS, Ferrendelli JA, Sprecher H. Triene prostaglandins: prostacyclin and thromboxane biosynthesis and unique biological properties. *Proc Natl Acad Sci USA*. Feb. 1979;76:944-948.
- Negre-Salvayre, A., et al., "Advanced lipid peroxidation end products in oxidative damage to proteins. Potential role in diseases and therapeutic prospects for the inhibitors." *British Journal of Pharmacology* 153:6-20 (Jan. 2008/epub Jul. 23, 2007).
- Nelson et al. "Icosapent Ethyl for Treatment of Elevated Triglyceride Levels," *Annals of Pharmacotherapy*, 47(11):1517-1523 (Nov. 2013/epub Nov. 5, 2013).
- Nelson JR, Wani O, May HT, Budoff M. Potential benefits of eicosapentaenoic acid on atherosclerotic plaques. *Vascul Pharmacol.* 91:1-9 (publication date Apr. 2017; epublication date Mar. 2, 2017).
- Nelson, G.J., et al., "The Effect of Dietary Docosahexaenoic Acid on Plasma Lipoproteins and Tissue Fatty Acid Composition in Humans", *Lipids*, 32(11):1137-1146, (Nov. 1997).
- Nemets, Boris, M.D., "Addition of Omega-3 Fatty Acid to Maintenance Medication Treatment for Recurrent Unipolar Depressive Disorder", *Am. J. Psychiatry*, 159(3):477-479, (Mar. 2002).
- Nemoto et al., "Ethyl-eicosapentaenoic Acid Reduces Liver Lipids and Lowers Plasma Levels of Lipids in Mice Fed a High-Fat Diet, *in vivo*," 23:685-690 (Sep./Oct. 2009).
- Nenseter, MS et al., "Effect of dietary supplementation with n-3 polyunsaturated fatty acids on physical properties and metabolism of low density lipoprotein in humans," *Arterioscler. Thromb. Vasc. Biol.*, 12:369-379 (Mar. 1992).
- Nestel, et al., "The n-3 fatty acids eicosapentaenoic acid and docosahexaenoic acid increase systemic arterial compliance in humans," *Am J Clin Nutr.*, 76:326-30 (Aug. 2002).
- Nestel, P.J., "Effects of N-3 fatty acids on lipid metabolism." *Ann Rev Nutr.*, 10:149-167 (1990).
- Nichols GA, Philip S, Reynolds K, Granowitz CB, Fazio S. Increased cardiovascular risk in hypertriglyceridemic patients with statin-controlled LDL cholesterol. *J Clin Endocrinol Metab* 103(8):3019-27 (publication date Aug. 1, 2018; epublication date May 29, 2018).
- Nichols GA, Philip S, Reynolds K, Granowitz CB, Fazio S. Increased residual cardiovascular risk in patients with diabetes and high vs.

US 10,568,861 B1

Page 22

(56)

References Cited

OTHER PUBLICATIONS

normal triglycerides despite statin-controlled LDL Cholesterol. *Diabetes Obes Metab* (publication date Sep. 17, 2018; epublication date Sep. 17, 2018).

Niemi M, Backman JT, Grantors M, Laitila J, Neuvonen M, Neuvonen PJ. Gemfibrozil considerably increases the plasma concentrations of rosiglitazone. *Diabetologia*. Oct. 2003/ Jul. 29, 2003; 46: 1319-1323.

Niemi M, Backman JT, Neuvonen PJ. Effects of trimethoprim and rifampin on the pharmacokinetics of the cytochrome P450 2C8 substrate rosiglitazone. *Clin. Pharmacol. Ther.*, Sep. 2004;76:239-249.

Nigon F, Lesnik P, Rouis M, Chapman MJ. Discrete subspecies of human low density lipoproteins are heterogeneous in their interaction with the cellular LDL receptor. *J. Lipid Res.*, Nov. 1991;32(11):1741-1753.

Nippon Rinsho, *Metabolic Syndrome 2nd Edition—Basics and New Clinical Findings*, Jan. 20, 2011, Special Issue 1 (vol. 992), pp. 503-506 (with English translation).

Nishikawa M. et al., "Effects of Eicosapentaenoic acid (EPA) on prostacyclin production in diabetics. GC/MS analysis of PG12 and PG13 levels" *Methods Find Exp Clin Pharmacol*. 19(6):429-33 (Jul./Aug. 1997).

Nobukata, H., et al., "Age-related changes in coagulation, fibrinolysis, and platelet aggregation in male WBN/Kob rats." *Thrombosis Research* 98: 507-516 (Jun. 15, 2000).

Nobukata, H., et al., "Long-term administration of highly purified eicosapentaenoic acid ethyl ester improves the dysfunction of vascular endothelial and smooth muscle cells in male WBN/Kob rats." *Metabolism*, 49(12): 1588-1591 (Dec. 2000).

Nobukata, H., et al., "Long-term administration of highly purified eicosapentaenoic acid ethyl ester prevents diabetes and abnormalities of blood coagulation in male WBN/Kob rats." *Metabolism*, 49(12): 912-919 (Jul. 2000).

Noguchi et al., "Chemoprevention of DMBA-induced mammary carcinogenesis in rats by low-dose EPA and DHA." *Br. J. Cancer* 75(3): 348-353 (1997).

Nomura et al., "The effects of pitavastatin, eicosapentaenoic acid and combined therapy on platelet-derived microparticles and adiponectin in hyperlipidemic, diabetic patients." *Platelets*, 20(1):16-22 (Feb. 2009).

Nomura S, Shouzu A, Omoto S, et al. Effects of eicosapentaenoic acid on endothelial cell-derived microparticles, angiopoietins and adiponectin in patients with type 2 diabetes. *J Atheroscler Thromb.*, Apr. 2009;16:83-90.

Nourooz-Zadeh, J., et al., "Urinary 8-epi-PGF2 α and its endogenous β -oxidation products (2,3-dinor and 2,3-dinor-5,6-dihydro) as biomarkers of total body oxidative stress." *Biochemical and Biophysical Research Communications* 330:731-736 (May 13, 2005).

Nozaki S. et al., "Effects of purified Eicosapentaenoic acid ethyl ester on plasma lipoproteins in primary hypercholesterolemia" *Int J Vitam Nutr Res*. 62(3):256-260 (1992).

Obata, et al., "Eicosapentaenoic acid inhibits prostaglandin D2 generation by inhibiting cyclo-oxygenase in cultured human mast cells". *Clin. & Experimental Allergy*, 29:1129-1135, (Aug. 1999).

O'Donnell, C.J., et al., "Leukocyte telomere length and carotid artery intimal medial thickness—the Framingham heart study." *Arteriosclerosis, Thrombosis, and Vascular Biology*.28:1165-1171 (Jun. 2008/epub Apr. 3, 2008).

Oemar BS, Tschudi MR, Godoy N, Brovkovich V, Malinski T, Luscher TF. Reduced endothelial nitric oxide synthase expression and production in human atherosclerosis. *Circulation.*, Jun. 30, 1998;97:2494-2498.

Oh, Robert C et al., *Management of Hypertriglyceridemia*, American Family Physician, LNKD-PUBMED: 17508532, vol. 75, No. 9, pp. 1365-1371 (May 1, 2007).

Ohara Y, Peterson TE, Harrison DG. Hypercholesterolemia increases endothelial superoxide anion production. *J. Clin. Invest.* Jun. 1983;91:2546-2551.

Ohashi, *Journal of Clinical and Experimental Medicine*, Feb. 14, 2009, vol. 228, No. 7, pp. 795-805 (with English translation).

Okuda, Y. et al., Eicosapentaenoic acid enhances nitric oxide production by cultured human endothelial cells. *Biochem. Biophys. Res. Commun.* 232: 487-491 (Mar. 17, 1997).

Okuda, Y., et al., "Long-term effects of eicosapentaenoic acid on diabetic peripheral neuropathy and serum lipids in patients with type II diabetes mellitus." *Journal of Diabetes and Its Complications* 10:280-287 (Sep./Oct. 1996).

Okumura, T., et al., "Eicosapentaenoic acid improves endothelial function in hypertriglyceridemic subjects despite increased lipid oxidizability." *Am J Med Sci* 324(5):247-253 (Nov. 2002).

Oliv, E.H., et al., "Biosynthesis of prostaglandins from 17(18)epoxy-eicosatetraenoic acid, a cytochrome P-450 metabolite of eicosapentaenoic acid." *Biochimica et Biophysica Acta*, 1126, 261-268 (Jun. 26, 1992).

Olofsson et al., "Apolipoprotein B: a clinically important apolipoprotein which assembles atherogenic lipoproteins and promotes the development of atherosclerosis" *Journal of Internal Medicine*, 258: 395-410 (Nov. 2005).

Omacor Summary of Product Characteristics, Pronova BioPharma Norge AS. Lysaker, Norway, Mar. 2015.

Omacor® Prescribing Information (Omega-3-acid ethyl esters, capsules) (2004). (9 pages).

Omacor®, Physicians' Desk Reference 2735 (60th ed. 2006)(3 pages).

Ona, V.O., et al., *Nature*, vol. 399, Inhibition of caspase-1 slows disease progression in a mouse model of Huntington's disease, pp. 263-267 (May 20, 1999).

Ooi EM, "Apolipoprotein C-III: Understanding an emerging cardiovascular risk factor", *Clin.Sci. (London)*, vol. 114, pp. 611-624 (May 2008).

Opalinska et al., "Increasing Level of Prostate-Specific Antigen and Prostate Cancer Risk Factors Among 193 Men Examined in Screening Procedure," *Ann. Univ. Curie Sklowska Med.*, 58(2):57-63 (Abstract Only)(2003)(2 pages).

Origin Trial Investigators (The). n-3 fatty acids and cardiovascular outcomes in patients with dysglycemia. *N Engl J Med* Jul. 6, 2012/epub Jun. 11, 2012; 367:309-318.

O'Riordan, "DHA and EPA have differential effects on LDL-cholesterol," May 24, 2011 [online][Retrieved on Aug. 21, 2015] Retrieved from website: <http://www.medscape.com/viewarticle/743305> (2 pages).

Osher et al., "Omega-3 Eicosapentaenoic Acid in Bipolar Depression: Report of a Small Open-Label Study," *J. Clin. Psych.* 66:726-729 (Jun. 2005).

Otvos et al., "Clinical Implications of Discordance Between LDL Cholesterol and LDL Particle Number," *J. Clin. Lipidol*, 5(2):105-113 (Mar.-Apr. 2011)(available online Mar. 1, 2011).

Ou Z, Ou J, Ackerman AW, Oldham KT, Pritchard KA, Jr. L-4f, an apolipoprotein a-1 mimetic, restores nitric oxide and superoxide anion balance in low-density lipoprotein-treated endothelial cells. *Circulation*. Mar. 25, 2003;107:1520-1524.

Ozaki M, Kawashima S, Yamashita T, Hirase T, Namiki M, Inoue N, Hirata K, Yasui H, Sakurai H, Yoshida Y, Masada M, Yokoyama M. Overexpression of endothelial nitric oxide synthase accelerates atherosclerotic lesion formation in apoe-deficient mice. *J. Clin. Invest.* Aug. 2002; 110:331-340.

Ozawa, Akio, Nakamura E, Jinbo H, Fujita T, Hirai A, Terano T, Hamazaki T et al., "Determination of higher fatty acids in various lipid fractions of human plasma, platelets, and erythrocyte membrane using thin layer chromatography and gas chromatography," *Bunseki Kagaku*, 32:174-8 (1982) (with English abstract).

Padgett et al., "Phylogenetic and immunological definition of four lipoylated proteins from *Novosphingobium aromaticivorans*, implications for primary biliary cirrhosis," *Journ. Autoimmunity* 24:209-219 (May 2005).

Park JH, Park DI, Kim HJ, et al. Metabolic syndrome is associated with erosive esophagitis. *World J. Gastroenterol.* Sep. 14, 2008 (35): 5442-7.

Park JY, Kim KA, Kang MH, Kim SL, Shin JG. Effect of rifampin on the pharmacokinetics of rosiglitazone in healthy subjects. *Clin. Pharmacol. Ther.*, Mar. 2004;75:157-162.

US 10,568,861 B1

Page 23

(56)

References Cited

OTHER PUBLICATIONS

- Park, Y., et al., "Omega-3 fatty acid supplementation accelerates chylomicron triglyceride clearance." *J. Lipid Res.* 44:455-463 (Mar. 2003).
- Pase M, Grima N, Sarris J. Do long-chain n-3 fatty acids reduce arterial stiffness? A meta-analysis of randomized controlled trials. *Br J Nutr.* Oct. 2011; 106:974-980.
- Patel et al., "Rosiglitazone monotherapy improves glycaemic control in patients with type 2 diabetes: a twelve-week, randomized, placebo-controlled study," *Diabetes, Obesity and Metabolism*, vol. 1, pp. 165-172 (May 1999).
- Paton, CM, Ntambi, JM., Biochemical and physiological function of stearoyl-CoA desaturase, *AM. J. Physiol. Endocrinol. Metab.* Jul. 2009/epub Dec. 9, 2008; 297:E28-E37.
- PCT/GB00/00164 International Search Report dated Oct. 20, 2000 (8 pages).
- PCT/US2011/062247 International Search Report and Written Opinion dated Jun. 14, 2012 (12 pages).
- PCT/US2013/020526 International Search Report dated Mar. 29, 2013 (2 pages).
- PCT/US2013/048241 International Search Report dated Dec. 13, 2013 (3 pages).
- PCT/US2013/048516 International Search Report dated Dec. 20, 2013 (3 pages).
- PCT/US2013/048559 International Search Report dated Dec. 13, 2013 (3 pages).
- PCT/US2013/068647 International Search Report and Written Opinion dated May 13, 2014 (18 pages).
- PCT/US2014/019454 International Search Report and Written Opinion dated Jun. 3, 2014 (12 pages).
- Pedersen RS, Damkier P, Brosen K. The effects of human CYP2C8 genotype and fluvoxamine on the pharmacokinetics of rosiglitazone in healthy subjects. *Br. J. Clin. Pharmacol.* Dec. 2006/epub Jul. 12, 2006; 62:682-689.
- Pedersen, T., et al., "Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease: the Scandinavian Simvastatin Survival Study (4S)", *The Lancet*, No. 19, vol. 344, 8934, p. 1383-1389 (Nov. 19, 1994).
- Peet et al., "A Dose-Ranging Study of the Effects of Ethyl-Eicosapentaenoate in Patients with Ongoing Depression Despite Apparently Adequate Treatment with Standard Drugs", *Arch. Gen. Psychiatry*, 59:913-919, (Oct. 2002).
- Peet, M., et al., Phospholipid Spectrum Disorder in Psychiatry pp. 1-19, (1999).
- Pejic et al., "Hypertriglyceridemia," *Journ. Amer. Board Fam. Med.*, vol. 19(3):310-316 (May/Jun. 2006).
- Pennathur S, Heinecke JW. Mechanisms for oxidative stress in diabetic cardiovascular disease. *Antioxid. Redox Signal.* Jul. 2007;9(7):955-969.
- Piccini, Monica, et al., *Genomics*, vol. 47, "FACL4, a new gene encoding long-chain acyl-CoA synthetase 4, is deleted in a family with Alport syndrome, elliptocytosis, and mental retardation," pp. 350-358 (Feb. 1998).
- Piche, "Tumor Necrosis Factor-Alpha, and Fibrinogen to Abdominal Adipose Tissue, Blood Pressure, and Cholesterol and Triglyceride Levels in Healthy Postmenopausal Women", *American Journal of Cardiology*, 2005, 96(1), 92-97.
- Pike, NB, "Flushing out the role of GPR109A (HM74V) in the clinical efficacy of nicotinic acid", *J. Clin. Invest.* 115:3400-3403, (Dec. 2005).
- PLUSEPA® Product brochure "Super Critically" Different from Other Omega-3 Fish Oil Supplements for Depression and ADHD, by Minami Nutrition (Apr. 2009, pp. 1-6).
- Pollin TI, Damcott CM, Shen H, et al. A null mutation in human APOC3 confers a favorable plasma lipid profile and apparent cardioprotection. *Science*. Dec. 12, 2008;322(5908):1702-1705.
- Pownall, H.J., et al., "Correlation of serum triglyceride and its reduction by ω -3 fatty acids with lipid transfer activity and the neutral lipid compositions of high-density and low-density lipoproteins." *Atherosclerosis* 143:285-297 (Apr. 1999).
- Press Release: Amarin Corporation Says Huntington's Disease Drug Failed in Trials, <http://www.fiercebiotech.com/node/6607/print> (Apr. 24, 2007) (Printed on Aug. 22, 2008)(2 pages).
- Pritchard KA, Ackerman AW, Ou J, Curtis M, Smalley DM, Fontana JT, Stemerman MB, Sessa WC. Native low-density lipoprotein induces endothelial nitric oxide synthase dysfunction: Role of heat shock protein 90 and caveolin-1. *Free Radic. Biol. Med.* Jul. 2002;33:52-62.
- Pritchard KA, Jr., Groszek L, Smalley DM, Sessa WC, Wu M, Villalon P, Wolin MS, Stemerman MB. Native low-density lipoprotein increases endothelial cell nitric oxide synthase generation of superoxide anion. *Circ. Res.* Sep. 1995;77:510-518.
- Puri et al., "Reduction in Cerebral Atrophy Associated with Ethyl-eicosapentaenoic Acid Treatment in Patients with Huntington's Disease," *Journ. Int'l. Med. Research*, 36:896-905 (Oct. 1, 2008).
- Puri, B., et al., "Eicosapentaenoic Acid in Treatment-Resistant Depression Associated with Symptom Remission, Structural Brain Changes and Reduced Neuronal Phospholipid Turnover," *Int J Clinical Practice*, 55:560-563 (Oct. 2001).
- Puri, B., et al., *Archives of General Psychiatry*, No. 55, "Sustained remission of positive and negative symptoms of schizophrenia following treatment with eicosapentaenoic acid," pp. 188-189, (Feb. 1998).
- Puri, B.K., et al., "Ethyl-EPA in Huntington Disease: A Double-Blind, Randomized, Placebo-Controlled Trial", *Neurology*, 65:286-292, (Jul. 26, 2005).
- Qi, K., et al., "Omega-3 fatty acid containing diets decrease plasma triglyceride concentrations in mice by reducing endogenous triglyceride synthesis and enhancing the blood clearance of triglyceride-rich particles." *Clinical Nutrition* 27(8):424-430 (Jun. 2008/epub Mar. 24, 2008).
- Rader, *Lipid Disorders*, in Eric J. Topol (ed.) *Textbook of Cardiovascular Medicine* pp. 55-75 (2007).
- Rahimy M, Hallen B, Narang P. Effect of tolterodine on the anticoagulant actions and pharmacokinetics of single-dose warfarin in healthy volunteers. *Arzneimittelforschung* 2002 52 (12): 890-5.
- Raitt, M.H., et al., "Fish oil supplementation and risk of ventricular tachycardia and ventricular fibrillation in patients with implantable defibrillators—a randomized controlled trial." *JAMA*. 293(23):2884-2891 (Jun. 15, 2005).
- Rambjor, Gro S., et al., "Eicosapentaenoic Acid is Primarily Responsible for Hypotriglyceridemic Effect of Fish Oil in Humans", *Fatty Acids and Lipids from Cell Biology to Human Disease: Proceedings of the 2nd international Congress of the ISSFAL (International Society for the Study of Fatty Acids and Lipids, AOCS Press*, 31:S-45-S-49, (Mar. 1, 1996).
- Randomised trial of cholesterol lowering in 4444 patients with coronary heart disease. The Scandinavian Simvastatin Survival Study, *Lancet*. 344: 1383-1389 (1994).
- Rao MN, Mullangi R, Katneni K, et al. Lack of effect of sucralfate on the absorption and pharmacokinetics of rosiglitazone. *J. Clin. Pharmacol.* Jun. 2002;42:670-675.
- Rauch B, Rudolf R, Schneider S, et al. OMEGA, a randomized, placebo-controlled trial to test the effect of highly purified omega-3 fatty acids on top of modern guideline-adjusted therapy after myocardial infarction. *Circulation*. Nov. 23, 2010 (epub Nov. 8, 2010); 122:2152-2159.
- Rees DD, Palmer RM, Moncada S. The role of endothelium-derived nitric oxide in the regulation of blood pressure. *Proc. Natl. Acad. Sci. USA*. May 1989;86:3375-3378.
- Reich, "Formulation and physical properties of soft capsules," *Pharmaceutical capsules*. (2004) Chapter 11:201-212.
- Reiffel, J.A., et al., "Antiarrhythmic effects of omega-3 fatty acids." *Am J Cardiol* 98:50i-60i (Aug. 21, 2006/epub May 26, 2006).
- Reiner Z, Catapano AL, De BG, et al. ESC/EAS Guidelines for the management of dyslipidaemias: the Task Force for the management of dyslipidaemias of the European Society of Cardiology (ESC) and the European Atherosclerosis Society (EAS). *Eur. Heart J. Jul.* 2011(epub Jun. 28, 2011); 32:1769-1818.

US 10,568,861 B1

Page 24

(56)

References Cited

OTHER PUBLICATIONS

- Richter, Werner O., "Hypertriglyceridämie: Ein klinischer Leitfaden," Wissenschaftliche Verlagsgesellschaft mbH Stuttgart, front page to p. V, pp. 2 to 55, 64 to 85, 90 to 97 (2008) (with English Summary).
- Ridker PM, Everett BM, Thuren T, et al. Antiinflammatory Therapy with canakinumab for atherosclerotic disease. *N Engl J Med* 377(12):1119-31 (publication date Sep. 21, 2017; epublication date Aug 27, 2017).
- Ridker, "C-Reactive Protein : A Simple Test to Help Predict Risk of Heart Attack and Stroke", *Circulation: Journal of the American Heart Association*, Sep. 23, 2003, 108, e81-e85.
- Riediger, N.D., et al., "A systemic review of the roles of n-3 fatty acids in health and disease." *J Am Diet Assoc*. 109:668-679. (Apr. 2009).
- Rifai, "High-Sensitivity C-Reactive Protein: A Novel and Promising Marker of Coronary Heart Disease", *Clinical Chemistry*, Mar. 2001, 47(3), 403-411.
- Risé, P., et al., "Effects of simvastatin on the metabolism of polyunsaturated fatty acids and on glycerolipid, cholesterol, and de novo lipid synthesis in THP-1 cells." *J. Lipid Res*. 38:1299-1307 (Jul. 1997).
- Risk and Prevention Study Collaborative Group, Roncaglioni MC, Tombesi M, et al. n-3 fatty acids in patients with multiple cardiovascular risk factors. *N Engl J Med*, May 9, 2013;368(19):1800-8.
- Rissanen et al., "Fish Oil-Derived Fatty Acids, Docosahexaenoic Acid and Docosapentaenoic Acid, and the Risk of Acute Coronary Events The Kuopio Ischaemic Heart Disease Risk Factor Study," *Circulation*. (Nov. 28, 2000)(102):2677-2679 doi:10.1161/01.CIR.102.22.2677.
- Rizzo M, Bemeis K. Low-density lipoprotein size and cardiovascular risk assessment. *Q. J. Med*. Jan. 2006; 99(1): 1-14.
- Roach, P.D., et al., "The effects of dietary fish oil on hepatic high density and low density lipoprotein receptor activities in the rat." *FEBS Lett*, 222: 159-162 (Sep. 28, 1987).
- Robinson, J.G., et al., "Meta-analysis of the relationship between non-high-density lipoprotein cholesterol reduction and coronary heart risk." *J Am Coll Cardiol*, 53: 316-322 (Jan. 27, 2009).
- Roche, H.M., et al., "Effect of long-chain n-3 polyunsaturated fatty acids on fasting and postprandial triacylglycerol metabolism." *Am J Clin Nutr* 71:232S-7S (Jan. 2000).
- Roche, H.M., et al., "Long-chain n-3 polyunsaturated fatty acids and triacylglycerol metabolism in the postprandial state." *Lipids* 34: S259-S265 (1999).
- Rodriguez, Y., et al., "Long-chain ω 6 polyunsaturated fatty acids in erythrocyte phospholipids are associated with insulin resistance in non-obese type 2 diabetics." *Clinica Chimica Acta* 354:195-199 (Apr. 2005).
- Roe MT, Armstrong PW, Fox KAA, et al; TRILOGY ACS Investigators. Prasugrel versus clopidogrel for acute coronary syndromes without revascularization. *N Engl J Med*. 367(14):1297-1309 (publication date Oct. 4, 2012; (epublication Aug. 25, 2012).
- Rogers, P. J., "No effect of n-3 long-chain polyunsaturated fatty acid (EPA and DHA) supplementation on depressed mood and cognitive function: a randomised controlled trial" *British Journal of Nutrition*, 99:421-431, (Feb. 2008/epub Oct. 24, 2007).
- Rost KL, Roots I. Nonlinear kinetics after high-dose omeprazole caused by saturation of genetically variable CYP2C19. *Hepatology* Jun. 23, 1996 (6): 1491-7.
- Rubins, HB, et al., "Distribution of lipids in 8,500 men with coronary artery disease: Department of Veterans Affairs HDL Intervention Trial Study Group," *Am. J. Cardiol*, 75:1196-1201, (Jun. 15, 1995).
- Rubins, HB, et al., "Gemfibrozil for the secondary prevention of coronary heart disease in men with low levels of high-density lipoprotein cholesterol: Veterans Affairs HDL-C Intervention Trial Study Group", *N. Eng. J. Med.*, 341:410-418, (Aug. 5, 1999).
- Ruiz-Narváez, E.A., et al., "Abdominal obesity and hyperglycemia mask the effect of a common APOC3 haplotype on the risk of myocardial infarction." *Am J Clin Nutr* 87:1932-8 (Jun. 2008).
- Ruocco MJ, Shipley GG. Interaction of cholesterol with galactocerebroside and galactocerebroside phosphatidylcholine bilayer membranes. *Biophys. J. Dec.* 1984; 46:695-707.
- Rupp, "Omega-3-Fettsäuren in der Sekundärprävention nach Myokardinfarkt," *Clin. Res. Cardiol.*, vol. 95:Suppl. 6, Vi/12-V1-16 (2006)(with English summary).
- Rustan, A.C., et al., "Eicosapentaenoic acid inhibits cholesterol esterification in cultured parenchymal cells and isolated microsomes from rat liver." *J. Bio. Chem.* 263(17):8126-32 (Jun. 15, 1988).
- Rustan, A.C., et al., "Eicosapentaenoic acid reduces hepatic synthesis and secretion of triacylglycerol by decreasing the activity of acyl-coenzyme A:1,2-diacylglycerol acyltransferase." *J. Lipid Res*. 29:1417-1426 (Nov. 1988).
- Rustan, A.C., et al., "Postprandial decrease in plasma unesterified fatty acids during n-3 fatty acid feeding is not caused by accumulation of fatty acids in adipose tissue." *Biochimica et Biophysica Acta* 1390:245-25 (Feb. 23, 1998).
- Ryan, A.M., et al., "Enteral nutrition enriched with eicosapentaenoic acid (EPA) preserves lean body mass following esophageal cancer surgery: results of a double-blinded randomized controlled trial." *Ann Surg* 249:355-363 (Mar. 2009).
- Ryan, A.S., et al., "Clinical overview of algal-docosahexaenoic acid: effects on triglyceride levels and other cardiovascular risk factors." *Am J Ther.* 16:183-192 (Mar./Apr. 2009).
- Sacks, Frank M., "The apolipoprotein story," *Atherosclerosis Supplements*, 23-27 (Aug. 2006/epub Jul. 5, 2006).
- Saito et al., "Effects of EPA on coronary artery disease in hypercholesterolemic patients with multiple risk factors: Sub-analysis of primary prevention cases from the Japan EPA Lipid Intervention Study (JELIS)," *Atherosclerosis*, 200:135-140 (Sep. 2008/epub Jun. 19, 2008).
- Saito et al., "Results of Clinical Usage of Improved Formulation (MND-21S) Epadel Capsule 300 with Respect to Hyperlipidemia," 26(12) *Jpn. Pharmacol. Ther.* 2047-62 (1998) (with English abstract).
- Saito, J., et al., "Mechanisms of enhanced production of PGI₂ in cultured rat vascular smooth muscle cells enriched with eicosapentaenoic acid." *Atherosclerosis* 131: 219-228 (Jun. 1997).
- Sampath H, Ntambi JM., Role of stearoyl-CoA desaturase in human metabolic disease, *Future Lipidol*. 2008;3:2,163-73.
- Sampath H, Ntambi JM., The Role of stearoyl-CoA desaturase in obesity, insulin resistance, and inflammation, *Ann. NY. Acad. Sci.* Dec. 2011; 1243:4 7-53.
- Samuels, Martin A., M. D., et al., "Huntington's Disease", *Office Practice of Neurology*, (122):654-655, (1996).
- Sanders, A. Hinds and C.C. Pereira, "Influence of n-3 fatty acids on blood lipids in normal subjects" *Journal of Internal Medicine*. 225:99-104,(1989).
- Sanders, et al., "Influence of an algal triacylglycerol containing docosahexaenoic acid (22:6n-3) and docosapentaenoic acid (22:5n-6) on cardiovascular risk factors in healthy men and women," *British Journal of Nutrition*, 95, 525-531 (Mar. 2006).
- Sanders, T.A., et al., "Effect of varying the ratio of n-6 to n-3 fatty acids by increasing the dietary intake of α -linolenic acid, eicosapentaenoic and docosahexaenoic acid, or both on fibrinogen and clotting factors VII and XII in persons aged 45-70 y: the OPTILIP Study." *Am J Clin Nutr* 84:513-22 (Sep. 2006).
- Sanders, T.A., et al., "Triglyceride-lowering effect of marine polyunsaturates in patients with hypertriglyceridemia." *Arterioscler. Thromb. Vasc. Biol.* 5:459-465 (Sep./Oct. 1985).
- Sarwar N, Danesh J, Eiriksdottir G, et al. Triglycerides and the risk of coronary heart disease: 10,158 incident cases among 262,525 participants in 29 Western prospective studies. *Circulation* 115:450-458, Jan. 30, 2007/epub Dec. 26, 2006.
- Sasaki J, Miwa T, Odawara M. Administration of highly purified eicosapentaenoic acid to stain-treated diabetic patients further improves vascular function. *Endocrine J.* Jan. 27, 2012; 59(4):297-304.
- Sasaki J, Yokoyama M, Matsuzaki M, et al. Relationship between coronary artery disease and non-HDL-C, and effect of highly purified EPA on the risk of coronary artery disease in hypercholesterolemic patients treated with statins: sub-analysis of the Japan EPA Lipid Intervention Study (JELIS). *J. Atheroscler. Thromb.* Dec. 17, 2012;19:194-204.

US 10,568,861 B1

Page 25

(56)

References Cited

OTHER PUBLICATIONS

- Sasaki, Y.F., et al., "Bio-anticlastogenic effects of unsaturated fatty acids included in fish oil—docosahexaenoic acid, docosapentaenoic acid, and eicosapentaenoic acid in cultured Chinese hamster cells." *Mutation Research*, 320: 9-22 (Jan. 1994).
- Sato et al., "General Pharmacological Studies on 5 8 11 14 17 Eicosapentaenoic Acid Ethyl Ester EPA-E", *Folia Pharmacol JPN*, 94 (1), 35-47, cited by other (Jul. 1989) (with English abstract).
- Sato, "Effects of Highly Purified Ethyl All-cis-5,8,11,14,17-icosapentaenoate (EPA-E) on Rabbit Platelets," *Biol. Pharm. Bull.*, 16(4):362-367 (Apr. 1993).
- Satoh et al., "Highly purified eicosapentaenoic acid reduces cardio-ankle vascular index in association with decreased serum amyloid A-LDL in metabolic syndrome," *Hypertension Research* (Nov. 2009/epub Sep. 18, 2009) (32):1004-1008.
- Satoh, N., et al., "Purified eicosapentaenoic acid reduces small dense LDL, remnant lipoprotein particles, and C-reactive protein in metabolic syndrome." *Diabetes Care*, 30(1): 144-146 (Jan. 2007).
- Satoh-Asahara N, Shimatsu A, Sasaki Y, Nakaoka H, Himeno A, Tochiya M, Kono S, Takaya T, Ono K, Wada H, Suganami T, Hasegawa K, Ogawa Y., "Highly purified eicosapentaenoic acid increases interleukin-10 levels of peripheral blood monocytes in obese patients with dyslipidemia." *Diabetes Care*. Dec. 2012/epub Aug. 21, 2012; 35(12):2631-2639.
- Schaefer, E.J., et al., "Effects of eicosapentaenoic acid, docosahexaenoic acid, and olive oil on cardiovascular disease risk factors [abstract 20007]." *Circulation*, 122:A20007 (2010) (Abstract only).
- Schechtman, G. & Hiatt, J., "Drug therapy for hypercholesterolemia in patients with cardiovascular disease: factors limiting achievement of lipid goals", *Am. J. Med.*, 100:197-204, (Feb. 1996).
- Schechtman, G., et al., "Dietary fish oil decreases low-density-lipoprotein clearance in nonhuman primates." *Am J Clin Nutr.*, 64:215-221 (Aug. 1996).
- Schechtman, G., et al., "Heterogeneity of Low Density Lipoprotein Responses to Fish-Oil Supplementation in Hypertriglyceridemic Subjects." *Arterioscler. Thromb. Vasc. Biol.* 9:345-354 (May/Jun. 1989).
- Schmidt, E.B., et al., "Lipoprotein-associated phospholipase A2 concentrations in plasma are associated with the extent of coronary artery disease and correlate to adipose tissue levels of marine n-3 fatty acids." *Atherosclerosis* 196: 420-424 (Jan. 2008).
- Schmitz PG, McCloud LK, Reikes ST, et al. Prophylaxis of hemodialysis graft thrombosis with fish oil: double-blind, randomized, prospective trial. *J. Am. Soc. Nephrol.* Jan. 13, 2002 (1): 184-90.
- Schmitz, G., et al., "The opposing effects of n-3 and n-6 fatty acids." *Progress in Lipid Research*, 47:147-155 (Mar. 2008/epub Dec. 27, 2007).
- Schreiner et al., "Lipoprotein[a] as a Risk Factor for Preclinical Atherosclerosis," *13 Atherosclerosis, Thrombosis & Vascular Biology* 6: 826-833 (1993).
- Schürmann, D.J. A comparison of the two one-sided tests procedure and the power approach for assessing the equivalence of average bioavailability. *J. Pharmacokinet. Biopharm.* 15(6), 657-680 (Dec. 1987).
- Schunkert H, König IR, Kathiresan S, et al. Large-scale association analysis identifies 13 new susceptibility loci for coronary artery disease. *Nat Genet.* Mar. 6, 2011;43(4):333-8.
- Schwartz GG, Bessac L, Berdan LG, et al. Effect of alirocumab, a monoclonal antibody to PCSK9, on long-term cardiovascular outcomes following acute coronary syndromes: rationale and design of the ODYSSEY outcomes trial. *Am Heart J* 168(5):682-9 (publication date Nov. 2014, epublication date Aug 7, 2017).
- Schwarz, S., et al., "Lycopene inhibits disease progression in patients with benign prostate hyperplasia." *J. Nutr.* 138: 49-53 (Jan. 2008).
- Schwellenbach et al., "The Triglyceride-Lowering Effects of a Modest Dose of Docosahexaenoic Acid Alone Versus in Combination with Low Dose Eicosapentaenoic Acid in Patients with Coronary Artery Disease and Elevated Triglycerides." *J. Am. Coll. Nutr.* 25(6):480-485 (Dec. 2006).
- Segrest et al., Structure of Apolipoprotein B-100 in Low Density Lipoproteins, *J. Lipid Res.* 42(9):1346-1367 (Sep. 2001).
- Self-Medlin Y, Byun J, Jacob RF, Mizuno Y, Mason RP. Glucose promotes membrane cholesterol crystalline domain formation by lipid peroxidation. *Biochim. Biophys. Acta.* Jun. 2009/epub Apr. 17, 2009; 1788(6): 1398-1403.
- Serhan C, Chiang N, Van Dyke T. Resolving inflammation: dual anti-inflammatory and pro-resolution lipid mediators. *Nat Rev Immunol.* May 2008; 8:3449-361.
- Serhan, C.N., et al., "Resolvins: A family of bioactive products of omega-3 fatty acid transformation circuits initiated by aspirin treatment that counter proinflammation signals." *J. Exp. Med.* 196:1025-1037 (Oct. 21, 2002).
- Sevanian A, Ursini F. Lipid peroxidation in membranes and low-density lipoproteins: similarities and differences. *Free Radic. Biol. Med.*, Aug. 2000;29(3-4):306-311.
- Shah, S. et al., "Eicosapentaenoic Acid (EPA) as an Adjunct in the Treatment of Schizophrenia", *Schizophrenia Research*, vol. 29, No. 1/02 (1998).
- Shan, Z., et al., "A combination study of spin-trapping, LC/ESR and LC/MS on carbon-centred radicals formed from lipoxygenase-catalysed peroxidation of eicosapentaenoic acid." *Free Radical Research*, 43(1):13-27 (Jan. 2009).
- Shearer et al., "Red Blood Cell Fatty Acid Patterns and Acute Coronary Syndrome," *PLoS ONE* 4(5): e5444 (doi:10.1371/journal/pone.0005444)(May 6, 2009).
- Sherratt SCR, Mason RP. Eicosapentaenoic acid and docosahexaenoic acid have distinct membrane locations and lipid interactions as determined by X-ray diffraction. *Chem Phys Lipids* 212:73-9 (publication date May 2018, epublication date Jan. 31, 2018).
- Shimizu et al., "Effects of Highly Purified Eicosapentaenoic Acid on Erythrocyte Fatty Acid Composition and Leukocyte and Colonic Mucosa Leukotriene B4 Production in Children with Ulcerative Colitis," *J. Pediatr. Gastroenterol. Nutr.*, vol. 37, No. 5, pp. 581-585 (Nov. 2003).
- Shimizu, H., et al., "Long-term effect of eicosapentaenoic acid ethyl (EPA-E) on albuminuria of non-insulin dependent diabetic patients." *Diabetes Research and Clinical Practice* 28: 35-40 (Apr. 1995).
- Shimokawa H, Flavahan NA, Vanhoutte PM. Loss of endothelial pertussis toxin-sensitive G protein function in atherosclerotic porcine coronary arteries. *Circulation.* Feb. 1991;83:652-660.
- Shinozaki K. et al., "The long-term effect of Eicosapentaenoic acid on serum levels of lipoprotein (a) and lipids in patients with vascular disease" *J Atheroscler Thromb.* 2(2):207-9 (1996).
- Shishehbor MH, Brenna ML, Aviles RJ, Fu X, Penn MS, Sprecher DL, Hazen SL. Statins promote potent systemic antioxidant effects through specific inflammatory pathways. *Circulation.* Jul. 29, 2003;108(4):426-431.
- Sicherer et al., "Prevalence of seafood allergy in the United States determined by a random telephone survey," *J. Allergy Clin. Immunol.*, 114(1):159-165 (Jul. 2004).
- Sierra, S., et al., "Dietary eicosapentaenoic acid and docosahexaenoic acid equally incorporate as decosahexaenoic acid but differ in inflammatory effects." *Nutrition* 24: 245-254 (Mar. 2008).
- Silvers, Karen M., et al., "Randomised double-blind placebo-controlled trial of fish oil in the treatment of depression", *Prostaglandins, Leukotrienes and Essential Fatty Acids*, 72:211-218, (Mar. 2005).
- Simmons, C.M., et al., "Inclusion of 10% fish oil in mixed medium-chain triacylglycerol-long chain triacylglycerol emulsions increases plasma triacylglycerol clearance and induces rapid eicosapentaenoic acid (20:5n-3) incorporation into blood cell phospholipids." *Am J Clin Nutr* 88: 282-8 (Aug. 2008).
- Simon, Joel A., et al., "Serum Fatty Acids and the Risk of Coronary Heart Disease", *American Journal of Epidemiology*, 142(5):469-476, (Sep. 1, 1995).
- Simopoulos, The Importance of the Omega-6/Omega-3 Fatty Acid Ratio in Cardiovascular Disease and Other Chronic Diseases, *Exp. Biol. Med.*, 233:674-688 (Jun. 1, 2008)(available online Jun. 1, 2008).
- Simopoulos, "Omega-3 fatty acids in health and disease and in growth and development," *Am. J. Clin. Nutr.* 54:438-63 (Sep. 1991).

US 10,568,861 B1

Page 26

(56)

References Cited

OTHER PUBLICATIONS

- Singer, Peter, "Fluvastatin plus fish oil are more effective on cardiovascular risk factors than fluvastatin alone," Letter to the Editor, *Prostaglandins, Leukotrienes and Essential Fatty Acids*, vol. 72, pp. 379-380 (May 2005).
- Singh, R.B., et al., "Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival—4," *Cardiovascular Drugs and Therapy* 11:485-491 (Jul. 1997).
- Sirtori, C.R., et al., "One-year treatment with ethyl esters of n-3 fatty acids in patients with hypertriglyceridemia and glucose intolerance—Reduced triglyceridemia, total cholesterol and increased HDL-C," *Atherosclerosis* 137: 419-427 (Apr. 1998).
- Skinner JS, Cooper A, & Feder GS and on behalf of the Guideline Development Group. "Secondary prevention for patients following a myocardial infarction; summary of NICE guidance," *Heart*, 93:862-864 (Jul. 2007).
- Slides for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, (158 pages).
- Smith et al., Pharmacokinetics and Pharmacodynamics of Epoetin Delta in Two Studies in Health Volunteers and Two Studies in Patients with Chronic Kidney Disease, *Clinical Therapeutics*/vol. 29, pp. 1368-1380 (Jul. 2007).
- Sniderman A, Kwiterovich PO. Update on the detection and treatment of atherogenic low-density lipoproteins. *Curr. Opin. Endocrinol. Diabetes Obes.* Apr. 20, 2013;20:140-147.
- Sohma, R., et al., "Protective effect of n-3 polyunsaturated fatty acid on primary culture of rat hepatocytes without glycemic alterations," *Journal of Gastroenterology and Hepatology* 22: 1965-1970 (Nov. 2007).
- Spector, A.A., "Arachidonic acid cytochrome P450 epoxygenase pathway," *Journal of Lipid Research*, 50: S52-S56 (2009) (published online on Oct. 23, 2008.).
- Spector, A.A., et al., "Epoxyeicosatrienoic acids (EETs): metabolism and biochemical function," *Progress in Lipid Research* 43: 55-90 (Jan. 2004).
- Springer, T.A., "Traffic signals for lymphocyte recirculation and leukocyte emigration: The multistep paradigm," *Cell*, 76: 301-314 (Jan. 28, 1994).
- Squires, RW, et al., "Low-dose, time release nicotinic acid: effects in selected patients with low concentrations of high density lipoprotein cholesterol", *Mayo Clinic Proc.*, 67:855-860, (Sep. 1992).
- Srinivas, et al., "Controlled release of lysozyme from succinylated gelatin microspheres," *J. Biomater. Sci., Polymer Ed.*, vol. 12(2):137-148 (2001).
- Stalenhoef, A.F.H., et al., "The effect of concentrated n-3 fatty acids versus gemfibrozil on plasma lipoproteins, low density lipoprotein heterogeneity and oxidizability in patients with hypertriglyceridemia," *Atherosclerosis* 153: 129-138 (Nov. 2000).
- Stampfer MJ, Krauss RM, Ma J, et al. A prospective study of triglyceride level, lowdensity lipoprotein particle diameter, and risk of myocardial infarction. *JAMA*. Sep. 1996;276:882-888.
- Stancu et al., "Statins: Mechanism of Action and Effects," *Journal of Cellular and Molecular Medicine* (Oct.-Dec. 2001), 5(4), 378-387.
- Stark, K.D. & Holub, B.J., Differential eicosapentaenoic acid elevations and altered cardiovascular disease risk factor responses after supplementation with docosahexaenoic acid in postmenopausal women receiving and not receiving hormone replacement therapy, *Am. J. Clin. Nutr.*, vol. 79, pp. 765-773 (May 2004).
- Stark, K.D., "The percentage of n-3 highly unsaturated fatty acids in total HUFA as a biomarker for omega-3 fatty acid status in tissues," *Lipids* 43:45-53 (Jan. 2008/epub Nov. 6, 2007).
- Stark, K.D., et al., "Effect of a fish-oil concentrate on serum lipids in postmenopausal women receiving and not receiving hormone replacement therapy in a placebo-controlled, double-blind trial," *Am J Clin Nutr* 72:389-94 (Aug. 2000).
- Steg PG, Bhatt DL, Wilson PWF, et al; REACH Registry Investigators. One-year cardiovascular event rates in outpatients with atherothrombosis. *JAMA*. 297(11):1197-1206 (publication date May 21, 2007).
- Stein et al., "Effect of Statin Therapy on Remnant Lipoprotein Cholesterol Levels in Patients with Combined Hyperlipidemia," *Arteriosclerosis, Thrombosis and Vascular Biology*, vol. 21, pp. 2026-2031 (Dec. 1, 2001).
- Steinberg D, Witztum JL. Is the oxidative modification hypothesis relevant to human atherosclerosis? Do the antioxidant trials conducted to date refute the hypothesis? *Circulation*. Apr. 30, 2002;105:2107-2111.
- Steinberg D, Lewis A. Conner Memorial Lecture: Oxidative modification of LDL and atherogenesis. *Circulation*. Feb. 18, 1997;95(4):1062-1071.
- Stepp DW, Ou J, Ackerman AW, Welak S, Klick D, Pritchard KA, Jr. Native Idl and minimally oxidized Idl differentially regulate superoxide anion in vascular endothelium in situ. *Am. J. Physiol.* Aug. 2002;283:H750-H759.
- Sternbach "The Glasgow Coma Scale," *The Journal of Emergency Medicine*, 19(1):67-71 (Feb. 8, 2000).
- Stielow et al., "Novel Nox Inhibitor of oxLDL-Induced Reactive Oxygen Specis Formation in Human Endothelial Cells," *Biochem. Biophys. Res. Comm.*, 344:200-205 (May 26, 2006/epub Mar. 26, 2006).
- Stiles, FDA approves EPA-only omega-3 PUFA capsule for high TG, Jul. 26, 2012, <http://www.medscape.com/viewarticle/791268>, accessed Dec. 17, 2014 (1 page).
- Stitzel N, Stirrups K, Masca N, et al. Supplement to: Coding variation in ANGPTL4, LPL, and SVEP1 and the risk of coronary disease. *N Engl J Med*. DOI: 10.1056/NEJMoa1507652; Mar. 24, 2016/epub Mar. 2, 2016.
- Stojancevic et al., "The impact of farnesoid X receptor activation on intestinal permeability in inflammatory bowel disease," *Can. J Gastroenterol.* 26(9):631-637 (Sep. 2012).
- Stoll, Andrew L. et al., "Omega 3 Fatty Acids in Bipolar Disorder", *Arch. Gen. Psychiatry*, 56:407-412, (May 1999).
- Stone NJ, Robinson J, Lichtenstein AH, et al. ACC/AHA Prevention Guideline: 2013 ACC/AHA Guideline on the Treatment of Blood Cholesterol to Reduce Atherosclerotic Cardiovascular Risk in Adults: A Report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines. erratum in *Circulation*. Jun. 24, 2014;129:S46-S48.
- Su, Kuan-Pin, et al., "Omega-3 Fatty Acids in Major Depressive Disorder A Preliminary Double-Blind, Placebo-Controlled Trial", *European Neuropsychopharmacology*, 13:267-271, (Aug. 2003).
- Sugiyama et al., "A Comparison of the Hypotensive Effects of Eicosapentaenoic Acid Ethyl (EPA) on Three Diseases (Occluded Arteriosclerosis, Hyperlipidemia, and These Two Diseases Combined) P2-504 Abstract," Annual Meeting of the Japanese Society of Pharmaceutical Health Care and Sciences 20:473 (Nov. 2010) (with English translation)(3 pages).
- Sugiyama, E., et al., "Eicosapentaenoic acid lowers plasma and liver cholesterol levels in the presence of peroxisome proliferators-activated receptor alpha," *Life Sciences*, 83:19-28 (Jul. 4, 2008/epub May 1, 2008).
- Superko et al., "Lipid Management to Reduce Cardiovascular Risk: A New Strategy is Required," *Circulation*, 117:560-568 (Jan. 29, 2008).
- Surette, M.E., et al., "Dependence on dietary cholesterol for n-3 polyunsaturated fatty acid induced changes in plasma cholesterol in the Syrian hamster," *J Lipid Res.*, 33:263-271 (Feb. 1992).
- Surette, M.E., et al., "Evidence for mechanisms of the hypotriglyceridemic effect of n-3 polyunsaturated fatty acids," *Biochimica et Biophysic Acta*, 1126: 199-205 (Jun. 22, 1992).
- Tagawa H, Shimokawa H, Tagawa T, et al. Long-term treatment with eicosapentaenoic acid augments both nitric oxide-mediated and non-nitric oxide-mediated endothelium-dependent forearm vasodilatation in patients with coronary artery disease. *J Cardiovasc Pharmacol* 33(4):633-40, Apr. 1999.
- Takaki A, Umamoto S, Ono K, Seki K, Ryoike T, Fujii A, Itagaki T, Harada M, Tanaka M, Yonezawa T, Ogawa H, Matsuzaki M. Add-on therapy of epa reduces oxidative stress and inhibits the progression

US 10,568,861 B1

Page 27

(56)

References Cited

OTHER PUBLICATIONS

of aortic stiffness in patients with coronary artery disease and statin therapy: A randomized controlled study. *J. Atheroscler. Thromb.* Jun. 23, 2011;18:857-866.

Takaku et al., Study on the Efficacy and Safety of Ethyl Eicosapentate (MND-21) in Treatment of Hyperlipidemia Based on a Long-Term Administration Test, 7 *J. Clin. Ther. Med.* 191 (1991) (with English Translation)(27 pages).

Talayero BG, Sacks FM. The role of triglycerides in atherosclerosis. *Curr. Cardiol. Rep.* 2011;13:544-552.

Tamura, et al., "Study of the Clinical Usefulness of Ethyl Eicosapentate (MND-21) in Long-Term Treatment of Hyperlipaemic Patients." *J Clin Thera & Medicines*, 7:1817-1834 (1991).

Tanaka et al., "Genome-Wide Association Study of Plasma Polyunsaturated Fatty Acids in the InCHIANTI Study." *PLoS Genetics* 5(1):1-8 (Jan. 2009).

Tanaka et al., "Suppression of prostaglandin synthesis by arachidonic acid or eicosapentaenoic acid in a macrophage-like cell line, RAW 264.7, treated with LPS," *Biol. Pharm. Bull.*, 22(10):1057-7 (Oct. 1999).

Tanaka et al., "Administration of high dose eicosapentaenoic acid enhances anti-inflammatory properties of high-density lipoprotein in Japanese patients with dyslipidemia," *Atherosclerosis*, 237(2):577-83 (Dec. 2014).

Tanaka et al., "Eicosapentaenoic Acid-Enriched High-Density Lipoproteins Exhibit Anti-Atherogenic Properties," *Circ. J.*, doi: 10.1253/circj.CJ-17-0294. [Epub ahead of print] (Jun. 23, 2017)(6 pages).

Tanaka, K.T., et al., "Reduction in the recurrence of stroke by eicosapentaenoic acid for hypercholesterolemic patients—Subanalysis of the JELIS trial." *Stroke*, 39(7):2052-8 (Jul. 2008/epub May 1, 2008).

Tatarczyk, et al., "Analysis of long-chain ω -3 fatty acid content in fish-oil supplements," *Wien Klin Wochenschr*, 119/13-14: 417-422 (2007).

Tatsuno et al., Efficacy and safety of TAK-085 compared with eicosapentaenoic acid in Japanese subjects with hypertriglyceridemia undergoing lifestyle modification: The omega-3 fatty acids randomized double-blind (ORL) study, *J. Clin. Lipid*; vol. 7(6), pp. 615-625 (Sep. 12, 2013).

Taylor et al., "Fish allergy: fish and products thereof," *Journal Food Science* (2004) 69.8 R175-R180.

Taylor, A.J., et al., "Arterial Biology for the Investigation of the Treatment Effects of Reducing Cholesterol (ARBITER) 2: a double-blind, placebo-controlled study of extended-release niacin on atherosclerosis progression in secondary prevention patients treated with statins", *Circulation*, 110:3512-3517, (Dec. 7, 2004/epub Nov. 10, 2004).

Tedgui, A., et al., "Anti-inflammatory mechanisms in the vascular wall." *Circ. Res.* 88:877-887 (May 11, 2001).

Teissier E, Nohara A, Chinetti G, Paumelle R, Cariou B, Fruchart JC, Brandes RP, Shah A, Staels B. Peroxisome proliferator-activated receptor alpha induces NADPH oxidase activity in macrophages, leading to the generation of LDL with PPAR-alpha activation properties. *Circ. Res.* Dec. 10, 2004/epub Nov. 11, 2004;95(12):1174-1182.

Teramoto T, Sasaki J, Ishibashi S, et al. Diagnosis of atherosclerosis. Executive Summary of the Japan Atherosclerosis Society (JAS) Guidelines for the Diagnosis and Prevention of Atherosclerotic Cardiovascular Diseases in Japan—2012 Version. *J Atheroscler Thromb.* 2014;21(4):296-8. Electronic publication Dec. 10, 2013.

Terano, et al., "Effect of Oral Administration of Highly Purified Eicosapentaenoic Acid on Platelet Function, Blood Viscosity and Red Cell Deformability in Healthy Human Subjects," *Atherosclerosis*, 46, 321-331 (Mar. 1983).

The TG and HDL Working Group of the Exome Sequencing Project, National Heart, Lung, and Blood Institute. Loss-of-function mutations in APOC3, triglycerides, and coronary disease. *N Engl J Med.* Jul. 3, 2014/epub Jun. 18, 2014; 371(1):22-31.

Theilla, M., et al., "A diet enriched in eicosapentaenoic acid, gamma-linolenic acid and antioxidants in the prevention of new pressure ulcer formation in critically ill patients with acute lung injury: A randomized, prospective, controlled study." *Clinical Nutrition* 26: 752-757 (Dec. 2007/epub Oct. 22, 2007).

Theobald et al., "LDL Cholesterol-Raising Effect of Low-Dose Docosahexaenoic Acid in Middle-Aged Men and Women," *Am. J. Clin. Nutr.* 79:558-63 (Apr. 2004).

Thies, F., et al., "Association of n-3 polyunsaturated fatty acids with stability of atherosclerotic plaques: a randomised controlled trial." *Lancet* 361: 477-85 (Feb. 8, 2003).

Thies, F., et al., "Dietary supplementation with eicosapentaenoic acid, but not with other long-chain n-3 or n-6 polyunsaturated fatty acids, decreases natural killer cell activity in healthy subjects aged >55 y." *Am J Clin Nutr* 73:539-48 (Mar. 2001).

Third Report of the NCEP Expert Panel on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III) Final Report, NIH Publication No. 02-5215 Sep. 2002 (220 pages in three parts).

Thomas et al., "Renal Failure—Measuring the Glomerular Filtration Rate," *Dtsch Arztebl Int.*, Dec. 18, 2009, 106(51-52); 849-54.

Thorwest M, Balling E, Kristensen SD, et al. Dietary fish oil reduces microvascular thrombosis in a porcine experimental model. *Thromb. Res.* Jul. 2000, 99(2): 203-8.

Thygesen K., Alpert J., Jaffe A., et al. Third Universal Definition of Myocardial Infarction. *J Am Coll Cardiol.*, Oct. 16, 2012/epub Sep. 5, 2012; 60(16):1581-1598.

Tilg H, Moschen AR. Inflammatory Mechanisms in the Regulation of Insulin Resistance. *Mol. Med.*, Mar./Apr. 2008;14(3-4):222-231.

Tirosh et al., "Changes in Triglyceride Levels and Risk for Coronary Heart Disease in Young Men," *American College of Physicians*, pp. 377-385 (Sep. 18, 2007).

Torreson, C. et al., "n-3 Fatty acids and cardiovascular disease: Actions and molecular mechanisms," *Prostaglandins Leukotrienes & Essent. Fatty Acids*, 77(5-6):319-26 Nov./Dec. 2007/epub Dec. 3, 2007. doi:10.1016/j.plefa.2007.10.014 (2007).

Toth PP, Granowitz C, Hull M, Liassou D, Anderson A, Philip S. High Triglycerides are associated with increased cardiovascular events, medical costs, and resource use: A real-world administrative claims analysis of statin-treated patients with high residual cardiovascular risk. *Journal of the American Heart Association*, 7(15):e008740 (publication date Jul. 25, 2018; epublication Aug. 7, 2018).

Transcript from Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, 76 pages.

TREND-HD Investigators, Randomized controlled trial of ethyl-eicosapentaenoic acid in Huntington disease: the TREND-HD study, *Arch Neurol.*, vol. 65(12): 1582-9 (Dec. 2008).

Tribble DL, Holl LG, Wood PD, Krauss RM. Variations in oxidative susceptibility among six low density lipoprotein subfractions of differing density and particle size. *Atherosclerosis*. Apr. 1992;93(3):189-199.

Tribble DL, Rizzo M, Chait A, Lewis DM, Blanche PJ, Krauss RM. Enhanced oxidative susceptibility and reduced antioxidant content of metabolic precursors of small, dense low-density lipoproteins. *Am. J. Med.* Feb. 1, 2001;110(2):103-110.

Trilipix Package Insert (Sep. 2010)(10 pages).

Tsimikas S, Witztum JL, Miller ER, Sasiela WJ, Szarek M, Olsson AG, Schwartz GG. High-dose atorvastatin reduces total plasma levels of oxidized phospholipids and immune complexes present on apolipoprotein B-100 in patients with acute coronary syndromes in the MIRACL trial. *Circulation.*, Sep. 14, 2004/epub Sep. 7, 2004; 110(11):1406-1412.

Tsuruta K., et al., "Effects of purified eicosapentaenoate ethyl ester on fibrinolytic capacity in patients with stable coronary artery disease and lower extremity ischaemia" *Coron Artery Dis.* 7(11):837-42 (Nov. 1996).

Tulenok TN, Chen M, Mason PE, Mason RP. Physical effects of cholesterol on arterial smooth muscle membranes: Evidence of immiscible cholesterol domains and alterations in bilayer width C during atherogenesis. *J. Lipid Res.* May 1998;39:947-956.

Tungsiripat, et al., "Dyslipidemia in HIV patients," *Cleveland Clinic Journal of Medicine*, v. 72, No. 12 (Dec. 2005).

US 10,568,861 B1

Page 28

(56)

References Cited

OTHER PUBLICATIONS

Turini et al., "Short-term fish oil supplementation improved innate immunity, but increased ex vivo oxidation of LDL in man—a pilot study." *Eur. J. Nutr.* 40:56-65 (Apr. 2001).

U.S. Appl. No. 14/245,499, filed Apr. 4, 2014 (now abandoned)(43 pages).

Ullian, M.E., "Fatty acid inhibition of angiotensin II-stimulated inositol phosphates in smooth muscle cells." *Am J Physiol Heart Circ Physiol.*, 264 (2 Pt 2):H595-603 (Feb. 1993).

Urakaze, Masaharu, et al., "Infusion of emulsified triicosapentaenylglycerol into rabbits. The effects on platelet aggregation, polymorphonuclear leukocyte adhesion, and fatty acid composition in plasma and platelet phospholipids", *Thromb. Res.*, 44(5):673-682 (Dec. 1986).

Urquhart et al., "Profile of eicosanoids produced by human saphenous vein endothelial cells and the effect of dietary fatty acids," *Prostaglandins Leukot. Essent. Fatty Acid*, 65(1):15-22 (Jul. 2001).

US Food and Drug Administration and Dept of Health and Human Services. Substances affirmed as generally recognized as safe: Menhaden Oil. *Fed Register*, 62:30751-30757 (Jun. 5, 1997).

Vaagenes et al., "The Hypolipidaemic Effect of EPA is Potentiated by 2- and 3-Methylation." In P. Quant & S. Eaton (eds.) *Current Views of Fatty Acid Oxidation and Ketogenesis from Organelles to Point Mutations; Advances in Experimental Medicine and Biology*, vol. 466, pp. 221-226 (1999).

Vaddadi, K.S., et al., "A Randomised, Placebo-Controlled, Double-Blind Study of Treatment of Huntington's Disease with Unsaturated Fatty Acids", *Clinical Neuroscience and Neuropathology*, 13(1):29-33, (Jan. 2002).

Vaduganathan M, Venkataramani AS, Bhatt DL. Moving toward global primordial prevention in cardiovascular disease: The heart of the matter. *J Am Coll Cardiol* Oct. 6, 2015;66(14):1535-7.

Van der Steeg, W.A., et al., "High-density lipoprotein cholesterol, high-density lipoprotein particle size, and apolipoprotein A-I: Significance for cardiovascular risk—the IDEAL and EPIC-Norfolk studies." *J. Am. Coll. Cardiol.* 51:634-642 (Feb. 12, 2008).

Van Do et al., "Allergy to fish parvalbumins: Studies on the cross-reactivity of allergens from 9 commonly consumed fish." *Journ. Allergy & Clin. Immunol.*, 16(6):1314-1320 (Dec. 1, 2005).

Van Wijk et al. Rosiglitazone improves postprandial triglyceride and free fatty acid metabolism in type 2 diabetes. *Diabetes Care*, vol. 28, No. 4, (Apr. 2005) pp. 844-849.

Varbo A, Benn M, Tybjaerg-Hansen A, Nordestgaard BG. Reply to letters regarding article, "Elevated remnant cholesterol causes both low-grade inflammation and ischemic heart disease, whereas elevated low-density lipoprotein cholesterol causes ischemic heart disease without inflammation". *Circulation*. Jun. 17, 2014; 129(24):e656.

Varbo et al., Remnant Cholesterol as a Causal Risk Factor for Ischemic Heart Disease, *J. Am. Coll. Cardiol.*, vol. 61(4), pp. 427-436 (Jan. 29, 2013/epub Dec. 19, 2012).

Varbo et al., Remnant cholesterol as a cause of ischemic heart disease: Evidence, definition, measurement, atherogenicity, high risk patients, and present and future treatment, *Pharmacol. Ther.*, vol. 141(3), pp. 358-367 (Mar. 2014/epub Nov. 26, 2013).

Vascepa [package insert], Bedminster, NJ: Amarin Pharma Inc.; Jul. 2012. (12 pages).

Vascepa [package insert]. Bedminster, NJ: Amarin Pharma Inc.; Nov. 2013. (11 pages).

Vasudevan et al., "Effective Use of Combination of Lipid Therapy", *Curr. Atheroscl. Rep.*, vol. 8, pp. 76-84 (Jan. 2006).

Vedin, I., et al., "Effects of docosahexaenoic acid—rich n-3 fatty acid supplementation on cytokine release from blood mononuclear leukocytes: the OmegAD study." *Am J Clin Nutr* 87:1616-22 (Jun. 2008).

Velliquette et al., "Regulation of human stearoyl-CoA desaturase by omega-3 and omega-6 fatty acids: Implications for the dietary management of elevated serum triglycerides," *Journal of Clinical Lipidology*. (Aug. 2009/epub Jun. 21, 2009) 3:281-288.

Vergnani L, Hatrik S, Ricci F, Passaro A, Manzoli N, Zuliani G, Brovkovich V, Fellin R, Malinski T. Effect of native and oxidized

low-density lipoprotein on endothelial nitric oxide and superoxide production : Key role of l-arginine availability. *Circulation*. March 21, 2000; 101 :1261-1266.

Verma S, Leiter LA, Bhatt DL. CANTOS ushers in a new calculus of inflammasome targeting for vascular protection-and maybe more. *Cell Metab* 26(5):703-5 (publication date Nov. 7, 2017; epublication date Oct. 19, 2017).

Vidal F, Colome C, Martinez-Gonzalez J, Badimon L. Atherogenic concentrations of native low density lipoproteins down-regulate nitric-oxide-synthase mma and protein levels in endothelial cells. *Eur. J. Biochem.* Mar. 15, 1998; 252:378-384.

Vidgren, H.M., et al., "Incorporation of n-3 fatty acids into plasma lipid fractions, and erythrocyte membranes and platelets during dietary supplementation with fish, fish oil, and docosahexaenoic acid-rich oil among healthy young men." *Lipids* 32: 697-705 (Jul. 1997).

Virani et al., "The Role of Lipoprotein-associated Phospholipase A2 as a marker for atherosclerosis" *Curr. Atheroscler. Rep.* 9(2): 97-103 (Aug. 2007).

Volcik, K.A., et al., "Peroxisome proliferator-activated receptor α genetic variation interacts with n-6 and long-chain n-3 fatty acid intake to affect total cholesterol and LDL-cholesterol concentrations in the Atherosclerosis Risk in Communities Study." *Am J Clin Nutr* 87:1926-31 (Jun. 2008).

Von Schacky C, Baumann K, Angerer P. The effect of n-3 fatty acids on coronary atherosclerosis: results from SCIMO, an angiographic study, background and implications. *Lipids* 2001 36 Suppl: S99-102.

Von Schacky, C., "A review of omega-3 ethyl esters for cardiovascular prevention and treatment of increased blood triglyceride levels." *Vascular Health and Risk Management* 2(3): 251-262 (2006).

Von Schacky, C., et al., "The Effect of Dietary ω -3 Fatty Acids on Coronary Atherosclerosis: A Randomized, Double-Blind, Placebo-Controlled Trial", *American College of Physicians-American Society of Internal Medicine*, 130(7):554-562, (Apr. 6, 1999).

Wada, M., et al., "Enzymes and receptors of prostaglandin pathways with arachidonic acid-derived versus eicosapentaenoic acid-derived substrates and products." *J. Biol. Chem.* 282(31): 22254-22266 (Aug. 3, 2007/epub May 22, 2007).

Wagner AH, Kohler T, Ruckschloss U, Just I, Hecker M. Improvement of nitric oxide-dependent vasodilation by hmg-coa reductase inhibitors through attenuation of endothelial superoxide anion formation. *Arterioscler. Thromb. Vasc. Biol.*, Jan. 2000;20:61-69.

Walker G, Mandagere A, Dufton C, et al. The pharmacokinetics and pharmacodynamics of warfarin in combination with ambrisentan in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2009/epub Feb. 4, 2009; 67 (5): 527-34.

Wall R, Ross RP, Fitzgerald G, Stanton C. Fatty acids from fish: the anti-inflammatory potential of long-chain omega-3 fatty acids. *Nutr Rev.* May 2010; 68:280-289.

Walldius, G., et al., "Editorial: Rationale for using apolipoprotein B and apolipoprotein A-I as indicators of cardiac risk and as targets for lipid-lowering therapy." *European Heart Journal* 26, 210-212 (Feb. 2005/epub Dec. 15, 2004).

Walter MF, Jacob RF, Bjork RE, Jeffers B, Buch J, Mizuno Y, Mason RP. Circulating lipid hydroperoxides predict cardiovascular events in patients with stable coronary artery disease: the PREVENT study. *J. Am. Coll. Cardiol.*, Mar. 25, 2008;51(12):1196-1202.

Walter MF, Jacob RF, Jeffers B, Ghadanfar MM, Preston GM, Buch J, Mason RP. Serum levels of thiobarbituric acid reactive substances predict cardiovascular events in patients with stable coronary artery disease: A longitudinal analysis of the PREVENT study. *J. Am. Coll. Cardiol.* Nov. 16, 2004; 44(10):1996-2002.

Wander, R.C., et al., "Influence of long chain polyunsaturated fatty acids on oxidation of low density lipoprotein." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 59(2):143-151 (Aug. 1998).

Wang Q, Liang X, Wang L, Lu X, Huang J, Cao J, Li H, Gu D. Effect of omega-3 fatty acids supplementation on endothelial function: A meta-analysis of randomized controlled trials. *Atheroscler.* Apr. 2012/epub Jan. 20, 2012; 221:563-543.

US 10,568,861 B1

Page 29

(56)

References Cited

OTHER PUBLICATIONS

- Wang, C., et al., "n-3 Fatty acids from fish or fish-oil supplements, but not α -linolenic acid, benefit cardiovascular disease outcomes in primary- and secondary-prevention studies: a systematic review." *Am J Clin Nutr* 84:5-17 (Jan. 2006).
- Wang, L., et al., "Triglyceride-rich lipoprotein lipolysis releases neutral and oxidized FFAs that induce endothelial cell inflammation." *J. Lipid Res.* 50:204-213 (Feb. 2009/epub Sep. 23, 2008).
- Warren, Stephen T., "The Expanding World of Trinucleotide Repeats", *Science*, 271:1374-1375, (Mar. 8, 1996).
- Wassmann S, Laufs U, Muller K, Konkol C, Ahlborn K, Baumer AT, Linz W, Bohm M, Nickenig G. Cellular antioxidant effects of atorvastatin in vitro and in vivo. *Arterioscler. Thromb. Vasc. Biol.* Feb. 1, 2002; 22:300-305.
- Watanabe et al., "Bile acids lower triglyceride levels via a pathway involving FXR, SHP, and SREBP-1c," *J Clin Invest.* 113(10): 1408-1418 (May 2004).
- Watanabe T, Ando K, Daidoji H, et al. A randomized controlled trial of eicosapentaenoic acid in patients with coronary heart disease on statins. *J Cardiol* 70(6):537-44 (publication date Dec. 2017; epublication date Aug. 31, 2017).
- Watanabe, Ikuyoshi, et al., "Usefulness of EPA-E (eicosapentaenoic acid ethyl ester) in preventing neointimal formation after vascular injury", *Kokyu to Junkan*, 42(7):673-677 (1994) (with English summary).
- Weaver, K.L., et al., "Effect of Dietary Fatty Acids on Inflammatory Gene Expression in Healthy Humans." *J. Biol. Chem.*, 284(23): 15400-15407 (2009) (published online Apr. 9, 2009).
- Webcast Information for the Oct. 16, 2013 Meeting of the Endocrinologic and Metabolic Drugs Advisory Committee, (1 page).
- Weber, P. "Triglyceride-lowering effect of n-3 long chain polyunsaturated fatty acid: eicosapentaenoic acid vs. docosahexaenoic acid." *Lipids* 34: S269 (1999).
- Wei et al., Effects of [EPA] Versus [DHA] on Serum Lipids: A Systematic Review and Meta-Analysis, 13 *Current Atherosclerosis Rep.* 13(6):474-483 (Dec. 2011).
- Wei L.J, Lin D.Y, Weissfeld L. Regression analysis of multivariate incomplete failure time data by modeling marginal distributions. *J Am Stat Assoc.* 84(408):1065-1073 (publication date Dec. 1989).
- Westerveld H.T. et al., "Effects of low-dose EPA-Eon glycemic control, lipid profile, lipoprotein(a), platelet aggregation, viscosity, and platelet and vessel wall interaction in NIDDM" *Diabetes Care* 16(5):683-8 (May 1993).
- Westphal, S., et al., "Postprandial chylomicrons and VLDLs in severe hypertriglyceridemia are lowered more effectively than are chylomicron remnants after treatment with n3 fatty acids." *Am J Clin Nutr* 71:914-20 (Apr. 2000).
- Whelan, "Evidence that dietary arachidonic acid increases circulating triglycerides." *Lipids* 30, 425-429 (May 1995).
- Wierzbicki, A.S., "Editorial: Newer, lower, better? Lipid drugs and cardiovascular disease—the continuing story." *Int J Clin Pract*, 61(7):1064-1067 (Jul. 2007).
- Wierzbicki, A.S., "Editorial: Raising HDL-C: back to the future?" *Int J Clin Pract*, 61(7): 1069-1071 (Jul. 2007).
- Wikipedia, "Diabetes mellitus," Dec. 12, 2016 (Dec. 12, 2016), retrieved on Jul. 30, 2018 from https://en.wikipedia.org/w/index.php?title=Diabetes_mellitus&oldid=754431573; entire document, especially p. 1, paragraph 1.
- Wikipedia, "Ethyl eicosapentaenoic acid," Apr. 1, 2016 (Apr. 1, 2016); retrieved on Jul. 27, 2018 from https://en.wikipedia.org/w/index.php?title=Ethyl_eicosapentaenoic_acid&oldid=713086755; entire document, especially p. 1, col. 2 and p. 3, paragraph 2.
- Williams et al., "NADPH Oxidase Inhibitors New Antihypertensive Agents?" *J. Cardiovasc Pharmacol* 50(1):9-16 (Jul. 1, 2007).
- Willumsen, N. et al., *Biochimica et Biophysica Acta*. vol. 1369, "On the effect of 2-deuterium- and 2-methyl-eicosapentaenoic acid derivatives on triglycerides, peroxisomal beta-oxidation and platelet aggregation in rats," pp. 193-203, (Mar. 2, 1998).
- Willumsen, N., et al., "Eicosapentaenoic acid, but not docosahexaenoic acid, increased, mitochondrial fatty acid oxidation and upregulates 2,3-dienoyl-CoA reductase gene expression in rats." *Lipids*, 31:579-592 (Jun. 1996).
- Wilson Omega 3 fish oil: EPA versus DHA (Dietivity.com, 1-16) (2006).
- Wilt, VM & Gumm, JG, "Isolated low high-density lipoprotein cholesterol", *Ann. Pharmacol.*, 31:89-97, (Jan. 1997).
- Wink, J. et al., "Effect of very-low-dose niacin on high-density lipoprotein in patients undergoing long-term statin therapy", *Am. Heart J.*, 143:514-518, (Mar. 2002).
- Wittrup HH, Tybjaerg-Hansen A, Nordestgaard BG. Lipoprotein lipase mutations, plasma lipids and lipoproteins, and risk of ischemic heart disease: a meta-analysis. *Circulation.*, Jun. 8, 1999;99:2901-2907.
- Witztum J.L. The oxidation hypothesis of atherosclerosis. *Lancet*, Sep. 17, 1994;344(8925):793-795.
- Wojczynski et al., "High-fat meal effect on LDL, HDL and VLDL particle size and number in the Genetics of Lipid-Lowering Drugs and Diet Network (GOLDN): an interventional study," *Lipids in Health and Disease* 10:181, pp. 1-11 (Oct. 18, 2011).
- Wojenski, C.M., et al., "Eicosapentaenoic acid ethyl ester as an antithrombotic agent: comparison to an extract of fish oil." *Biochimica et Biophysica Acta*. 1081:33-38 (Jan. 4, 1991).
- Wong, S.H., et al., "Effects of eicosapentaenoic and docosahexaenoic acids on Apolipoprotein B mRNA and secretion of very low density lipoprotein in HepG2 cells." *Arterioscler. Thromb. Vasc. Biol.* 9:836-841 (Nov./Dec. 1989).
- Wood et al., "Carbohydrate Restriction Alters Lipoprotein Metabolism by Modifying VLDL, LDL and HDL Subfraction Distribution and Size in Overweight Men," *Journ. of Nutrition*, 136(2):384-9 (Feb. 2006).
- Woodman et al., "Effects of Purified Eicosapentaenoic and Docosahexaenoic Acids on Glycemic Control, Blood Pressure, and Serum Lipids in Type 2 Diabetic Patients with Treated Hypertension", *The American Journal of Clinical Nutrition: Official Journal of the American Society for Clinical Nutrition, Inc.*, 76(5):1007-1015 (Nov. 1, 2002).
- Woodman, R.J., et al., "Effects of purified eicosapentaenoic acid and docosahexaenoic acid on platelet, fibrinolytic and vascular function in hypertensive type 2 diabetic patients." *Atherosclerosis* 166: 85-93 (Jan. 2003).
- Wu et al., "Diabetic dyslipidemia," *Metabolism Clinical and Experimental*, 63:1469-1479 (Dec. 2014)(available online Aug. 29, 2014).
- Wu, W.H., et al., "Effects of docosahexaenoic acid supplementation on blood lipids, estrogen metabolism, and in vivo oxidative stress in postmenopausal vegetarian women." *Eur J Clin Nutr.*, 60:386-392 (Mar. 2006).
- Xiao, Y.F., et al., "Inhibitory effect of n-3 fish oil fatty acids on cardiac Na⁺/Ca²⁺ exchange currents in HEK293t cells." *Biochemical and Biophysical Research Communications* 321: 116-123 (Aug. 13, 2004).
- Xiao, Y-F., et al., "Blocking effects of polyunsaturated fatty acids on Na⁺ channels of neonatal rat ventricular myocytes." *Proc. Natl. Acad. Sci.* 92: 11000-11004 (Nov. 21, 1995).
- Xiao, Y-F., et al., "Fatty acids suppress voltage-gated Na⁺ currents in HEK293t cells transfected with the α -subunit of the human cardiac Na⁺ channel." *Proc. Natl. Acad. Sci.* 95: 2680-2685 (Mar. 3, 1998).
- Xydakis, AM et al., "Combination therapy for combined dyslipidemia," *American Journal of Cardiology*, Nov. 20, 2002 US, vol. 90, No. 10 Suppl. 2, p. 21 K-29K (Nov. 20, 2002).
- Yacyszyn BR, Thomson AB. The clinical importance of proton pump inhibitor pharmacokinetics. *Digestion* 2002 66 (2): 67-78.
- Yadav D, Pitchumoni CS. Issues in Hyperlipidemic Pancreatitis. *J Clin Gastroenterol* 236(1):54-62, Jan. 2003.
- Yagi K. Assay for blood plasma or serum. *Methods Enzymol.* 1984;105:328-331.
- Yamagishi K, Nettleton J, Folsom A. Plasma fatty acid composition and incident heart failure in middle-aged adults: The Atherosclerosis Risk in Communities (ARIC) Study. *Am Heart J.*, Nov. 2008/epub Aug. 29, 2008; 156:965-974.

(56)

References Cited

OTHER PUBLICATIONS

- Yamakawa K, Shimabukuro M, Higa N, Asahi T, Ohba K, Arasaki O, Higa M, Oshiro Y, Yoshida H, Higa T, Saito T, Ueda S, Masuzaki H, Sata M. Eicosapentaenoic Acid Supplementation Changes Fatty Acid Composition and Corrects Endothelial Dysfunction in Hyperlipidemic Patients. *Cardiol Res Practice*. Dec. 26, 2012; epub Article ID 754181.
- Yamamoto, H. et al., Improvement of coronary vasomotion with Eicosapentaenoic acid does not inhibit acetylcholine-induced coronary vasospasm in patients with variant angina: *Jpn Cir J*. 59(9):608-16 (Sep. 1995).
- Yamamoto, K., et al., "4-Hydroxydocosaheptaenoic acid, a potent Peroxisome Proliferator-Activated Receptor C agonist alleviates the symptoms of DSS-induced colitis." *Biochemical and Biophysical Research Communications* 367: 566-572 (Mar. 14, 2008/epub Jan. 10, 2008).
- Yamano T, Kubo T, Shiono Y, et al. Impact of eicosapentaenoic acid treatment on the fibrous cap thickness in patients with coronary atherosclerotic plaque: an optical coherence tomography study. *J Atheroscler Thromb*. 2015/epub Aug. 15, 2014;22:52-61.
- Yamashita et al., *J. Biochem.*, vol. 122, No. 1, "Acyl-transferases and Transacylases Involved in Fatty Acid Remodeling of Phospholipids and Metabolism of Bioactive Lipids in Mammalian Cells", pp. 1-16 (Jul. 1997).
- Yamashita, N., et al., "Inhibition of natural killer cell activity of human lymphocytes by eicosapentaenoic acid." *Biochem. Biophys. Res. Comm.* 138(3): 1058-1067 (Aug. 25, 1986).
- Yamazaki et al., "Changes in fatty acid composition in rat blood and organs after infusion of eicosapentaenoic acid ethyl ester", *Biochim. Biophys. ACTA*, 1128(1):35-43, (Sep. 22, 1992).
- Yamazaki, et. al., "Dissolution tests by RDC method for soft gelatin capsules containing ethyl icosapentate", *Pharm. Tech. Japan*, vol. 15, No. 4, pp. 595-603 Abstract (Apr. 1999) (with English abstract).
- Yang, S.P., et al., "Eicosapentaenoic acid attenuates vascular endothelial growth factor-induced proliferation via inhibiting Flk-1 receptor expression in bovine carotid artery endothelial cells." *J. Cell. Physio.* 176:342-349 (Aug. 1998).
- Yano T, Mizuguchi K, Takasugi K, Tanaka Y, Sato M. "Effects of ethyl all-cis-5,8,11,14,17-icosapentaenoate on low density lipoprotein in rabbits," *Yakugaku Zasshi*, 115:843-51 (Oct. 1995)
- Yano, T., et al., "Effects of ethyl-all-cis-5,8,11,14,17-icosapentaenoate (EPA-E), pravastatin and their combination on serum lipids and intimal thickening of cuff-sheathed carotid artery in rabbits." *Life Sciences*, 61(20):2007-2015 (1997).
- Yao et al., "Oxidized high density lipoprotein induces macrophage apoptosis via toll-like receptor 4-dependent CHOIP pathway," *Journ. Lipid Res.*, 58:164-177 (Jan. 2017)(First published Nov. 28, 2016).
- Yates RA, Wong J, Seiberling M, et al. The effect of anastrozole on the single-dose pharmacokinetics and anticoagulant activity of warfarin in healthy volunteers. *Br. J. Clin. Pharmacol.* May 2001 51(5): 429-35.
- Yerram, N.R., et al., "Eicosapentaenoic acid metabolism in brain microvessel endothelium: effect on prostaglandin formation." *J. Lipid Res.*30:1747-1757 (Nov. 1989).
- Yokoyama et al., "Effects of eicosapentaenoic acid on cardiovascular events in Japanese patients with hypercholesterolemia: Rationale, design, and baseline characteristics of the Japan EPA Lipid Intervention Study (JELIS)," *Amer. Heart Journal* 146(4):613-620 (Oct. 2003).
- Yokoyama et al., "Effects of eicosapentaenoic acid on major coronary events in hypercholesterolaemic patients (JELIS): a randomized open-label, blinded endpoint analysis", *Lancet*, vol. 369, pp. 1090-1098 (Mar. 31, 2007).
- Yorioka, N, "Lipid-lowering therapy and coagulation/fibrinolysis parameters in patients on peritoneal dialysis," *The International Journal of Artificial Organs*, vol. 23(1):27-32 (Jan. 2000).
- Yoshimura et al., "Effects of highly purified eicosapentaenoic acid on plasma beta thromboglobulin level and vascular reactivity to angiotensin II", *Artery*, 14(5):295-303 (1987).
- Zaima, N., et al., "Trans geometric isomers of EPA decrease LXRA-induced cellular triacylglycerol via suppression of SREBP-1c and PGC-1 β ," *J. Lipid Res.* 47: 2712-2717 (Dec. 2006).
- Zalewski et al., Role of Lipoprotein-Associated Phospholipase A2 in Atherosclerosis: Biology, Epidemiology, and Possible Therapeutic Target, *Arteriosclerosis, Thrombosis, & Vascular Biology* 25(5):923-931 (May 2005/epub Feb. 24, 2005).
- Zanarini, et al., "Omega-3 Fatty Acid Treatment of Women with Borderline Personality Disorder: A Double-Blind, Placebo-Controlled Pilot Study," *Am J Psychiatry*, 160:167-169 (Jan. 2003).
- Zhan, S. et. al. "Meta-analysis of the effects of soy protein containing isoflavones on the lipid profile," *Am. J. Clin. Nutr.* (Feb. 2005), 81, p. 397-408.
- Zhang, M., et al., "Effects of eicosapentaenoic acid on the early stage of type 2 diabetic nephropathy in KKAY/Ta mice: involvement of anti-inflammation and antioxidative stress." *Metabolism Clinical and Experimental* 55:1590-1598 (Dec. 2006).
- Zhang, Y.W., et al., "Inhibitory effects of eicosapentaenoic acid (EPA) on the hypoxia/reoxygenation-induced tyrosine kinase activation in cultured human umbilical vein endothelial cells." *Prostaglandins, Leukotrienes and Essential FattyAcids* 67(4):253-261 (Oct. 2002).
- Zhang, Y.W., et al., "Pretreatment with eicosapentaenoic acid prevented hypoxia/reoxygenation-induced abnormality in endothelial gap junctional intercellular communication through inhibiting the tyrosine kinase activity." *Prostaglandins, Leukotrienes and Essential Fatty Acids* 61(1): 33-40 (Jul. 1999).
- Zhao et al., "Polyunsaturated Fatty Acids are FXR Ligands and Differentially Regulate Expression of FXR Targets," *DNA and Cell Biology*, 23(8):519-526 (Aug. 25, 2004).
- Zhao, G. et al., "Dietary α -linolenic acid inhibits proinflammatory cytokine production by peripheral blood mononuclear cells in hypercholesterolemic subjects." *Am J Clin Nutr* 85:385-91 (Feb. 2007).
- Zhao, G., et al., "Dietary α -linolenic acid reduces inflammatory and lipid cardiovascular risk factors in hypercholesterolemic men and women." *J. Nutr.* 134: 2991-2997 (Nov. 2004).
- Zheng et al., "Function of ω -3 long chain unsaturated fatty acid in metabolic syndrome," *Chinese Journal of Endocrinology and Metabolism*, vol. 27, No. 9, pp. 787-790 (Sep. 30, 2011)(with English translation).
- Ziegler, D., et al., "Treatment of symptomatic diabetic polyneuropathy with the antioxidant α -lipoic acid: A 7-month multicenter randomized controlled trial (ALADIN III Study)." *Diabetes Care* 22:1296-1301 (Aug. 1999).
- Zimmerman JJ, Raible DG, Harper DM, et al. Evaluation of a potential tigecycline-warfarin drug interaction. *Pharmacotherapy* Jul. 28, 2008 (7): 895-905.
- Zuijgeest-van Leeuwen, et al., "N-3 Fatty Acids Administered as Triacylglycerols or as Ethyl Esters Have Different Effects on Serum Lipid Concentrations in Healthy Subjects," *N-3 Fatty Acids, Lipid Metabolism and Cancer*, pp. 89-100 (2000).
- Zuijgeest-van Leeuwen, S.D., et al., "Incorporation and washout of orally administered n-3 fatty acid ethyl esters in different plasma lipid fractions." *British Journal of Nutrition* 82:481-488 (1999).
- Zuijgeest-van Leeuwen, SD, et al., "Eicosapentaenoic acid inhibits lipolysis in weight-losing cancer patients as well as in healthy volunteers," *Eur J Gastroenterol & Hepatol.*, 10(12):A67 (1998).
- Zvyaga T, Chang SY, Chen C, et al. Evaluation of six proton pump inhibitors as inhibitors of various human cytochromes P450: focus on cytochrome P450 2C19. *Drug Metab. Dispos.* Sep. 2012 40(9): 1698-711.
- Zimmer et al., "Danger signaling in Atherosclerosis," *Circ. Res.*, 2015; 116:323-340.

US 10,568,861 B1

1

**METHODS OF REDUCING THE RISK OF A
CARDIOVASCULAR EVENT IN A SUBJECT
AT RISK FOR CARDIOVASCULAR DISEASE**

PRIORITY CLAIM

This application is a divisional of U.S. patent application Ser. No. 16/502,621 filed Jul. 3, 2019, which is a continuation of U.S. patent application Ser. No. 16/287,157 filed Feb. 27, 2019 (now U.S. Pat. No. 10,383,840), which is a continuation of U.S. patent application Ser. No. 16/005,852 filed Jun. 12, 2018 (now U.S. Pat. No. 10,278,935), which is a continuation of U.S. patent application Ser. No. 15/886,422 filed Feb. 1, 2018 (now U.S. Pat. No. 10,016,386), which is a continuation application of U.S. patent application Ser. No. 15/607,084 filed May 26, 2017 (now U.S. Pat. No. 9,918,955), which is a continuation of U.S. patent application Ser. No. 15/427,238 filed Feb. 8, 2017 (now U.S. Pat. No. 9,693,986), which is a continuation application of U.S. patent application Ser. No. 15,333,991 filed Oct. 25, 2016 (now U.S. Pat. No. 9,610,272), which is a continuation of U.S. patent application Ser. No. 14/411,815, filed Dec. 29, 2014 (now U.S. Pat. No. 9,603,826), which is a 371 national stage application of PCT/US2013/048559 filed Jun. 28, 2013, and which claims priority to U.S. provisional patent application Ser. No. 61/666,447, filed Jun. 29, 2012, the entire contents of which are incorporated herein by reference.

BACKGROUND

Cardiovascular disease is one of the leading causes of death in the United States and most European countries. It is estimated that over 70 million people in the United States alone suffer from a cardiovascular disease or disorder including but not limited to high blood pressure, coronary heart disease, dyslipidemia, congestive heart failure and stroke.

Lovaza®, a lipid regulating agent, is indicated as an adjunct to diet to reduce triglyceride levels in adult patients with very high triglyceride levels. Unfortunately, Lovaza® can significantly increase LDL-C and/or non-HDL-C levels in some patients. A need exists for improved treatments for cardiovascular diseases and disorders.

SUMMARY

In various embodiments, the present invention provides methods of reducing the risk of a cardiovascular event in a subject on statin therapy. In one embodiment, the method comprises administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of eicosapentaenoic acid ethyl ester or a derivative thereof. In another embodiment, the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL. In another embodiment, the composition contains not more than 10%, by weight, docosahexaenoic acid or derivative thereof, substantially no docosahexaenoic acid or derivative thereof, or no docosahexaenoic acid or derivative thereof. In another embodiment, eicosapentaenoic acid ethyl ester comprises at least 96%, by weight, of all fatty acids present in the composition; the composition contains not more than 4%, by weight, of total fatty acids other than eicosapentaenoic acid ethyl ester; and/or the composition contains about 0.1% to about 0.6% of at least one fatty acid other than eicosapentaenoic acid ethyl ester and docosahexaenoic acid.

2

In another embodiment, the invention provides a method of treating hypertriglyceridemia comprising administering a composition as described herein to a subject in need thereof one to about four times per day.

These and other embodiments of the present invention will be disclosed in further detail herein below.

DETAILED DESCRIPTION

While the present invention is capable of being embodied in various forms, the description below of several embodiments is made with the understanding that the present disclosure is to be considered as an exemplification of the invention, and is not intended to limit the invention to the specific embodiments illustrated. Headings are provided for convenience only and are not to be construed to limit the invention in any manner. Embodiments illustrated under any heading may be combined with embodiments illustrated under any other heading.

The use of numerical values in the various quantitative values specified in this application, unless expressly indicated otherwise, are stated as approximations as though the minimum and maximum values within the stated ranges were both preceded by the word "about." Also, the disclosure of ranges is intended as a continuous range including every value between the minimum and maximum values recited as well as any ranges that can be formed by such values. Also disclosed herein are any and all ratios (and ranges of any such ratios) that can be formed by dividing a disclosed numeric value into any other disclosed numeric value. Accordingly, the skilled person will appreciate that many such ratios, ranges, and ranges of ratios can be unambiguously derived from the numerical values presented herein and in all instances such ratios, ranges, and ranges of ratios represent various embodiments of the present invention.

Compositions

In one embodiment, a composition of the invention is administered to a subject in an amount sufficient to provide a daily dose of eicosapentaenoic acid of about 1 mg to about 10,000 mg, 25 about 5000 mg, about 50 to about 3000 mg, about 75 mg to about 2500 mg, or about 100 mg to about 1000 mg, for example about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225 mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about

US 10,568,861 B1

3

2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, about 2500 mg, about 2525 mg, about 2550 mg, about 2575 mg, about 2600 mg, about 2625 mg, about 2650 mg, about 2675 mg, about 2700 mg, about 2725 mg, about 2750 mg, about 2775 mg, about 2800 mg, about 2825 mg, about 2850 mg, about 2875 mg, about 2900 mg, about 2925 mg, about 2950 mg, about 2975 mg, about 3000 mg, about 3025 mg, about 3050 mg, about 3075 mg, about 3100 mg, about 3125 mg, about 3150 mg, about 3175 mg, about 3200 mg, about 3225 mg, about 3250 mg, about 3275 mg, about 3300 mg, about 3325 mg, about 3350 mg, about 3375 mg, about 3400 mg, about 3425 mg, about 3450 mg, about 3475 mg, about 3500 mg, about 3525 mg, about 3550 mg, about 3575 mg, about 3600 mg, about 3625 mg, about 3650 mg, about 3675 mg, about 3700 mg, about 3725 mg, about 3750 mg, about 3775 mg, about 3800 mg, about 3825 mg, about 3850 mg, about 3875 mg, about 3900 mg, about 3925 mg, about 3950 mg, about 3975 mg, about 4000 mg, about 4025 mg, about 4050 mg, about 4075 mg, about 4100 mg, about 4125 mg, about 4150 mg, about 4175 mg, about 4200 mg, about 4225 mg, about 4250 mg, about 4275 mg, about 4300 mg, about 4325 mg, about 4350 mg, about 4375 mg, about 4400 mg, about 4425 mg, about 4450 mg, about 4475 mg, about 4500 mg, about 4525 mg, about 4550 mg, about 4575 mg, about 4600 mg, about 4625 mg, about 4650 mg, about 4675 mg, about 4700 mg, about 4725 mg, about 4750 mg, about 4775 mg, about 4800 mg, about 4825 mg, about 4850 mg, about 4875 mg, about 4900 mg, about 4925 mg, about 4950 mg, about 4975 mg, about 5000 mg, about 5025 mg, about 5050 mg, about 5075 mg, about 5100 mg, about 5125 mg, about 5150 mg, about 5175 mg, about 5200 mg, about 5225 mg, about 5250 mg, about 5275 mg, about 5300 mg, about 5325 mg, about 5350 mg, about 5375 mg, about 5400 mg, about 5425 mg, about 5450 mg, about 5475 mg, about 5500 mg, about 5525 mg, about 5550 mg, about 5575 mg, about 5600 mg, about 5625 mg, about 5650 mg, about 5675 mg, about 5700 mg, about 5725 mg, about 5750 mg, about 5775 mg, about 5800 mg, about 5825 mg, about 5850 mg, about 5875 mg, about 5900 mg, about 5925 mg, about 5950 mg, about 5975 mg, about 6000 mg, about 6025 mg, about 6050 mg, about 6075 mg, about 6100 mg, about 6125 mg, about 6150 mg, about 6175 mg, about 6200 mg, about 6225 mg, about 6250 mg, about 6275 mg, about 6300 mg, about 6325 mg, about 6350 mg, about 6375 mg, about 6400 mg, about 6425 mg, about 6450 mg, about 6475 mg, about 6500 mg, about 6525 mg, about 6550 mg, about 6575 mg, about 6600 mg, about 6625 mg, about 6650 mg, about 6675 mg, about 6700 mg, about 6725 mg, about 6750 mg, about 6775 mg, about 6800 mg, about 6825 mg, about 6850 mg, about 6875 mg, about 6900 mg, about 6925 mg, about 6950 mg, about 6975 mg, about 7000 mg, about 7025 mg, about 7050 mg, about 7075 mg, about 7100 mg, about 7125 mg, about 7150 mg, about 7175 mg, about 7200 mg, about 7225 mg, about 7250 mg, about 7275 mg, about 7300 mg, about 7325 mg, about 7350 mg, about 7375 mg, about 7400 mg, about 7425 mg, about 7450 mg, about 7475 mg, about 7500 mg, about 7525 mg, about 7550 mg, about 7575 mg, about 7600 mg, about 7625 mg, about 7650 mg, about 7675 mg, about 7700 mg, about 7725 mg, about 7750 mg, about 7775 mg, about 7800 mg, about 7825 mg, about 7850 mg, about 7875 mg, about 7900 mg, about 7925 mg, about 7950 mg, about 7975 mg, about 8000 mg, about 8025 mg, about 8050 mg, about 8075 mg, about 8100 mg, about 8125 mg, about 8150 mg, about 8175 mg, about 8200 mg, about 8225 mg, about 8250 mg, about 8275 mg, about 8300 mg, about 8325 mg, about 8350 mg, about 8375 mg, about 8400 mg, about 8425 mg, about 8450 mg, about 8475 mg, about 8500 mg, about 8525 mg,

4

about 8550 mg, about 8575 mg, about 8600 mg, about 8625 mg, about 8650 mg, about 8675 mg, about 8700 mg, about 8725 mg, about 8750 mg, about 8775 mg, about 8800 mg, about 8825 mg, about 8850 mg, about 8875 mg, about 8900 mg, about 8925 mg, about 8950 mg, about 8975 mg, about 9000 mg, about 9025 mg, about 9050 mg, about 9075 mg, about 9100 mg, about 9125 mg, about 9150 mg, about 9175 mg, about 9200 mg, about 9225 mg, about 9250 mg, about 9275 mg, about 9300 mg, about 9325 mg, about 9350 mg, about 9375 mg, about 9400 mg, about 9425 mg, about 9450 mg, about 9475 mg, about 9500 mg, about 9525 mg, about 9550 mg, about 9575 mg, about 9600 mg, about 9625 mg, about 9650 mg, about 9675 mg, about 9700 mg, about 9725 mg, about 9750 mg, about 9775 mg, about 9800 mg, about 9825 mg, about 9850 mg, about 9875 mg, about 9900 mg, about 9925 mg, about 9950 mg, about 9975 mg, or about 10,000 mg.

In one embodiment, a composition for use in methods of the invention comprises eicosapentaenoic acid, or a pharmaceutically acceptable ester, derivative, conjugate or salt thereof, or mixtures of any of the foregoing, collectively referred to herein as "EPA." The term "pharmaceutically acceptable" in the present context means that the substance in question does not produce unacceptable toxicity to the subject or interaction with other components of the composition.

In another embodiment, the EPA comprises an eicosapentaenoic acid ester. In another embodiment, the EPA comprises a C₁-C₅ alkyl ester of eicosapentaenoic acid. In another embodiment, the EPA comprises eicosapentaenoic acid ethyl ester, eicosapentaenoic acid methyl ester, eicosapentaenoic acid propyl ester, or eicosapentaenoic acid butyl ester.

In another embodiment, the EPA is in the form of ethyl-EPA, lithium EPA, mono-, di- or triglyceride EPA or any other ester or salt of EPA, or the free acid form of EPA. The EPA may also be in the form of a 2-substituted derivative or other derivative which slows down its rate of oxidation but does not otherwise change its biological action to any substantial degree.

In another embodiment, EPA is present in a composition useful in accordance with methods of the invention in an amount of about 50 mg to about 5000 mg, about 75 mg to about 2500 mg, or about 100 mg to about 1000 mg, for example about 75 mg, about 100 mg, about 125 mg, about 150 mg, about 175 mg, about 200 mg, about 225 mg, about 250 mg, about 275 mg, about 300 mg, about 325 mg, about 350 mg, about 375 mg, about 400 mg, about 425 mg, about 450 mg, about 475 mg, about 500 mg, about 525 mg, about 550 mg, about 575 mg, about 600 mg, about 625 mg, about 650 mg, about 675 mg, about 700 mg, about 725 mg, about 750 mg, about 775 mg, about 800 mg, about 825 mg, about 850 mg, about 875 mg, about 900 mg, about 925 mg, about 950 mg, about 975 mg, about 1000 mg, about 1025 mg, about 1050 mg, about 1075 mg, about 1100 mg, about 1125 mg, about 1150 mg, about 1175 mg, about 1200 mg, about 1225 mg, about 1250 mg, about 1275 mg, about 1300 mg, about 1325 mg, about 1350 mg, about 1375 mg, about 1400 mg, about 1425 mg, about 1450 mg, about 1475 mg, about 1500 mg, about 1525 mg, about 1550 mg, about 1575 mg, about 1600 mg, about 1625 mg, about 1650 mg, about 1675 mg, about 1700 mg, about 1725 mg, about 1750 mg, about 1775 mg, about 1800 mg, about 1825 mg, about 1850 mg, about 1875 mg, about 1900 mg, about 1925 mg, about 1950 mg, about 1975 mg, about 2000 mg, about 2025 mg, about 2050 mg, about 2075 mg, about 2100 mg, about 2125 mg, about 2150 mg, about 2175 mg, about 2200 mg, about 2225

US 10,568,861 B1

5

mg, about 2250 mg, about 2275 mg, about 2300 mg, about 2325 mg, about 2350 mg, about 2375 mg, about 2400 mg, about 2425 mg, about 2450 mg, about 2475 mg, about 2500 mg, about 2525 mg, about 2550 mg, about 2575 mg, about 2600 mg, about 2625 mg, about 2650 mg, about 2675 mg, about 2700 mg, about 2725 mg, about 2750 mg, about 2775 mg, about 2800 mg, about 2825 mg, about 2850 mg, about 2875 mg, about 2900 mg, about 2925 mg, about 2950 mg, about 2975 mg, about 3000 mg, about 3025 mg, about 3050 mg, about 3075 mg, about 3100 mg, about 3125 mg, about 3150 mg, about 3175 mg, about 3200 mg, about 3225 mg, about 3250 mg, about 3275 mg, about 3300 mg, about 3325 mg, about 3350 mg, about 3375 mg, about 3400 mg, about 3425 mg, about 3450 mg, about 3475 mg, about 3500 mg, about 3525 mg, about 3550 mg, about 3575 mg, about 3600 mg, about 3625 mg, about 3650 mg, about 3675 mg, about 3700 mg, about 3725 mg, about 3750 mg, about 3775 mg, about 3800 mg, about 3825 mg, about 3850 mg, about 3875 mg, about 3900 mg, about 3925 mg, about 3950 mg, about 3975 mg, about 4000 mg, about 4025 mg, about 4050 mg, about 4075 mg, about 4100 mg, about 4125 mg, about 4150 mg, about 4175 mg, about 4200 mg, about 4225 mg, about 4250 mg, about 4275 mg, about 4300 mg, about 4325 mg, about 4350 mg, about 4375 mg, about 4400 mg, about 4425 mg, about 4450 mg, about 4475 mg, about 4500 mg, about 4525 mg, about 4550 mg, about 4575 mg, about 4600 mg, about 4625 mg, about 4650 mg, about 4675 mg, about 4700 mg, about 4725 mg, about 4750 mg, about 4775 mg, about 4800 mg, about 4825 mg, about 4850 mg, about 4875 mg, about 4900 mg, about 4925 mg, about 4950 mg, about 4975 mg, or about 5000 mg.

In another embodiment, a composition useful in accordance with the invention contains not more than about 10%, not more than about 9%, not more than about 8%, not more than about 7%, not more than about 6%, not more than about 5%, not more than about 4%, not more than about 3%, not more than about 2%, not more than about 1%, or not more than about 0.5%, by weight, docosahexaenoic acid (DHA), if any. In another embodiment, a composition of the invention contains substantially no docosahexaenoic acid. In still another embodiment, a composition useful in the present invention contains no docosahexaenoic acid and/or derivative thereof.

In another embodiment, EPA comprises at least 70%, at least 80%, at least 90%, at least 95%, at least 96%, at least 97%, at least 98%, at least 99%, or 100%, by weight, of all fatty acids present in a composition that is useful in methods of the present invention.

In some embodiments, the composition comprises at least 96% by weight of eicosapentaenoic acid ethyl ester and less than about 2% by weight of a preservative. In some embodiments, the preservative is a tocopherol such as all-racemic α -tocopherol.

In another embodiment, a composition useful in accordance with methods of the invention contains less than 10%, less than 9%, less than 8%, less than 7%, less than 6%, less than 5%, less than 4%, less than 3%, less than 2%, less than 1%, less than 0.5% or less than 0.25%, by weight of the total composition or by weight of the total fatty acid content, of any fatty acid other than EPA. Illustrative examples of a "fatty acid other than EPA" include linolenic acid (LA), arachidonic acid (AA), docosahexaenoic acid (DHA), alpha-linolenic acid (ALA), stearadonic acid (STA), eicosatrienoic acid (ETA) and/or docosapentaenoic acid (DPA). In another embodiment, a composition useful in accordance with methods of the invention contains about 0.1% to about 4%, about

6

0.5% to about 3%, or about 1% to about 2%, by weight, of total fatty acids other than EPA and/or DHA.

In another embodiment, a composition useful in accordance with the invention has one or more of the following features: (a) eicosapentaenoic acid ethyl ester represents at least about 96%, at least about 97%, or at least about 98%, by weight, of all fatty acids present in the composition; (b) the composition contains not more than about 4%, not more than about 3%, or not more than about 2%, by weight, of total fatty acids other than eicosapentaenoic acid ethyl ester; (c) the composition contains not more than about 0.6%, not more than about 0.5%, or not more than about 0.4% of any individual fatty acid other than eicosapentaenoic acid ethyl ester; (d) the composition has a refractive index (20° C.) of about 1 to about 2, about 1.2 to about 1.8 or about 1.4 to about 1.5; (e) the composition has a specific gravity (20° C.) of about 0.8 to about 1.0, about 0.85 to about 0.95 or about 0.9 to about 0.92; (f) the composition contains not more than about 20 ppm, not more than about 15 ppm or not more than about 10 ppm heavy metals, (f) the composition contains not more than about 5 ppm, not more than about 4 ppm, not more than about 3 ppm, or not more than about 2 ppm arsenic, and/or (g) the composition has a peroxide value of not more than about 5 meq/kg, not more than about 4 meq/kg, not more than about 3 meq/kg, or not more than about 2 meq/kg.

In another embodiment, compositions useful in accordance with methods of the invention are orally deliverable. The terms "orally deliverable" or "oral administration" herein include any form of delivery of a therapeutic agent or a composition thereof to a subject wherein the agent or composition is placed in the mouth of the subject, whether or not the agent or composition is swallowed. Thus "oral administration" includes buccal and sublingual as well as esophageal administration. In one embodiment, the composition is present in a capsule, for example a soft gelatin capsule.

A composition for use in accordance with the invention can be formulated as one or more dosage units. The terms "dose unit" and "dosage unit" herein refer to a portion of a pharmaceutical composition that contains an amount of a therapeutic agent suitable for a single administration to provide a therapeutic effect. Such dosage units may be administered one to a plurality (i.e. 1 to about 10, 1 to 8, 1 to 6, 1 to 4 or 1 to 2) of times per day, or as many times as needed to elicit a therapeutic response.

In one embodiment, compositions of the invention, upon storage in a closed container maintained at room temperature, refrigerated (e.g. about 5 to about 5-10° C.) temperature, or frozen for a period of about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, or 12 months, exhibit at least about 90%, at least about 95%, at least about 97.5%, or at least about 99% of the active ingredient(s) originally present therein.

Therapeutic Methods

In one embodiment, the invention provides a method for treatment and/or prevention of cardiovascular-related disease and disorders. The term "cardiovascular-related disease and disorders" herein refers to any disease or disorder of the heart or blood vessels (i.e. arteries and veins) or any symptom thereof. Non-limiting examples of cardiovascular-related disease and disorders include hypertriglyceridemia, hypercholesterolemia, mixed dyslipidemia, coronary heart disease, vascular disease, stroke, atherosclerosis, arrhythmia, hypertension, myocardial infarction, and other cardiovascular events.

The term "treatment" in relation a given disease or disorder, includes, but is not limited to, inhibiting the disease

US 10,568,861 B1

7

or disorder, for example, arresting the development of the disease or disorder; relieving the disease or disorder, for example, causing regression of the disease or disorder; or relieving a condition caused by or resulting from the disease or disorder, for example, relieving, preventing or treating symptoms of the disease or disorder. The term "prevention" in relation to a given disease or disorder means: preventing the onset of disease development if none had occurred, preventing the disease or disorder from occurring in a subject that may be predisposed to the disorder or disease but has not yet been diagnosed as having the disorder or disease, and/or preventing further disease/disorder development if already present.

In various embodiments, the present invention provides methods of reducing a risk of a cardiovascular event in a subject on statin therapy. In some embodiments, the method comprises (a) identifying a subject on statin therapy and having a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL, wherein said subject has established cardiovascular disease or has a high risk of developing cardiovascular disease; and (b) administering to the subject a pharmaceutical composition comprising about 1 g to about 4 g of eicosapentaenoic acid ethyl ester per day, wherein the composition contains substantially no docosa-hexaenoic acid.

In some embodiments, the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL, for example 135 mg/dL to 500 mg/dL, 150 mg/dL to 500 mg/dL, or 200 mg/dL to <500 mg/dL. In some embodiments, the subject or subject group has a baseline triglyceride level (or median baseline triglyceride level in the case of a subject group), fed or fasting, of about 135 mg/dL, about 140 mg/dL, about 145 mg/dL, about 150 mg/dL, about 155 mg/dL, about 160 mg/dL, about 165 mg/dL, about 170 mg/dL, about 175 mg/dL, about 180 mg/dL, about 185 mg/dL, about 190 mg/dL, about 195 mg/dL, about 200 mg/dL, about 205 mg/dL, about 210 mg/dL, about 215 mg/dL, about 220 mg/dL, about 225 mg/dL, about 230 mg/dL, about 235 mg/dL, about 240 mg/dL, about 245 mg/dL, about 250 mg/dL, about 255 mg/dL, about 260 mg/dL, about 265 mg/dL, about 270 mg/dL, about 275 mg/dL, about 280 mg/dL, about 285 mg/dL, about 290 mg/dL, about 295 mg/dL, about 300 mg/dL, about 305 mg/dL, about 310 mg/dL, about 315 mg/dL, about 320 mg/dL, about 325 mg/dL, about 330 mg/dL, about 335 mg/dL, about 340 mg/dL, about 345 mg/dL, about 350 mg/dL, about 355 mg/dL, about 360 mg/dL, about 365 mg/dL, about 370 mg/dL, about 375 mg/dL, about 380 mg/dL, about 385 mg/dL, about 390 mg/dL, about 395 mg/dL, about 400 mg/dL, about 405 mg/dL, about 410 mg/dL, about 415 mg/dL, about 420 mg/dL, about 425 mg/dL, about 430 mg/dL, about 435 mg/dL, about 440 mg/dL, about 445 mg/dL, about 450 mg/dL, about 455 mg/dL, about 460 mg/dL, about 465 mg/dL, about 470 mg/dL, about 475 mg/dL, about 480 mg/dL, about 485 mg/dL, about 490 mg/dL, about 495 mg/dL, or about 500 mg/dL.

In some embodiments, the subject or subject group is also on stable therapy with a statin (with or without ezetimibe). In some embodiments, the subject or subject group also has established cardiovascular disease, or is at high risk for establishing cardiovascular disease. In some embodiments, the subject's statin therapy includes administration of one or more statins. For example and without limitation, the subject's statin therapy may include one or more of: atorvastatin, fluvastatin, lovastatin, pitavastatin, pravastatin, rosuvastatin, and simvastatin. In some embodiments, the subject

8

is additionally administered one or more of: amlodipine, ezetimibe, niacin, and sitagliptin. In some embodiments, the subject's statin therapy includes administration of a statin and ezetimibe. In some embodiments, the subject's statin therapy includes administration of a statin without ezetimibe.

In some embodiments, the subject's statin therapy does not include administration of 200 mg or more per day of niacin and/or fibrates. In some embodiments, the subject is not on concomitant omega-3 fatty acid therapy (e.g., is not being administered or co-administered a prescription and/or over-the-counter composition comprising an omega-3 fatty acid active agent). In some embodiments, the subject is not administered or does not ingest a dietary supplement comprising an omega-3 fatty acid.

In some embodiments, the subject has established cardiovascular disease ("CV disease" or "CVD"). The status of a subject as having CV disease can be determined by any suitable method known to those skilled in the art. In some embodiments, a subject is identified as having established CV disease by the presence of any one of: documented coronary artery disease, documented cerebrovascular disease, documented carotid disease, documented peripheral arterial disease, or combinations thereof. In some embodiments, a subject is identified as having CV disease if the subject is at least 45 years old and: (a) has one or more stenosis of greater than 50% in two major epicardial coronary arteries; (b) has had a documented prior MI; (c) has been hospitalized for high-risk NSTEMI ACS with objective evidence of ischemia (e.g., ST-segment deviation and/or biomarker positivity); (d) has a documented prior ischemic stroke; (e) has symptomatic artery disease with at least 50% carotid arterial stenosis; (f) has asymptomatic carotid artery disease with at least 70% carotid arterial stenosis per angiography or duplex ultrasound; (g) has an ankle-brachial index ("ABI") of less than 0.9 with symptoms of intermittent claudication; and/or (h) has a history of aorto-iliac or peripheral arterial intervention (catheter-based or surgical).

In some embodiments, the subject or subject group being treated in accordance with methods of the invention has a high risk for developing CV disease. For example and without limitation, a subject or subject group has a high risk for developing CV disease if the subject or subject in a subject group is age 50 or older, has diabetes mellitus (Type 1 or Type 2), and at least one of: (a) is a male age 55 or older or a female age 65 or older; (b) is a cigarette smoker or was a cigarette smoker who stopped less than 3 months prior; (c) has hypertension (e.g., a blood pressure of 140 mmHg systolic or higher, or greater than 90 mmHg diastolic); (d) has an HDL-C level of ≤ 40 mg/dL for men or ≤ 50 mg/dL for women; (e) has an hs-CRP level of > 3.0 mg/L; (f) has renal dysfunction (e.g., a creatinine clearance ("CrCL") of greater than 30 mL/min and less than 60 mL/min); (g) has retinopathy (e.g., defined as any of: non-proliferative retinopathy, preproliferative retinopathy, proliferative retinopathy, maculopathy, advanced diabetic eye disease, or history of photocoagulation); (h) has microalbuminuria (e.g., a positive micral or other strip test, an albumin/creatinine ratio of ≥ 2.5 mg/mmol, or an albumin excretion rate on timed collection of ≥ 20 mg/min all on at least two successive occasions); (i) has macroalbuminuria (e.g., albumix or other dip stick evidence of gross proteinuria, an albumin/creatinine ratio of ≥ 25 mg/mmol, or an albumin excretion rate on timed collection of ≥ 200 mg/min all on at least two successive occasions); and/or (j) has an ankle-brachial index of < 0.9 without symptoms of intermittent claudication.

US 10,568,861 B1

9

In some embodiments, the subject's baseline lipid profile is measured or determined prior to administering the pharmaceutical composition to the subject. Lipid profile characteristics can be determined by any suitable method known to those skilled in the art including, for example, by testing a fasting or non-fasting blood sample obtained from the subject using standard blood lipid profile assays. In some embodiments, the subject has one or more of: a baseline non-HDL-C value of about 200 mg/dL to about 300 mg/dL; a baseline total cholesterol value of about 250 mg/dL to about 300 mg/dL; a baseline VLDL-C value of about 140 mg/dL to about 200 mg/dL; a baseline HDL-C value of about 10 to about 30 mg/dL; and/or a baseline LDL-C value of about 40 to about 100 mg/dL.

In some embodiments, the cardiovascular event for which risk is reduced is one or more of: cardiovascular death; nonfatal myocardial infarction; nonfatal stroke; coronary revascularization; unstable angina (e.g., unstable angina determined to be caused by myocardial ischemia by, for example, invasive or non-invasive testing, and requiring hospitalization); cardiac arrest; peripheral cardiovascular disease requiring intervention, angioplasty, bypass surgery or aneurysm repair; death; and onset of new congestive heart failure.

In some embodiments, the subject is administered about 1 g to about 4 g of the pharmaceutical composition per day for about 4 months, about 1 year, about 2 years, about 3 years, about 4 years, about 5 years, or more than about 5 years. Thereafter, in some embodiments the subject exhibits one or more of

- (a) reduced triglyceride levels compared to baseline;
- (b) reduced Apo B levels compared to baseline;
- (c) increased HDL-C levels compared to baseline;
- (d) no increase in LDL-C levels compared to baseline;
- (e) a reduction in LDL-C levels compared to baseline;
- (f) a reduction in non-HDL-C levels compared to baseline;
- (g) a reduction in VLDL levels compared to baseline;
- (h) a reduction in total cholesterol levels compared to baseline;
- (i) a reduction in high sensitivity C-reactive protein (hs-CRP) levels compared to baseline; and/or
- (j) a reduction in high sensitivity troponin (hsTnT) levels compared to baseline.

In some embodiments, the subject exhibits one or more of: (a) a reduction in triglyceride level of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 55% as compared to baseline;

(b) a less than 30% increase, less than 20% increase, less than 10% increase, less than 5% increase or no increase in non-HDL-C levels or a reduction in non-HDL-C levels of at least about 1%, at least about 3%, at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% as compared to baseline;

(c) an increase in HDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, or at least about 50% as compared to baseline; and/or

(d) a less than 30% increase, less than 20% increase, less than 10% increase, less than 5% increase or no increase in LDL-C levels or a reduction in LDL-C levels of at least about 5%, at least about 10%, at least about 15%, at least

10

about 20%, at least about 25%, at least about 30%, at least about 35%, at least about 40%, at least about 45%, at least about 50%, or at least about 55% as compared to baseline.

In one embodiment, the subject or subject group being treated has a baseline EPA blood level on a (mol %) basis of less than 2.6, less than 2.5, less than 2.4, less than 2.3, less than 2.2, less than 2.1, less than 2, less than 1.9, less than 1.8, less than 1.7, less than 1.6, less than 1.5, less than 1.4, less than 1.3, less than 1.2, less than 1.1 or less than 1.

In another embodiment, the subject or subject group being treated has a baseline triglyceride level (or median baseline triglyceride level in the case of a subject group), fed or fasting, of about 135 mg/dL to about 100 mg/dL. In some embodiments, the subject or subject group being treated in accordance with methods of the invention is on stable therapy with a statin (with or without ezetimibe). As used herein, the phrase "on stable therapy with a statin" means that the subject or subject group has been on the same daily dose of the same statin for at least 28 days and, if applicable, the same daily dose of ezetimibe for at least 28 days. In some embodiments, the subject or subject group on stable statin therapy has an LDL-C level of about 40 mg/dL to about 100 mg/dL.

In some embodiments, safety laboratory tests of subject blood samples include one or more of: hematology with complete blood count ("CBC"), including RBC, hemoglobin (Hgb), hematocrit (Hct), white cell blood count (WBC), white cell differential, and platelet count; and biochemistry panel including total protein, albumin, alkaline phosphatase, alanine aminotransferase (ALT/SGPT), aspartate aminotransferase (AST/SGOT), total bilirubin, glucose, calcium, electrolytes, (sodium, potassium, chloride), blood urea nitrogen (BUN), serum creatinine, uric acid, creatine kinase, and HbA_{1c}.

In some embodiments, a fasting lipid panel associated with a subject includes TG, TC, LDL-C, HDL-C, non-HDL-C, and VLDL-C. In some embodiments, LDL-C is calculated using the Friedewald equation, or is measured by preparative ultracentrifugation (Beta Quant) if the subject's triglyceride level is greater than 400 mg/dL. In some embodiments, LDL-C is measured by ultracentrifugation (Beta Quant) at randomization and again after about one year after randomization.

In some embodiments, a biomarker assay associated with blood obtained from a subject includes hs-CRP, Apo B and hsTnT.

In some embodiments, a medical history associated with a subject includes family history, details regarding all illnesses and allergies including, for example, date(s) of onset, current status of condition(s), and smoking and alcohol use.

In some embodiments, demographic information associated with a subject includes day, month and year of birth, race, and gender.

In some embodiments, vital signs associated with a subject include systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature (e.g., oral body temperature).

In some embodiments, a physical examination of a subject includes assessments of the subject's general appearance, skin, head, neck, heart, lung, abdomen, extremities, and neuromusculature.

In some embodiments, the subject's height and weight are measured. In some embodiments, the subject's weight is recorded with the subject wearing indoor clothing, with shoes removed, and with the subject's bladder empty.

In some embodiments, a waist measurement associated with the subject is measured. In some embodiments, the

US 10,568,861 B1

11

waist measurement is determined with a tape measure at the top of the subject's hip bone.

In some embodiments, an electrocardiogram associated with the subject is obtained. In some embodiments, an ECG is obtained every year during the treatment/follow-up portion of the study. In some embodiments, the ECG is a 12-lead ECG. In some embodiments, the ECG is analyzed for detection of silent MI.

In some embodiments, subjects randomly assigned to the treatment group receive 4 g per day of a composition comprising at least 96% by weight of eicosapentaenoic acid ethyl ester. In some embodiments, the composition is encapsulated in a gelatin capsule. In some embodiments, subjects in this treatment group continue to take 4 g per day of the composition for about 1 year, about 2 years, about 3 years, about 4 years, about 4.75 years, about 5 years, about 6 years, about 7 years, about 8 years, about 9 years, about 10 years, or more than about 10 years. In some embodiments, a median treatment duration is planned to be about 4 years.

In some embodiments, the present invention provides a method of reducing a risk of cardiovascular events in a subject. In some embodiments, the method comprises administering to the subject a composition comprising at least 96% by weight of eicosapentaenoic acid ethyl ester. In some embodiments, the subject is administered about 1 g to about 4 g of the composition per day.

In some embodiments, the reduced risk of CV events is indicated or determined by comparing an amount of time (e.g., an average amount of time) associated with a subject or subject group from first dosing to a first CV event selected from the group consisting of: CV death, nonfatal MI, nonfatal stroke, coronary revascularization, and hospitalization (e.g., emergent hospitalization) for unstable angina determined to be caused by myocardial ischemia (e.g., by invasive or non-invasive testing), to an amount of time (e.g., an average amount of time) associated with a placebo or untreated subject or group of subjects from first dosing with a placebo to a first CV event selected from the group consisting of: CV death, nonfatal MI, nonfatal stroke, coronary revascularization, and hospitalization (e.g., emergent hospitalization) for unstable angina determined to be caused by myocardial ischemia (e.g., by invasive or non-invasive testing), wherein said placebo does not include eicosapentaenoic acid ethyl ester. In some embodiments, the amount of time associated with the subject or group of subjects are compared to the amount of time associated with the placebo or untreated subject or group of subjects are compared using a log-rank test. In some embodiments, the log-rank test includes one or more stratification factors such as CV Risk Category, use of ezetimibe, and/or geographical region.

In some embodiments, the present invention provides a method of reducing risk of CV death in a subject on stable statin therapy and having CV disease or at high risk for developing CV disease, comprising administering to the subject a composition as disclosed herein.

In another embodiment, the present invention provides a method of reducing risk of recurrent nonfatal myocardial infarction (including silent MI) in a subject on stable statin therapy and having CV disease or at high risk for developing CV disease, comprising administering to the patient one or more compositions as disclosed herein.

In some embodiments, the present invention provides a method of reducing risk of nonfatal stroke in a subject on stable statin therapy and having CV disease or at high risk for developing CV disease, comprising administering to the subject a composition as disclosed herein.

12

In some embodiments, the present invention provides a method of reducing risk of coronary revascularization in a subject on stable statin therapy and having CV disease or at high risk for developing CV disease, comprising administering to the subject a composition as disclosed herein.

In some embodiments, the present invention provides a method of reducing risk of developing unstable angina caused by myocardial ischemia in a subject on stable statin therapy and having CV disease or at high risk for developing CV disease, comprising administering to the subject a composition as disclosed herein.

In another embodiment, any of the methods disclosed herein are used in treatment or prevention of a subject or subjects that consume a traditional Western diet. In one embodiment, the methods of the invention include a step of identifying a subject as a Western diet consumer or prudent diet consumer and then treating the subject if the subject is deemed a Western diet consumer. The term "Western diet" herein refers generally to a typical diet consisting of, by percentage of total calories, about 45% to about 50% carbohydrate, about 35% to about 40% fat, and about 10% to about 15% protein. A Western diet may alternately or additionally be characterized by relatively high intakes of red and processed meats, sweets, refined grains, and desserts, for example more than 50%, more than 60% or more or 70% of total calories come from these sources.

In another embodiment, a composition as described herein is administered to a subject once or twice per day. In another embodiment, 1, 2, 3 or 4 capsules, each containing about 1 g of a composition as described herein, are administered to a subject daily. In another embodiment, 1 or 2 capsules, each containing about 1 g of a composition as described herein, are administered to the subject in the morning, for example between about 5 am and about 11 am, and 1 or 2 capsules, each containing about 1 g of a composition as described herein, are administered to the subject in the evening, for example between about 5 pm and about 11 pm.

In some embodiments, the risk of a cardiovascular event in a subject is reduced compared to a control population. In some embodiments, a plurality of control subjects to a control population, wherein each control subject is on stable statin therapy, has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL, and has established cardiovascular disease or a high risk of developing cardiovascular disease, and wherein the control subjects are not administered the pharmaceutical composition comprising about 1 g to about 4 g of eicosapentaenoic acid ethyl ester per day.

In some embodiments, a first time interval beginning at (a) an initial administration of a composition as disclosed herein to the subject to (b) a first cardiovascular event of the subject is greater than or substantially greater than a first control time interval beginning at (a') initial administration of a placebo to the control subjects to (b') a first cardiovascular event in the control subjects. In some embodiments, the first cardiovascular event of the subject is a major cardiovascular event selected from the group consisting of: cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, coronary revascularization, and unstable angina caused by myocardial ischemia. In some embodiments, the first cardiovascular event of the control subjects is a major cardiovascular event selected from the group consisting of: cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, coronary revascularization, and unstable angina caused by myocardial ischemia. In some embodiments, the first cardiovascular event of the subject and the first cardio-

US 10,568,861 B1

13

vascular event of the control subjects is any of: death (from any cause), nonfatal myocardial infarction, or nonfatal stroke. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any of: death from a cardiovascular cause, nonfatal myocardial infarction, coronary revascularization, unstable angina, peripheral cardiovascular disease, or cardiac arrhythmia requiring hospitalization. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any of: death from a cardiovascular cause and nonfatal myocardial infarction. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is death (from any cause). In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any of: fatal myocardial infarction and nonfatal myocardial infarction (optionally including silent MI). In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is coronary revascularization. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is hospitalization (e.g. emergent hospitalization) for unstable angina (optionally unstable angina caused by myocardial ischemia). In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any one of: fatal stroke or nonfatal stroke. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any one of: new coronary heart failure, new coronary heart failure leading to hospitalization, transient ischemic attack, amputation for coronary vascular disease, and carotid revascularization. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is any one of: elective coronary revascularization and emergent coronary revascularization. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is an onset of diabetes. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is cardiac arrhythmia requiring hospitalization. In some embodiments, the first cardiovascular event of the subject and the first cardiovascular event of the control subjects is cardiac arrest.

In some embodiments, a second time interval beginning at (a) an initial administration of the pharmaceutical composition to the subject to (c) a second cardiovascular event of the subject is greater than or substantially greater than a second control time interval beginning at (a') initial administration of a placebo to the control subjects to (c') a second cardiovascular event in the control subjects. In some embodiments, the second cardiovascular event of the subject and the second cardiovascular event of the control subjects is a major cardiovascular event selected from the group consisting of: cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, coronary revascularization, and unstable angina caused by myocardial ischemia.

In some embodiments, the subject has diabetes mellitus and the control subjects each have diabetes mellitus. In some embodiments, the subject has metabolic syndrome and the control subjects each have metabolic syndrome.

14

In some embodiments, the subject exhibits one or more of (a) reduced triglyceride levels compared to the control population; (b) reduced Apo B levels compared to the control population; (c) increased HDL-C levels compared to the control population; (d) no increase in LDL-C levels compared to the control population; (e) a reduction in LDL-C levels compared to the control population; (f) a reduction in non-HDL-C levels compared to the control population; (g) a reduction in VLDL levels compared to the control population; (h) a reduction in total cholesterol levels compared to the control population; (i) a reduction in high sensitivity C-reactive protein (hs-CRP) levels compared to the control population; and/or (j) a reduction in high sensitivity troponin (hsTnT) levels compared to the control population.

In some embodiments, the subject's weight after administration of the composition is less than a baseline weight determined before administration of the composition. In some embodiments, the subject's waist circumference after administration of the composition is less than a baseline waist circumference determined before administration of the composition.

In methods of the present invention in which a time interval is determined or assessed, the time interval may be for example an average, a median, or a mean time interval. For example, in embodiments wherein a first control time interval is associated with a plurality of control subjects, the first control time interval is an average, a median, or a mean of a plurality of first control time intervals associated with each control subject. Similarly, in embodiments wherein a second control time interval is associated with a plurality of control subjects, the second control time interval is an average, a median, or a mean of a plurality of second control time intervals associated with each control subject.

In some embodiments, the reduced risk of cardiovascular events is expressed as a difference in incident rates between a study group and a control population. In some embodiments, the subjects in the study group experience a first major cardiovascular event after an initial administration of a composition as disclosed herein at a first incidence rate which is less than a second incidence rate, wherein the second incidence rate is associated with the rate of cardiovascular events in the subjects in the control population. In some embodiments, the first major cardiovascular event is any one of: cardiovascular death, nonfatal myocardial infarction, nonfatal stroke, coronary revascularization, and hospitalization for unstable angina (optionally determined to be caused by myocardial ischemia). In some embodiments, the first and second incidence rates are determined for a time period beginning on the date of the initial administration and ending about 4 months, about 1 year, about 2 years, about 3 years, about 4 years, or about 5 years after the date of initial administration.

In another embodiment, the invention provides use of any composition described herein for treating hypertriglyceridemia in a subject in need thereof, comprising: providing a subject having a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL and administering to the subject a pharmaceutical composition as described herein. In one embodiment, the composition comprises about 1 g to about 4 g of eicosapentaenoic acid ethyl ester, wherein the composition contains substantially no docosahexaenoic acid.

EXAMPLES

A phase 3, multi-center, placebo-controlled randomized, double-blind, 12-week study with an open-label extension is

US 10,568,861 B1

15

performed to evaluate the efficacy and safety of AMR101 in patients with fasting triglyceride levels ≥ 150 mg/dL and < 500 mg/dL. The primary objective is, in patients at LDL-C goal while on statin therapy, with established cardiovascular disease (CVD) or at high risk for CVD, and hypertriglyceridemia (fasting triglycerides, TG, ≥ 200 mg/dL and < 500 mg/dL, determine the efficacy of AMR101 4 g daily, compared to placebo, in preventing the occurrence of a first major cardiovascular event of the composite endpoint that includes:

- cardiovascular ("CV") death;
- nonfatal myocardial infarction ("MI");
- nonfatal stroke;
- coronary revascularization; and
- unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization.

The secondary objectives of this study are the following:

To evaluate the effect of therapy on the composite of death from CV causes, nonfatal MI, coronary revascularization, unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization, nonfatal stroke, or peripheral CV disease requiring intervention, angioplasty, bypass surgery, and aneurysm repair;

To evaluate the effect of therapy on combinations of each of the clinical events listed in secondary objective #1, supra, in addition to cardiac arrhythmia requiring hospitalization, cardiac arrest, peripheral CV disease requiring intervention, angioplasty, bypass surgery, aneurysm repair, and total mortality;

To evaluate the effect of therapy on the occurrence of a second, third, fourth and fifth major cardiovascular event (e.g., occurrence of CV death, nonfatal MI, nonfatal stroke, coronary revascularization, and unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization after a first occurrence of any of same);

To evaluate the effect of therapy on the first occurrence of a major cardiovascular event in subgroups of patients including (a) those with diabetes mellitus, and (b) those with metabolic syndrome (e.g., as defined by the NCEP ATP III or future criteria as may evolve therefrom);

To evaluate the effect of therapy on new congestive heart failure ("CHF"), on new CHF as a primary cause of hospitalization, on transient ischemic attack, on amputation for CV disease, and on carotid revascularization;

To evaluate the effect of therapy on occurrence of elective coronary revascularization and emergent coronary revascularization;

To evaluate the effects of therapy on lipids, lipoproteins and inflammatory markers including triglycerides, total cholesterol, low-density lipoprotein cholesterol ("LDL-C"), high-density lipoprotein cholesterol ("HDL-C"), non-HDL-C, very low-density lipoprotein cholesterol ("VLDL-C"), apolipoprotein B ("apo B"), high-sensitivity C-reactive protein ("hs-CRP"), and high-sensitivity troponin ("hsTnT") as follows:

- Evaluation of the effect of therapy on each marker;

- Evaluation of the effect of the baseline value of each marker on therapy effects; and

- Evaluation of the effect of therapy for preventing clinical events as defined above among all patients in the study and in sub-groups such as patients with diabetes mellitus and patients with substantial on-treatment changes of any of the markers;

16

To evaluate the effect of therapy on new onset diabetes; and

To explore the effect of therapy on weight and waist circumference.

Study Population

The population for this study is men and women ≥ 45 years of age with established CVD, or men and women ≥ 50 years of age with diabetes in combination with one additional risk factor for CVD. In addition, all patients will have atherogenic dyslipidemia defined as on treatment for hypercholesterolemia (but at treatment goal for LDL-C, by treatment with a statin) and hypertriglyceridemia. More details are listed in the inclusion criteria.

The patients will need to provide consent to participate in the study and be willing and able to comply with the protocol and the study procedures.

Study Periods

This study consists of the following study periods:

Screening Period: During the screening period, patients will be evaluated for inclusion/exclusion criteria.

At the first visit to the Research Unit (Visit 1), study procedures will be performed for evaluation of patient's eligibility in the study. At this screening visit, patients will sign an informed consent form before any study procedure is performed; the informed consent form will cover the treatment/follow-up period. Based on the evaluation from Visit 1, the following situations may occur:

Patients who are eligible for participation based on the study procedures on Visit 1 will return to the Research Unit for Visit 2 (randomization visit) to start the treatment/follow-up period. This case includes, for example, patients at Visit 1 who are on a stable dose of a statin, are planning to stay on the same statin and the same dose of the statin, and who not need to wash out any non-statin lipid-altering medications.

Patients who are not eligible for participation based on the study procedures on Visit 1 and are unlikely to become eligible in the next 28 days (for example: unlikely to stabilize statin dose, unable to wash out non-statin lipid-altering medications, etc.): these patients will be screen failed after Visit 1.

Patients not eligible for participation in the study based on the study procedures on Visit 1 may possibly become eligible in the next 28 days: these patients may return at the discretion of the investigator for a second optional screening visit (Visit 1.1) at which time the procedures needed for re-evaluation of the previously failed inclusion/exclusion criteria will be repeated. This case includes, for example, patients who are started on a statin at Visit 1, whose statin dose is changed at Visit 1, and/or needed to wash out non-statin lipid-altering medications. The following applies for these patients:

Patients with a change in the statin or statin dose on Visit 1 will need to be on a stable statin dose for at least 28 days before the lipid qualifying measurements at Visit 1.1. Other concomitant medications (antidiabetic therapy, for example) can be optimized or stabilized during this period.

Patients starting a washout at Visit 1 will have a washout period of at least 28 days (only 7 days for bile acid sequestrants) before the lipid qualifying measurements at Visit 1.1.

Patients at Visit 1 who are on a stable dose of a statin, are planning to stay on the same statin at the same dose, and who do not need any medication washout, but were asked to return for Visit 1.1 to repeat one or more of the other study procedures not related to concomitant medications

US 10,568,861 B1

17

Patients who become eligible for participation based on the additional study procedures at Visit 1.1 will return to the Research Unit for Visit 2 (randomization visit) to start the treatment/follow-up period.

At the end of the screening period, patients will need to meet all inclusion/exclusion criteria before they can be randomized. Patients who are not eligible for participation after the screening period (based on study procedures at Visit 1 and/or Visit 1.1) may return at a later date for rescreening. These patients will need to re-start with all procedures starting with Visit 1. This includes patients who need more time to stabilize one or more conditions or therapies (for example: statin, antidiabetic, antihypertensive, thyroid hormone, HIV-protease inhibitor therapy).

Treatment/Follow-Up Period: Within 42 days after the first screening visit (Visit 1) or within 60 days after the first screening visit (Visit 1) for those patients that have a second screening visit (Visit 1.1), eligible patients will enter the treatment/follow-up period. During this period, the patients will receive study drug during the planned visits at the Research Site and take the study drug while away from the Research Site.

During the visits, study procedures will be performed for evaluation of efficacy and safety. A detailed schedule of procedures is provided in Table 1.

Study Duration

The estimated study duration includes a planned 18-month enrollment period followed by a follow-up period of approximately 3.5 years in expected duration (approximately 5 years in total). Patients will be randomized at different times during the enrollment period but will all end the study at the same date (study end date). It is planned that all randomized patients will receive study medication and be followed-up until the study end date. This is an event-driven trial and patients will continue in the trial if the trial runs longer than expected, or will terminate earlier if the trial runs shorter than expected.

The total duration of the trial is based on a median 4-year follow-up period across patients. The first patient randomized would be followed for 4.75 years (the longest individual follow-up duration), and the last patient randomized would be followed for 3.25 year (the shortest individual follow-up duration).

Study Groups

At Visit 2 (Day 0), eligible study patients will be randomly assigned to the following treatment groups:

Group 1: AMR101 4 g daily (four 1000 mg capsules daily)

Group 2: placebo (four capsules daily)

The four AMR101 or placebo capsules daily will be taken as two capsules in the morning and two capsules in the evening (twice-per-day dosing regimen).

Number of Patients

This is an event-driven trial: It is expected that a minimum of 1612 primary efficacy endpoint events will be required during the study. A total of approximately 7990 patients will be entered into the study to either receive AMR101 or placebo (approximately 3995 patients per treatment group) in order to observe an estimated 1612 events that make up the primary composite endpoint for efficacy.

Number of Study Sites

Participants will be enrolled at multiple Research Sites in multiple countries.

Randomization

On Day 0, eligible patients will be randomized to one of 2 study groups using a computer-generated randomization

18

schema. Randomized treatment assignment to either AMR101 or placebo in a 1:1 ratio will be provided using the internet (IWR).

Blinding

This is a double-blind study. Patients, investigators, pharmacists and other supporting staff at the Research Sites, personnel and designees of the Sponsor, study administrators and personnel at the organization(s) and vendors supporting the study will be unaware of the randomization code (i.e., they will not know which study participants are receiving the experimental drug and which are receiving the placebo drug). The study medication AMR101 and placebo capsules will be similar in size and appearance to maintain blinding.

During the double-blind treatment/follow-up period, everyone (patients, investigators, pharmacists and other supporting staff at the Research Sites, personnel and designees of the Sponsor, study administrators and personnel at the organization(s) and vendors managing/supporting the study), with the exception of the laboratory personnel performing the analysis, will be blinded to individual results of the efficacy laboratory measurements (including lipid values). Individual results from the lipid profile may be unblinded in the event of an emergency for a patient.

Stratification

Participants will be assigned to treatment groups stratified by CV risk category, use of ezetimibe and by geographical region (Westernized, Eastern European, and Asia Pacific countries). There are two CV risk categories:

CV Risk Category 1: patients with established CVD defined in the inclusion criteria. Patients with diabetes and established CVD are included in this category.

CV Risk Category 2: patients with diabetes and at least one additional risk factor for CVD, but no established CVD.

Stratification will be recorded in the IWR at the time of enrollment. Approximately 70% of randomized patients will be in the CV Risk Category 1 and approximately 30% of randomized patients will be in the CV Risk Category 2. Enrollment with patients of a CV risk category will be stopped when the planned number of patients in that risk category is reached.

Study Population

Inclusion Criteria

Patients meeting the following criteria will be eligible to participate in the study:

Fasting TG levels of ≥ 200 mg/dL (2.26 mmol/L) and < 500 mg/dL (5.64 mmol/L).

LDL-C > 40 mg/dL (1.04 mmol/L) and ≤ 100 mg/dL (2.60 mmol/L) and on stable therapy with a statin (with or without ezetimibe) for at least 4 weeks prior to the LDL-C/TG baseline qualifying measurements for randomization

Stable therapy is defined as the same daily dose of the same statin for at least 28 days before the lipid qualification measurements (TG and LDL-C) and, if applicable, the same daily dose of ezetimibe for at least 28 days before the lipid qualification measurements (TG and LDL-C). Patients who have their statin therapy or use of ezetimibe initiated at Visit 1, or have their statin, statin dose and/or ezetimibe dose changed at Visit 1, will need to go through a stabilization period of at least 28 days since initiation/change and have their qualifying lipid measurements measured (TG and LDL-C) after the washout period (at Visit 1.1).

Statins may be administered with or without ezetimibe.

If patients qualify at the first qualification visit (Visit 1) for TG and LDL-C, and meet all other inclusion/exclusion criteria, they may be randomized at Visit 2. If patients don't qualify at the first qualifying visit (Visit 1), a second

US 10,568,861 B1

19

re-qualifying visit (Visit 1.1) is allowed. For some patients, because they need to stabilize medications and/or need to washout medications, the second re-qualifying visit (Visit 1.1) will be needed after the stabilization/washout period.

Either having established CVD (in CV Risk Category 1) or at high risk for CVD (in CV Risk Category 2). The CV risk categories are defined as follows:

CV Risk Category 1: defined as men and women ≥ 45 years of age with one or more of the following:

Documented coronary artery disease (CAD; one or more of the following primary criteria must be satisfied):

Documented multivessel CAD ($>50\%$ stenosis in at least two major epicardial coronary arteries—with or without antecedent revascularization)

Documented prior MI

Hospitalization for high-risk NSTEMI-ACS (with objective evidence of ischemia: ST-segment deviation or biomarker positivity)

Documented cerebrovascular or carotid disease (one of the following primary criteria must be satisfied):

Documented prior ischemic stroke

Symptomatic carotid artery disease with $\geq 50\%$ carotid arterial stenosis

Asymptomatic carotid artery disease with $\geq 70\%$ carotid arterial stenosis per angiography or duplex ultrasound

History of carotid revascularization (catheter-based or surgical)

Documented peripheral arterial disease (PAD; one or more of the following primary criteria must be satisfied):

ABI < 0.9 with symptoms of intermittent claudication

History of aorto-iliac or peripheral arterial intervention (catheter-based or surgical)

OR

CV Risk Category 2: defined as patients with:

Diabetes mellitus (Type 1 or Type 2) requiring treatment with medication AND

Men and women ≥ 50 years of age AND

One of the following at Visit 1 (additional risk factor for CVD):

Men ≥ 55 years of age or women ≥ 65 years of age;

Cigarette smoker or stopped smoking within 3 months before Visit 1;

Hypertension (blood pressure ≥ 140 mmHg systolic OR ≥ 90 mmHg diastolic) or on antihypertensive medication;

HDL-C ≤ 40 mg/dL for men or ≤ 50 mg/dL for women;

Hs-CRP > 3.00 mg/L (0.3 mg/dL);

Renal dysfunction: CrCL > 30 and < 60 mL/min (> 0.50 and < 1.00 mL/sec);

Retinopathy, defined as any of the following: non-proliferative retinopathy, preproliferative retinopathy, proliferative retinopathy, maculopathy, advanced diabetic eye disease or a history of photocoagulation;

Micro- or macroalbuminuria. Microalbuminuria is defined as either a positive micral or other strip test (may be obtained from medical records), an albumin creatinine ratio ≥ 2.5 mg/mmol or an albumin excretion rate on timed collection ≥ 20 mg/min all on at least two successive occasions; macroalbuminuria, defined as albustix or other dipstick evidence of gross proteinuria, an albumin:creatinine ratio ≥ 25 mg/mmol or an albumin excretion rate on timed collection ≥ 200 mg/min all on at least two successive occasions;

ABI < 0.9 without symptoms of intermittent claudication (patients with ABI < 0.9 with symptoms of intermittent claudication are counted under CV Risk Category 1).

Patients with diabetes with CVD as defined above are eligible based on the CVD requirements and will be counted

20

under CV Risk Category 1. Only patients with diabetes and no documented CVD as defined above need at least one additional risk factor as listed, and will be counted under CV Risk Category 2.

Women may be enrolled if all 3 of the following criteria are met:

They are not pregnant;

They are not breastfeeding;

They do not plan on becoming pregnant during the study.

Women of child-bearing potential must have a negative urine pregnancy test before randomization.

Women are not considered to be of childbearing potential if they meet one of the following criteria as documented by the investigator:

They have had a hysterectomy, tubal ligation or bilateral oophorectomy prior to signing the informed consent form;

They are post-menopausal, defined as ≥ 1 year since their last menstrual period or have a follicle-stimulating hormone (FSH) level in a menopausal range.

Women of childbearing potential must agree to use an acceptable method of avoiding pregnancy from screening to the end of the study, unless their sexual partner(s) is/are surgically sterile or the woman is abstinent.

Understanding of the study procedures, willing to adhere to the study schedules, and agreement to participate in the study by giving informed consent prior to screening.

Agree to follow a physician recommended diet and to maintain it through the duration of the study.

Exclusion Criteria

Patients are excluded from participation in the study if any of the following criteria apply:

Severe (class IV) heart failure.

Any life-threatening disease expected to result in death within the next 2 years (other than CVD).

Active severe liver disease (evaluated at Visit 1): cirrhosis, active hepatitis, ALT or AST $> 3 \times$ ULN, or biliary obstruction with hyperbilirubinemia (total bilirubin $> 2 \times$ ULN).

Hemoglobin A1c $> 10.0\%$ (or 86 mmol/mol IFCC units) at screening (Visit 1). If patients fail this criterion (HbA1c $> 10.0\%$ or 86 mmol/mol IFCC units) at Visit 1, they may have their antidiabetic therapy optimized and be retested at Visit 1.1.

Poorly controlled hypertension: blood pressure ≥ 200 systolic mmHg OR ≥ 100 mmHg diastolic (despite antihypertensive therapy).

Planned coronary intervention (such as stent placement or heart bypass) or any non-cardiac major surgical procedure. Patients can be (re)evaluated for participation in the trial (starting with Visit 1.1) after their recovery from the intervention/surgery.

Known familial lipoprotein lipase deficiency (Fredrickson Type I), apolipoprotein C-II deficiency, or familial dysbeta-lipoproteinemia (Fredrickson Type III).

Participation in another clinical trial involving an investigational agent within 90 days prior to screening (Visit 1). Patients cannot participate in any other investigational medication or medical device trial while participating in this study (participation in a registry or observational study without an additional therapeutic intervention is allowed).

Intolerance or hypersensitivity to statin therapy.

Known hypersensitivity to any ingredients of the study product or placebo; known hypersensitivity to fish and/or shellfish.

History of acute or chronic pancreatitis.

Malabsorption syndrome and/or chronic diarrhea (Note: patients who have undergone gastric/intestinal bypass sur-

US 10,568,861 B1

21

gery are considered to have malabsorption, hence are excluded; patients who have undergone gastric banding are allowed to enter the trial).

Non-study drug related, non-statin, lipid-altering medications, supplements or foods:

Patients are excluded if they used niacin >200 mg/day or fibrates during the screening period (after Visit 1) and/or plan to use during the study; patients who are taking niacin >200 mg/day or fibrates during the last 28 days before Visit 1 need to go through washout of at least 28 days after their last use and have their qualifying lipids measured (TG and LDL-C) after the washout period (Visit 1.1);

Patients are excluded if they take any omega-3 fatty acid medications (prescription medicines containing EPA and/or DHA) during the screening period (after Visit 1) and/or plan to use during the treatment/follow-up period of the study. To be eligible for participation in the study, patients who are taking omega-3 fatty acid medications during the last 28 days before Visit 1 (except patients in The Netherlands), need to go through a washout period of at least 28 days after their last use and have their qualifying lipids measured (TG and LDL-C) after the washout period (at Visit 1.1);

For patients in The Netherlands only: patients being treated with omega-3 fatty acid medications containing EPA and/or DHA are excluded; no washout is allowed.

Patients are excluded if they use dietary supplements containing omega-3 fatty acids (e.g., flaxseed, fish, krill, or algal oils) during the screening period (after Visit 1) and/or plan to use during the treatment/follow-up period of the study. To be eligible for participation in the study, patients who are taking >300 mg/day omega-3 fatty acids (combined amount of EPA and DHA) within 28 days before Visit 1 (except patients in The Netherlands), need to go through a washout period of at least 28 days since their last use and have their qualifying lipid measurements measured (TG and LDL-C) after the washout period (at Visit 1.1);

For patients in The Netherlands only: patients being treated with dietary supplements containing omega-3 fatty acids of >300 mg/day EPA and/or DHA are excluded; no washout is allowed.

Patients are excluded if they use bile acid sequestrants during the screening period (after Visit 1) and/or plan to use during the treatment/follow-up period of the study. To be eligible for participation in the study, patients who are taking bile acid sequestrants within 7 days before Visit 1, need to go through a washout period of at least 7 days since their last use and have their qualifying lipid measurements measured (TG and LDL-C) after the washout period (at Visit 1.1);

Other medications (not indicated for lipid alteration):

Treatment with tamoxifen, estrogens, progestins, thyroid hormone therapy, systemic corticosteroids (local, topical, inhalation, or nasal corticosteroids are allowed), HIV-protease inhibitors that have not been stable for ≥ 28 days prior to the qualifying lipid measurements (TG and LDL-C) during screening. To be eligible for participation in the study, patients who are not taking a stable dose of these medications within 28 days before Visit 1, need to go through a stabilization period of at least 28 days since their last dose change and have their qualifying lipid measurements measured (TG and LDL-C) after the washout period (at Visit 1.1).

Patients are excluded if they use cyclophosphamide or systemic retinoids during the screening period (after Visit 1) and/or plan to use during the treatment/follow-up period of the study. To be eligible for participation in the study, patients who are taking these medications within 28 days before Visit 1, need to go through a washout period of at

22

least 28 days since their last use and have their qualifying lipid measurements measured (TG and LDL-C) after the washout period (at Visit 1.1).

Known to have AIDS (patients who are HIV positive without AIDS are allowed).

Requirement for peritoneal dialysis or hemodialysis for renal insufficiency or if creatinine clearance (CrCL) <30 mL/min (0.50 mL/sec).

Unexplained creatine kinase concentration >5 \times ULN or creatine kinase elevation due to known muscle disease (e.g., polymyositis, mitochondrial dysfunction) at Visit 1.

Any condition or therapy which, in the opinion of the investigator, might pose a risk to the patient or make participation in the study not in the patient's best interest.

Drug or alcohol abuse within the past 6 months, and unable/unwilling to abstain from drug abuse and excessive alcohol consumption during the study or drinking 5 units or more for men or 4 units or more for women in any one hour (episodic excessive drinking or binge drinking). Excessive alcohol consumption is on average >2 units of alcohol per day. A unit of alcohol is defined as a 12-ounce (350 mL) beer, 5-ounce (150 mL) wine, or 1.5-ounce (45 mL) of 80-proof alcohol for drinks.

Mental/psychological impairment or any other reason to expect patient difficulty in complying with the requirements of the study or understanding the goal and potential risks of participating in the study (evaluated at Visit 1).

Study Procedures

Assessment Schedule

Screening Period

Screening Visit (Visit 1)

Patients will come to the Research Site for Visit 1. They will be instructed to fast for at least 10 hours before their visit.

If patients qualify for randomization based on the procedures at Visit 1, they need to be randomized within 60 days after Visit 1. The following procedures will be performed at the screening visit:

Obtain signed informed consent

Assign the patient a patient number

Obtain medical, surgical and family history

Record demographics

Obtain height, weight, and body mass index

Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)

Obtain a 12-lead electrocardiogram

Evaluate inclusion/exclusion criteria

This includes procedures and (fasting) blood samples (for example, hs-CRP, calculated creatinine clearance) as needed to determine the CV risk category (see inclusion criteria)

Obtain fasting blood samples for chemistry and hematology testing

Obtain a fasting blood sample for the lipid profile (TG, TC, HDL-C, LDL-C, non-HDL-C, VLDL-C)

Perform a urine pregnancy test on women of childbearing potential

Record concomitant medication(s)

Instruct patient to fast for at least 10 hours prior to the next visit

Screening Visit (Visit 1.1)

Some patients will skip Visit 1.1: Patients who qualify for study participation after Visit 1 because they meet all inclusion criterion and none of the exclusion criteria, may return to the Research Site for Visit 2 to be randomized and to start the treatment/follow-up period of the study. For these patients, Visit 2 will occur soon after Visit 1.

US 10,568,861 B1

23

Patients, who do not qualify at Visit 1, may return to the Research Site for a second qualifying visit (Visit 1.1) at the discretion of the investigator. At Visit 1.1, procedures that caused failure of eligibility at Visit 1 will be repeated. Patients will be eligible for randomization after Visit 1.1 if they meet all inclusion criteria and if they no longer fail the exclusion criteria. If patients are evaluated at Visit 1.1 and qualify for randomization based on the repeated procedures at Visit 1.1, they need to be randomized within 60 days after Visit 1.

For some patients, Visit 1.1 will be mandatory at least 28 days after Visit 1 in order to check eligibility. These are patients who at Visit 1 started treatment with a statin, changed their statin, changed the daily dose of their statin, started to washout prohibited medications or started a stabilization period with certain medications (see inclusion/exclusion criteria for details). Any of these changes at Visit 1 may affect the qualifying lipid levels and therefore, patients will need to have Visit 1.1 to determine whether they qualify based on lipid level requirements (TG and LDL-C) determined at Visit 1. Other procedures that caused failure of eligibility at Visit 1 will also be repeated at Visit 1.1.

The following procedures will be performed at the screening visit:

Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)

Evaluate inclusion/exclusion criteria; only those evaluations will be repeated that deemed the patient not eligible on Visit 1.

Obtain fasting blood samples for chemistry and hematology testing. Only those samples will be obtained that deemed the patient not eligible on Visit 1.

Obtain a fasting blood sample for the lipid profile (TG, TC, HDL-C, LDL-C, non-HDL-C, VLDL-C) if the patient was deemed not eligible on Visit 1. This includes patients who at Visit 1 started treatment with a statin, changed their statin, changed the daily dose of their statin, started to washout prohibited medications or started a stabilization period with certain medications (see inclusion/exclusion criteria for details). These patients will have a fasting blood sample collected at Visit 1.1 for the qualifying lipid values (TG and LDL-C), and the TG and LDL-C inclusion criteria will be evaluated.

Record concomitant medication(s)

Treatment/Follow-Up Period

Every attempt should be made to complete the follow-up visits during the defined window periods.

Randomization visit (Visit 2; Day 0)

Qualified patients will return to the Research Site for Visit 2.

The following procedures will be performed at Visit 2:

Perform physical examination

Obtain weight

Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)

Measure waist circumference (one of the factors to diagnose metabolic syndrome)

Obtain a 12-lead electrocardiogram

Evaluate inclusion/exclusion criteria

Obtain fasting blood samples for:

Chemistry and hematology testing

Lipid profile (baseline)

Biomarker assays (baseline)

Genetic testing (optional blood sample)

Archiving (in countries and at sites approved by IRB/IEC and dependent on country regulations)

24

Perform a urine pregnancy test on women of childbearing potential (must be negative for randomization)

Dispense study drug and record randomization number

Instruct patient on how to take study drug

Administer study drug—Note: Study drug should be taken orally with food following the collection of all fasting blood samples

Assess for and record adverse events

Record concomitant medication(s)

Instruct patient:

To bring all study supplies with them to the next visit

Not to take study drug on the morning of their next visit

To fast for ≥ 10 hours prior to the next visit

Visit 3 (Day 120; ~4 Months)

Patients will return to the Research Site for Visit 3 on Day 120 \pm 10 days.

The following procedures will be performed:

Perform physical examination

Obtain weight

Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)

Obtain fasting blood samples for:

Chemistry and hematology testing

Lipid profile

Review study drug compliance by unused capsule count; discuss with and counsel patients about compliance if needed

Administer study drug—Note: Study drug should be taken orally with food following the collection of all fasting blood samples

Assess and record efficacy events

Assess for and record adverse events

Record concomitant medication(s)

Instruct patient:

To bring all study supplies with them to the next visit

Not to take study drug on the morning of their next visit

To fast for ≥ 10 hours prior to the next visit

Visits 4, 5, 6 and 7

At Visit 4: Day 360 \pm 10; Visit 5: Day 720 \pm 10; Visit 6: Day 1080 \pm 10; and Visit 7: Day 1440 \pm 10, the following procedures will be performed:

Perform physical examination

Obtain weight

Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)

Measure waist circumference (collected at Visit 5 only)

Obtain a 12-lead electrocardiogram

Obtain fasting blood samples for:

Chemistry and hematology testing

Lipid profile

Biomarker assays (collected at Visit 5 only)

Archiving (in countries and at sites approved by IRB/IEC and dependent on country regulations)

Review study drug compliance by unused capsule count; discuss with and counsel patients about compliance if needed

Administer study drug—Note: Study drug should be taken orally with food following the collection of all fasting blood samples

Assess and record efficacy events

Assess for and record adverse events

Record concomitant medication(s)

Instruct patient:

To bring all study supplies with them to the next visit

Not to take study drug on the morning of their next visit

To fast for ≥ 10 hours prior to the next visit

US 10,568,861 B1

25

Additional Visits

The end date of the study is expected for Day 1800 but the actual end date will be dependent on the determination of the study end date by the DMC. The study end date is determined to be when approximately 1612 primary efficacy events have occurred. If the actual study end date is later than the expected end date, additional visits will be planned between Visit 7 and the Last Visit with a maximum of 360±10 days between visits. If the actual study end date is sooner than the expected end date, fewer visits will occur, and the last visit (See Section 6.1.2.5) will occur sooner.

On additional visits the same procedures will be performed as listed in Section 6.1.2.3. Irrespective of the number of additional visits, after the DMC has established the end of the study date, there will be a last visit with procedures as listed in Section 6.1.2.5.

Last Visit—End of Study

All patients will complete the study at the same time (within a 30-day window after the study end date), irrespective of the date that they were randomized. The end date of the study is planned for Day 1800 but the actual end date will be dependent on the determination of the study end date when approximately 1612 primary efficacy events have occurred (event-driven trial). For each patient, the last visit may occur within 30 day after the actual study end date. However, for the efficacy endpoints based on CV events, only events occurring up to and including the scheduled actual study end date will be included in the efficacy analyses.

A final follow-up visit is required for all patients. In the rare cases that a final follow-up visit cannot occur within the 30-day timeframe following the study end date, any attempt to contact the patient must be recorded on a special contact form, until/unless appropriate information is obtained.

At the Last Visit, the following procedures will be performed:

- Perform physical examination
- Obtain weight
- Obtain vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature)
- Measure waist circumference
- Obtain a 12-lead electrocardiogram
- Obtain fasting blood samples for:
 - Chemistry and hematology testing
 - Lipid profile
 - Biomarker assays
- Archiving (in countries and at sites approved by IRB/IEC and dependent on country regulations)
- Determine study drug compliance by unused capsule count

- Assess and record efficacy events
- Assess for and record adverse events
- Record concomitant medication(s)
- Telephone Follow-up Contact
- Site personnel will contact each patient by telephone on the following study days:

- Day 60±3 days
- Day 180±5 days
- Day 270±5 days
- Day 450±5 days
- Day 540±5 days
- Day 630±5 days
- Day 810±5 days
- Day 900±5 days
- Day 990±5 days
- Day 1170±5 days
- Day 1260±5 days

26

Day 1350±5 days

Day 1530±5 days

Day 1620±5 days

Day 1710±5 days

If the treatment/follow-up period of the study is extended beyond the expected end date (Day 1800), additional follow-up phone calls will be made every 3 months in-between additional visits±5 days. If the treatment/follow period of the study is shorter than the expected end date, less follow-up phone calls will be needed.

Every attempt will be made to talk to each patient within this time frame.

The following information will be collected from the patient:

Possible efficacy endpoints related to CV events. Patients will be asked to return to the Research Site to assess for any endpoints or events identified.

Adverse events

Concomitant medications

Current address and contact information (update if changed or will be changing)

Patients will be reminded about the following items:

To take the study medication according to the dosing schedule assigned, with food

When to return to the Research Center for the next visit

To bring the unused study medication to the next visit

To not take study drug on the morning of their next visit

To fast for at least 10 hours prior to the next visit

Laboratory Procedures**Clinical Laboratory Procedures**

All clinical laboratory determinations for screening and safety will be performed by a certified clinical laboratory under the supervision of the Sponsor or its designee.

Whenever possible and appropriate, samples for the clinical laboratory procedures will be collected after fasting for at least 10 hours. For the purposes of this study, fasting is defined as nothing by mouth except water (and any essential medications).

The investigator must review and sign all laboratory test reports. At screening, patients who have laboratory values that are outside the exclusionary limits specified in the exclusion criteria may not be enrolled in the study (patients can be considered for the study if values are classified as not clinically significant by the investigator). After randomization, the investigator will be notified if laboratory values are outside of their normal range. In this case, the investigator will be required to conduct clinically appropriate follow-up procedures.

Safety Laboratory Tests

The safety laboratory tests include:

Hematology with complete blood count (CBC), including RBC, hemoglobin (Hgb), hematocrit (Hct), white cell blood count (WBC), white cell differential, and platelet count

Biochemistry panel including total protein, albumin, alkaline phosphatase, alanine aminotransferase (ALT/SGPT), aspartate aminotransferase (AST/SGOT), total bilirubin, glucose, calcium, electrolytes (sodium, potassium, chloride), blood urea nitrogen (BUN), serum creatinine, uric acid, creatine kinase, and HbA1c.

Fasting Lipid Profile

The fasting lipid panel includes: TG, TC, LDL-C, HDL-C, non-HDL-C, and VLDL-C.

At all visits, LDL-C will be calculated using the Friedewald equation. At Visit 1 and Visit 1.1 Direct LDL-C will be used if at the same visit TG>400 mg/dL (4.52 mmol/L). These LDL-C values will be used for the evaluation of the LDL-C inclusion criterion (LDL-C qualifying measure-

US 10,568,861 B1

27

ments for randomization) and for the assessment of changes in the statin therapy when LDL-C is not at goal. At all remaining visits (except Visit 2 and Visit 4) LDL-C will be measured by Direct LDL Cholesterol or by Preparative Ultracentrifugation if at the same visit TG>400 mg/dL (4.52 mmol/L). In addition, irrespective of the TG levels, at Visit 2 (0 Months of Follow-up, baseline) and at Visit 4 (12 Months of Follow-up), LDL-C will be measured by Preparative Ultracentrifugation. These Preparative Ultracentrifugation LDL-C measurements will be used in the statistical analysis including the calculation of the percent change from baseline (1 year versus baseline).

Genetic Testing

A fasting blood sample will be stored for future genetic testing at the discretion of the sponsor. The specifics of this test will be determined at a later date. This sample is optional as local regulations may prohibit genetic samples to be collected or shipped outside the country, or patients may not consent.

Research on genetic testing will look for links between genes and certain diseases, including their treatment(s) such as medicines and medical care. The blood samples will be collected in the study center with the regular protocol-required labs. Each patient tube with sample for genetic testing will be labeled with patient number only. The site will maintain a Subject Code Identification List for cross-reference. The patient number does not contain any identifiable information (i.e. Patient initials, date of birth, etc). Un-analyzed samples will be stored frozen by the sponsor for a period of up to 2 years following the end of the study, at which time they will be destroyed. If samples are tested, results will not be reported to the patient, parents, relatives, or attending physician and will not be recorded in the patient's medical records. There will be no follow-up contact with the sites or patients regarding this sample. The subject can withdraw their consent for genetic testing at any time up to analysis, even after the sample has been obtained. The subject can notify the site in writing that they withdraw their consent for the genetic testing portion of the study, and it will be documented by the site in the subject chart, as well as captured in the CRF. The lab will be notified to pull the sample and destroy it.

Biomarkers Assays

The biomarker assays include: hs-CRP, Apo B and hsTnT. Additional laboratory tests

Additional laboratory tests include:

A urine pregnancy test will be administered to women of childbearing potential at certain visits as listed in schedule of procedures (Table 1). The urine pregnancy tests will be performed at the Research Site utilizing marketed test kits, or at a certified clinical laboratory.

A fasting blood sample (12 mL) for archiving. This sample will be collected only at sites in countries where allowed by local regulations and at sites for which approved by the IRB or IEC. The plasma from the archiving sample will be stored frozen in 2 separate equal aliquots, and will be used at the Sponsor's discretion to perform repeat analyses described in the protocol or to perform other tests related to cardiovascular health.

Blinding of Laboratory Results

All efficacy laboratory results during the double-blind period of the trial will be blinded (values not provided) to patients, investigators, pharmacists and other supporting staff at the Research Sites, personnel and designees of the Sponsor, study administrators and personnel at the organization(s) and vendors managing and/or supporting the study,

28

with the exception of the laboratory personnel conducting the assays. To ensure patient safety, hsTnT values will be reported to the site.

Flagging of Critical Lab Values

Critical lab values are values that may warrant medical intervention to avoid possible harm to a patient. Critical lab values will be defined in the Laboratory Manual for the study, and the Research Site will be notified of the occurrence of a critical lab value (critical high or critical low) by a special annotation (flag) in the laboratory reports provided to the Research Sites. Although laboratory values that are part of the efficacy endpoints during the double-blind period of the study will not be provided to the Research Site (see Section 6.3.1.6), the sites will be notified when the TG value of a patient sample is >1000 mg/dL (11.29 mmol/L) (critical high TG value) or if the LDL-C values of a patient sample is >130 mg/dL (3.37 mmol/L) (critical high LDL-C value). These critical high values will need to be confirmed by a repeat measurement (new fasting blood sample) within 7 days. TG value of >2000 mg/dL (22.58 mmol/L) will also be flagged, so that appropriate medical action can be taken by the investigator as soon as possible.

If TG values are confirmed critically high, patients may be discontinued from study drug with the option to remain on study. The investigator should use the best clinical judgment for each patient which could include the use of approved TG-lowering medications after patients have been discontinued from study drug.

If LDL-C values are confirmed critically high, the investigator may need to take appropriate medical action which could include: reinforce/intensify therapeutic lifestyle changes (including diet and physical activity), increase the dose of the present statin therapy, add ezetimibe, or prescribe a more potent statin to lower LDL-C. The investigator should use the best clinical judgment for each patient.

Medical Procedures

Medical, Surgical and Family History

Medical history, including family history and details regarding all illnesses and allergies, date(s) of onset, status of current condition, and smoking and alcohol use will be collected on all patients.

Demographics

Demographic information including day, month, and year of birth, race, and gender will be collected for all patients.

Vital Signs

Vital signs include systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature. Blood pressure will be measured using a standardized process:

Patient should sit for ≥5 minutes with feet flat on the floor and measurement arm supported so that the midpoint of the manometer cuff is at heart level.

Use a mercury sphygmomanometer or automatic blood pressure device with an appropriately sized cuff with the bladder centered over the brachial artery.

Blood pressure should be recorded to the nearest 2 mmHg mark on the manometer or to the nearest whole number on an automatic device. A blood pressure reading should be repeated 1 to 2 minutes later, and the second reading should also be recorded to the nearest 2 mmHg mark.

Physical Examination

A physical examination must include source documentation of general appearance, skin, and specific head and neck, heart, lung, abdomen, extremities, and neuromuscular assessments.

US 10,568,861 B1

29

Height, Weight and Body Mass Index

Height and weight will be measured. Measurement of weight should be performed with the patient dressed in indoor clothing, with shoes removed, and bladder empty.

Waist Circumference

Waist circumference will be measured with a tape measure, as follows: Start at the top of the hip bone then bring the tape measure all the way around—level with the navel. Make sure the tape measure is snug, but without compressing the skin, and that it is parallel with the floor.

Patients should not hold their breath while measuring waist circumference.

Electrocardiogram (ECG)

ECGs (standard 12-lead) will be obtained annually. Site personnel should make every attempt to perform a patient's ECG using the same equipment at each visit. ECGs will be reviewed by the site for the detection of silent MI. Silent MIs will be sent for event adjudication.

Treatment and Restrictions**Treatment****Treatment Regimen, Dosage, and Duration**

Eligible study patients will be randomly assigned on Day 0 to one of the 2 treatment groups. Patients in each group will receive either 4 g/day AMR101 or placebo for up to 4.75 years (4 years planned median treatment duration) according to Table 2.

The daily dose of study drug is 4 capsules per day taken as two capsules take on two occasions per day (2 capsules given twice daily).

TABLE 2

Dosing Schedule during the Treatment Period		
Treatment Group	Daily Dose	Number of Capsules per Day
1	4 g	4 capsules of 1000 mg AMR101
2	Placebo	4 capsules of matching placebo

Patients will be instructed to take study drug with food (i.e., with or at the end of their morning and evening meals). On days that patients are scheduled for study visits, the daily dose of study drug will be administered by site personnel with food provided by the site following collection of all fasting blood samples. For the purposes of this study, fasting is defined as nothing by mouth except water (and any essential medications) for at least 10 hours.

Treatment Assignment**Identification Number**

A unique patient identification number (patient number) will be established for each patient at each site. The patient number will be used to identify the patient throughout the study and will be entered on all documentation. If a patient is not eligible to receive treatment, or if a patient discontinues from the study, the patient number cannot be reassigned to another patient. The patient number will be used to assign patients to one of the 2 treatment groups according to the randomization schedule.

Drug Randomization

Only qualified patients who meet all of the inclusion criteria and none of the exclusion criteria will be randomized and will receive study medication starting at Visit 2 (Day 0). Eligible patients will be randomly assigned to one of the 2 treatment groups. Randomization will be stratified by CV risk category, use of ezetimibe and by geographical region (Westernized, Eastern European, and Asia Pacific countries) (See Section 3.10). Approximately 70% of randomized

30

patients will be in the CV Risk Category 1, including patients with established CVD, and approximately 30% of randomized patients will be in the CV Risk Category 2, including patients with diabetes and at least one additional risk factor but no established CVD. Enrollment with patients of a CV risk category will be stopped when the planned number of patients in that risk category is reached.

Emergency Unblinding

In an emergency, when knowledge of the patient's treatment assignment is essential for the clinical management or welfare of the patient, the investigator may request the patient's treatment assignment for unblinding. Prior to unblinding the patient's individual treatment assignment, the investigator should assess the relationship of an adverse event to the administration of the study drug (Yes or No). If the blind is broken for any reason, the investigator must record the date and reason for breaking the blind on the appropriate Case Report Form (CRF) and source documents.

Compliance Control

It is recommended that, unless clear contraindications arise, patients be strongly encouraged to adhere to their treatment regimen with the study drug for the duration of the trial. Any interruptions of therapy should, if possible, be brief (e.g., <4 weeks) and only for clinically indicated reasons, such as adverse events. Discontinuations will be discouraged as much as possible. Any discontinuations should be based on compelling clinical reasons.

For every patient, an assessment of compliance to the study drug treatment regimen must be obtained at each scheduled visit. Study medication will be dispensed in amounts exceeding the amount required for the study. Patients will be instructed to return all unused study medication at the next visit. Compliance to the study drug regimen will be evaluated at each visit by counting unused capsules. Discrepancies will be evaluated and discussed with each patient to assess compliance. If compliance is unsatisfactory, the patient will be counseled about the importance of compliance to the dosing regimen. At the end of the study, the final study medication compliance will be determined by unused capsule count.

Study Restrictions**Concomitant Medications During Treatment/Follow-Up Period**

Any medications administered during the study period must be documented on the Concomitant Medication CRF. Patients must not have taken any investigational agent within 90 days prior to screening. Patients cannot participate in any other investigational medication trial while participating in this study.

The following non-study drug related, non-statin, lipid-altering medications and supplements, and foods are prohibited during the study (from Visit 1 until after the Last Visit-End of Study), except for compelling medical reasons in ODIS patients:

- niacin >200 mg/day;
- fibrates;
- prescription omega-3 fatty acid medications;
- dietary supplements containing omega-3 fatty acids (e.g., flaxseed, fish, krill, or algal oils);
- bile acid sequestrants;
- cyclophosphamide;
- systemic retinoids

If any of these products would be used during the treatment/follow-up period of the study, it should be for compelling medical reasons in ODIS patients, and it should be documented in the Concomitant Medication CRF. If the

US 10,568,861 B1

31

ODIS patient agrees to restart study medication, the use of excluded medication must be discontinued.

Foods enriched with omega-3 fatty acids are strongly discouraged after Visit 1 for the duration of the study (does not apply to The Netherlands or Canada only. Therefore, all centers in The Netherlands and Canada must ignore this request).

The following products are allowed: statins, ezetimibe, and herbal products & dietary supplements not containing omega-3 fatty acids.

Statins:

The same statin at the same dose should be continued until the end of the study, unless deemed medically necessary to change because of an adverse event or lack of efficacy (LOE). It is preferred that if LOE is the determining factor that ezetimibe be added to the present dose.

Switching between a brand name statin and the generic version of the same statin is allowed at any time during the study.

Statins may be administered with or without ezetimibe.

Based on the FDA recommendation, simvastatin 80 mg be used only in patients who have been taking this dose for 12 months or more and have not experienced any muscle toxicity. (See reference: FDA Drug Safety Communication: Ongoing safety review of high-dose Zocor (simvastatin) and increased risk of muscle injury. (<http://www.fda.gov/Drugs/DrugSafety/PostmarketDrugSafetyInformationforPatientsandProviders/ucm204882.htm>))

Changing of the type of statin or the statin dose during the treatment/follow-up period of the study should only be done for compelling medical reasons and must be documented in the CRF.

LDL-C Rescue:

If the level of LDL-C exceeds 130 mg/dL (3.37 mmol/L) during the study (initial measurement and confirmed by a second determination at least 1 week later), the investigator may either increase the dose of the present statin therapy or may add ezetimibe to lower LDL-C. The investigator should use the best clinical judgment for each patient.

No data are available with regard to potential interactions between ethyl-EPA and oral contraceptives. There are no reports suggesting that omega-3 fatty acids, including ethyl-EPA, would decrease the efficacy of oral contraceptives.

Patient Restrictions

Beginning at the screening visit, all patients should be instructed to refrain from excessive alcohol consumption, to follow a physician recommended diet and to maintain it through the duration of the study. Excessive alcohol consumption is on average 2 units of alcohol per day or drinking 5 units or more for men or 4 units or more for women in any one hour (episodic excessive drinking or binge drinking). A unit of alcohol is defined as a 12-ounce (350 mL) beer, 5-ounce (150 mL) wine, or 1.5-ounce (45 mL) of 80-proof alcohol for drinks.

Investigational Product

Clinical Trial Material

The following will be supplied by the Sponsor:

AMR101 1000 mg capsules

Placebo capsules

The Sponsor will supply sufficient quantities of AMR101 1000 mg capsules and placebo capsules to allow for completion of the study. The lot numbers of the drugs supplied will be recorded in the final study report.

Records will be maintained indicating the receipt and dispensation of all drug supplies. At the conclusion of the study, any unused study drug will be destroyed.

32

Pharmaceutical Formulations

AMR101 1000 mg and placebo capsules (paraffin) are provided in liquid-filled, oblong, gelatin capsules. Each capsule is filled with a clear liquid (colorless to pale yellow in color). The capsules are approximately 25.5 mm in length with a diameter of approximately 9.5 mm.

Labeling and Packaging

Study medication will be packaged in high-density polyethylene bottles. Labeling and packaging will be performed according to GMP guidelines and all applicable country-specific requirements. The bottles will be numbered for each patient based on the randomization schedule. The patient randomization number assigned by IWR or a designee of the Sponsor for the study (if no IWR system is used), will correspond to the number on the bottles. The bottle number for each patient will be recorded in the Electronic Data Capture (EDC) system for the study.

Dispensing Procedures and Storage Conditions

Dispensing Procedures

At Visit 2 (Day 0), patients will be assigned study drug according to their treatment group determined by the randomization schedule. Once assigned to a treatment group, patients will receive study drug supplies. At each visit, patients will bring unused drug supplies dispensed to them earlier. From the drug supplies assigned to each patient, site personnel will administer drug while the patients are at the Research Site.

The investigator or designee must contact the IWR system or a designee of the Sponsor for the study (if no IWR system is used) when any unscheduled replacements of study medication are needed.

During the last visit during the treatment period, patients will bring the unused drug supplies for site personnel to calculate the final study medication compliance by unused capsule count.

Storage Conditions

At the Research Sites, study drugs must be stored at room temperature, 68° F. to 77° F. (20° C. to 25° C.). Do not allow storage temperature to go below 59° F. (15° C.) or above 86° F. (30° C.). Store in the original package.

Study drugs must be stored in a pharmacy or locked and secure storage facility, accessible only to those individuals authorized by the investigator to dispense the drug. The investigator or designee will keep accurate dispensing records. At the conclusion of the study, study site personnel will account for all used and unused study drug. Any unused study drug will be destroyed. The investigator agrees not to distribute study drug to any patient, except those patients participating in the study.

Efficacy Assessments

Specification of Variables and Procedures

The primary endpoint and the majority of the secondary and tertiary endpoints are based on clinical events related to CVD and mortality. All events occurring between randomization and the study end date (inclusive) must be recorded. Only adjudicated events will be included in the final analyses. Further details on the assessment of clinical events and their definitions will be found in the CEC charter.

Efficacy Endpoints

Primary Efficacy Endpoint

Time from randomization to the first occurrence of the composite of the following clinical events:

CV death,

Nonfatal MI (including silent MI; ECGs will be performed annually for the detection of silent MIs),

Nonfatal stroke,

Coronary revascularization

US 10,568,861 B1

33

Hospitalization for unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing.

The first occurrence of any of these major adverse vascular events during the follow-up period of the study will be included in the incidence.

Secondary Efficacy Endpoints

The key secondary efficacy endpoint is:

The composite of death from CV causes, nonfatal MI, coronary revascularization, unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization, nonfatal stroke, or peripheral CVD requiring intervention, angioplasty, bypass surgery, or aneurysm repair.

Other secondary efficacy endpoints are as follows (to be tested in said order):

The composite of total mortality, nonfatal MI, or nonfatal stroke;

The composite of death from CV causes, nonfatal MI, coronary revascularization, unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization, peripheral CVD requiring intervention, or cardiac arrhythmia requiring hospitalization;

The composite of death from CV causes, nonfatal MI, coronary revascularization, or unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing and requiring emergent hospitalization;

The composite of death from CV causes or nonfatal MI; Total mortality;

Fatal and nonfatal MI (including silent MI);

Coronary Revascularization;

Hospitalization for unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing;

Fatal and nonfatal stroke.

For the secondary endpoints that count a single event, the first occurrence of this type of event will be counted in each patient. For secondary endpoints that are composites of two or more types of events, the first occurrence of any of the event types included in the composite will be counted in each patient.

Tertiary Efficacy Endpoints:

The second, third, fourth, and fifth major CV event of the primary composite endpoint. The type of (nonfatal) events may occur in any order.

Primary endpoint in subset of patients with diabetes mellitus;

Primary endpoint in subset of patients with metabolic syndrome;

New CHF, new CHF leading to hospitalization, transient ischemic attack, amputation for CVD and carotid revascularization;

Elective coronary revascularization and emergent coronary revascularization;

New onset diabetes;

Fasting TG, TC, LDL-C, HDL-C, non-HDL-C, VLDL-C, apo B, hs-CRP, and hsTnT: effect of baseline and on-treatment change of biomarkers on primary and key secondary endpoints;

CV mortality;

Cardiac Arrhythmias requiring hospitalization;

Cardiac Arrest;

To explore the effect of AMR101 on weight and waist circumference.

For the tertiary endpoints that count a single event, the first occurrence of this type of event will be counted in each

34

patient. For tertiary endpoints that are composites of two or more types of events, the first occurrence of any of the event types included in the composite will be counted in each patient (except when stated otherwise, for the second, third, fourth, and fifth major CV event).

Safety Assessments

Specification of Variables and Procedures

Safety assessments will include adverse events, clinical laboratory measurements (chemistry, hematology), 12-lead ECGs, vital signs (systolic and diastolic blood pressure, heart rate, respiratory rate, and body temperature), and physical examinations as per Study Procedures/Table 1.

A complete medical, surgical and family history will be completed at Visit 1.

All laboratory test results must be evaluated by the investigator as to their clinical significance. Any observations at physical examinations or laboratory values considered by the investigator to be clinically significant should be considered an adverse event.

Adverse Events

An adverse event is defined as any untoward medical occurrence, which does not necessarily have a causal relationship with the medication under investigation. An adverse event can therefore be any unfavorable and/or unintended sign (including an abnormal laboratory finding), symptom, or disease temporally associated with the use of an investigational medication product, whether or not related to the investigational medication product. All adverse events, including observed or volunteered problems, complaints, or symptoms, are to be recorded on the appropriate CRF. Each adverse event is to be evaluated for duration, intensity, and causal relationship with the study medication or other factors.

Adverse events, which include clinical laboratory test variables, will be monitored from the time of informed consent until study participation is complete. Patients should be instructed to report any adverse event that they experience to the investigator. Beginning with Visit 2, investigators should assess for adverse events at each visit and record the event on the appropriate adverse event CRF.

Wherever possible, a specific disease or syndrome rather than individual associated signs and symptoms should be identified by the investigator and recorded on the CRF. However, if an observed or reported sign or symptom is not considered a component of a specific disease or syndrome by the investigator, it should be recorded as a separate adverse event on the CRF.

Any medical condition that is present when a patient is screened or present at baseline that does not deteriorate should not be reported as an adverse event. However, medical conditions or signs or symptoms present at baseline and that change in severity or seriousness at any time during the study should be reported as an adverse event.

Clinically significant abnormal laboratory findings or other abnormal assessments that are detected during the study or are present at baseline and significantly worsen will be reported as adverse events or SAEs. The investigator will exercise his or her medical and scientific judgment in deciding whether an abnormal laboratory finding or other abnormal assessment is clinically significant.

The investigator will rate the severity (intensity) of each adverse event as mild, moderate, or severe, and will also categorize each adverse event as to its potential relationship to study drug using the categories of Yes or No.

Severity:

Mild—An event that is usually transient in nature and generally not interfering with normal activities.

US 10,568,861 B1

35

Moderate—An event that is sufficiently discomforting to interfere with normal activities.

Severe—An event that is incapacitating with inability to work or do usual activity or inability to work or perform normal daily activity.

Causality Assessment:

The relationship of an adverse event to the administration of the study drug is to be assessed according to the following definitions:

No (unrelated, not related, no relation)—The time course between the administration of study drug and the occurrence or worsening of the adverse event rules out a causal relationship and another cause (concomitant drugs, therapies, complications, etc.) is suspected.

Yes—The time course between the administration of study drug and the occurrence or worsening of the adverse event is consistent with a causal relationship and no other cause (concomitant drugs, therapies, complications, etc.) can be identified.

The following factors should also be considered:

The temporal sequence from study medication administration

The event should occur after the study medication is given. The length of time from study medication exposure to event should be evaluated in the clinical context of the event.

Underlying, concomitant, intercurrent diseases

Each report should be evaluated in the context of the natural history and course of the disease being treated and any other disease the patient may have.

Concomitant medication

The other medications the patient is taking or the treatment the patient receives should be examined to determine whether any of them might be recognized to cause the event in question.

Known response pattern for this class of study medication

Clinical and/or preclinical data may indicate whether a particular response is likely to be a class effect.

Exposure to physical and/or mental stresses

The exposure to stress might induce adverse changes in the patient and provide a logical and better explanation for the event.

The pharmacology and pharmacokinetics of the study medication

The known pharmacologic properties (absorption, distribution, metabolism, and excretion) of the study medication should be considered.

Unexpected Adverse Events—An unexpected adverse event is an adverse event either not previously reported or where the nature, seriousness, severity, or outcome is not consistent with the current Investigator's Brochure.

Serious Adverse Events

A serious adverse event (SAE) is defined as an adverse event that meets any of the following criteria:

Results in death

Is life-threatening—Note: The term "life-threatening" in the definition of "serious" refers to an event in which the patient was at risk of death at the time of the event. It does not refer to an event, which hypothetically might have caused death, if it were more severe.

Requires hospitalization or prolongation of existing hospitalization—Note: In general, hospitalization for treatment of a pre-existing condition(s) that did not worsen from baseline is not considered adverse events and should not be reported as SAEs.

36

Results in disability/incapacity

Is a congenital anomaly/birth defect;

Is an important medical event—Note: Important medical events that may not result in death, be life threatening, or require hospitalization may be considered an SAE when, based upon appropriate medical judgment, they may jeopardize the patient and may require medical or surgical intervention to prevent one of the outcomes listed above. Examples of such medical events include allergic bronchospasm requiring intensive treatment in an emergency room or at home, blood dyscrasias or convulsions that do not result in inpatient hospitalizations, or the development of drug dependency.

By design of this study SAEs that are endpoint events will only be recorded for the endpoint determination and not captured as SAEs. The intention is that the endpoint events are not reported to IRBs as SAEs, unless the IRB requires that these are reported. Investigators should specifically inform their institution/IRB of this plan and confirm whether or not they want the endpoint events reported. By agreement with the US FDA, these endpoints will also not be reported to the US FDA as SAEs; rather they will be reported as endpoint events. Following adjudication if the event is determined to not meet the criteria for an event, the event will be evaluated as an SAE beginning with that day as Day 0.

Serious Adverse Event Reporting—Procedure for Investigators

Initial Reports

All SAEs occurring from the time of informed consent until 28 days following the last administration of study medication must be reported to the Sponsor or designee within 24 hours of the knowledge of the occurrence (this refers to any adverse event that meets any of the aforementioned serious criteria). SAEs that the investigator considers related to study medication occurring after the 28-day follow-up period will also be reported to the Sponsor or designee.

The investigator is required to submit SAE reports to the Institutional Review Board (IRB) or Independent Ethics Committee (IEC) in accordance with local requirements. All investigators involved in studies using the same investigational medicinal product (IMP) will receive any Suspected Unexpected Serious Adverse Reaction (SUSAR) reports for onward submission to their local IRB as required. All reports sent to investigators will be blinded.

In addition, regulatory agencies will be notified of SAEs per the requirements of the specific regulatory jurisdiction regulations and laws.

Follow-Up Reports

The investigator must continue to follow the patient until the SAE has subsided, or until the condition becomes chronic in nature, stabilizes (in the case of persistent impairment), or the patient dies. Within 24 hours of receipt of follow-up information, the investigator must update the SAE form electronically in the EDC system for the study and submit any supporting documentation (e.g., laboratory test reports, patient discharge summary, or autopsy reports) to the Sponsor or designee via fax or email.

Reporting by the Sponsor

IRBs and IECs will be informed of SUSARs according to local requirements. Cases will be unblinded for reporting purposes as required.

Exposure In Utero During Clinical Trials

If a patient becomes pregnant during the study, the investigator should report the pregnancy to the Sponsor or designee within 24 hours of being notified. The Sponsor or designee will then forward the Exposure In Utero form to the investigator for completion.

US 10,568,861 B1

37

The patient should be followed by the investigator until completion of the pregnancy. If the pregnancy ends for any reason before the anticipated date, the investigator should notify the Sponsor or designee. At the completion of the pregnancy, the investigator will document the outcome of the pregnancy. If the outcome of the pregnancy meets the criteria for immediate classification as an SAE (i.e., post-partum complication, spontaneous abortion, stillbirth, neonatal death, or congenital anomaly), the investigator should follow the procedures for reporting an SAE.

Treatment Discontinuation/Patient Withdrawal

Patients may withdraw from the study at any time and for any reason. Study drug administration may also be discontinued at any time, at the discretion of the investigator. In any case, follow-up for efficacy and safety should be continued.

Reasons for Early Study Drug Discontinuation

Study drug discontinuation should be avoided as much as possible, but may be done for any of the following reasons:

Patient withdraws consent or requests early discontinuation from the study for any reason. Patients should be encouraged to continue to participate in the study for the entire duration of the study even if they choose not to take study medication any longer.

Occurrence of a clinical or laboratory adverse event, either serious or non-serious, at the discretion of the investigator. The Sponsor or designee should be notified if a patient is discontinued because of an adverse event or laboratory abnormality. It is recommended that, unless clear contraindications arise, patients be strongly encouraged to adhere to their treatment regimen with the study drug for the duration of the trial. Any interruptions of therapy should, if possible, be brief (e.g., <4 weeks) and only for clinically indicated reasons, such as adverse events. The following should be considered reason for discontinuation:

ALT>3×ULN and bilirubin>1.5×ULN

ALT>5×ULN

ALT>3×ULN and appearance or worsening of hepatitis

ALT>3×ULN persisting for >4 weeks

ALT>3×ULN and cannot be monitored weekly for 4 weeks

Any medical condition or personal circumstance that, in the opinion of the investigator, exposes the patient to risk by continuing in the study or precludes adherence to the protocol.

Sponsor discontinues the study.

A TG value that is flagged as critically high, i.e., >1000 mg/dL (11.29 mmol/L), and confirmed as critically high by a repeat measurement (new fasting blood sample) within 7 days. In this case, a patient may be discontinued from study drug (with the option to remain ODIS) and other lipid-altering medications may be (re)initiated. If the TG value is flagged as >2000 mg/dL (22.58 mmol/L) then appropriate medical action can be taken by the investigator as soon as possible.

Occurrence of an outcome event according to the judgment of the investigator is not considered a valid reason for study drug discontinuation.

Patients whose treatment with study medication is discontinued early, and have not withdrawn consent, will stay in study and will be monitored until the end of the study. Patients that continue in the study after indefinite cessation of therapy will be characterized as Off Drug In Study (ODIS). ODIS patients should be asked to return to the study site for an interim visit once the patient has been off study drug for >30 days. Procedures at this visit are consistent with

38

those at Visit 5. If not contraindicated, patients will also have the option to restart study medication at any point once characterized as ODIS.

The reason for study drug discontinuation or interruption will be recorded on the CRF.

Follow-Up after Early Study Drug Discontinuation/Lost to Follow-Up

Patients who prematurely discontinue study drug are not to be replaced.

All randomized patients must be followed up according to the study flowchart until the study end date or death, regardless of whether they discontinue study drug prematurely or not. Any event occurring after early study drug discontinuation will be recorded up through the study end date.

In order to follow the medical status of the patients, especially when they discontinued the study, investigators are encouraged to obtain information from the patient's primary care practitioner (physician or any other medical care provider). Investigators are also requested to try as much as possible to re-contact those patients at the end of the trial to obtain at least their vital status as well as their status with respect to the primary endpoint, and thus avoid lost to follow-up for the efficacy assessment.

If patients are lost to follow-up, the CRF must be completed up to the last visit or contact.

Statistics

Analysis Populations

Randomized Population

The randomized population will include all patients who sign the informed consent form and are assigned a randomization number at Visit 2 (Day 0).

Intent-to-Treat Population

The Intent-to-Treat (ITT) population will consist of all randomized patients who take at least one dose of study drug. The ITT population is the primary analysis population. All efficacy analyses will be performed on the ITT population.

Per-Protocol Population

The per-protocol (PP) population will include all ITT patients without any major protocol deviations, and who had ≥80% compliance with study drug while on treatment (up to discontinuation for patients whose treatment is terminated early). The per-protocol efficacy analysis for CV events will be restricted to each patient's time on study drug plus 30 days thereafter.

Safety Population

All safety analyses will be conducted based on the safety population, which is defined as all randomized patients who receive at least one dose of study drug. This is the same as the ITT population.

Statistical Methods

Safety and efficacy variables will be analyzed using appropriate statistical methods to be described in detail in a separate Statistical Analysis Plan (SAP). The SAP will be finalized before study unblinding.

Patient Disposition and Demographic/Baseline Characteristics

The numbers of patients screened, the number of patients randomized per treatment group (randomized population), and the number of patients in the ITT and PP populations by treatment group will be listed.

For randomized patients who discontinued treatment with study drug, the primary reason for discontinuation will be listed and summarized by treatment group.

Summary statistics (mean, standard deviation, median, minimum and maximum) will be provided by treatment

US 10,568,861 B1

39

group for demographic characteristics (e.g., age, sex, race, and ethnicity) and baseline characteristics (e.g., body weight, height, and body mass index) in the ITT and PP populations.

Demographic data and baseline characteristics will be compared among treatment groups for the ITT and PP population. Differences in demographic and baseline characteristics will be tested using a chi-square test (for categorical variables) or a 1-way analysis of variance model with treatment as a factor (for continuous variables). The p-values will be used as descriptive statistics, primarily as an assessment of the adequacy of randomization.

Study Medication Exposure and Compliance

The final compliance to study drug will be calculated as the percent of doses taken relative to doses scheduled to be taken. Overall percent compliance will be calculated per patient in the ITT and PP populations and summarized by treatment group using summary statistics (n, mean, standard deviation, median, minimum, and maximum).

Concomitant Therapies

Concomitant medication/therapy verbatim terms will be coded using the latest version of the World Health Organization Drug Dictionary. The numbers and percentages of patients in each treatment group taking concomitant medications will be summarized by anatomic and therapeutic chemical classification and preferred term.

Analysis of Efficacy

For efficacy endpoints including CV events, only adjudicated events will be included in the final statistical analyses.

Summary Statistics

Summary statistics (n, mean, standard deviation, median, minimum, and maximum) for the baseline and post-baseline measurements, the percent changes, or changes from baseline will be presented by treatment group and by visit for all efficacy variables to be analyzed. The summary statistics will include changes in body weight and body mass index from baseline by treatment group and by visit.

Primary Endpoint

The primary efficacy endpoint is the time from randomization to the first occurrence of any component of the composite of the following clinical events:

- CV death,
- Nonfatal MI (including silent MI),
- Nonfatal stroke,
- Coronary revascularization,

Hospitalization for unstable angina determined to be caused by myocardial ischemia by invasive/non-invasive testing.

The analysis of the primary efficacy endpoint will be performed using the log-rank test comparing the 2 treatment groups (AMR101 and placebo) and including the stratification factor "CV risk category", use of ezetimibe and geographical region (Westernized, Eastern European, and Asia Pacific countries) (each as recorded in the IWR at the time of enrollment) as covariates. Treatment difference will be tested at alpha level of 0.0476 accounting for one interim efficacy analysis. The hazard ratio for treatment group (AMR101 vs. placebo) from a Cox proportional hazard model that includes the stratification factor will also be reported, along with the associated 95% confidence interval. Kaplan-Meier estimates from randomization to the time to the primary efficacy endpoint will be plotted.

The size and direction of the treatment effects of the individual components of the composite endpoint and their relative contribution to the composite endpoint will be determined as well.

40

Secondary Endpoints

The statistical analyses of the secondary endpoints will be analyzed by the same log-rank test specified above for the primary efficacy endpoint. Treatment differences will be tested at alpha level of 0.05 using a sequential procedure for controlling type 1 error starting with the key secondary variable. The remaining secondary variables will be tested in the order specified in Section 9.2.2. Estimates of the hazard ratios from the Cox proportional hazard model and the associated 95% confidence intervals will also be provided. Kaplan-Meier estimates from randomization the time to the secondary efficacy endpoints will be plotted.

Tertiary Endpoints

For event rates, the statistical analyses of the tertiary endpoints will be similar to the analysis of the secondary efficacy endpoints. All tertiary analyses will be conducted for the ITT population. No adjustments for multiple testing will be made.

For measurements of lipids, lipoproteins and inflammatory markers the change from baseline will be analyzed in the units of each marker, and the percent change from baseline. Since these biomarkers are typically not normally distributed, the Wilcoxon rank-sum test will be used for treatment comparisons of the percent change from baseline, and medians and quartiles will be provided for each treatment group. The medians of the differences between the treatment groups and 95% confidence intervals will be estimated with the Hodges-Lehmann method.

New onset diabetes is defined as Type 2 diabetes newly diagnosed during the treatment/follow-up period (i.e. patients with no history of diabetes at randomization).

For purposes of this study, a diagnosis of diabetes is made based on the observation of:

1. $HbA_{1c} \geq 6.5\%$. The test should be performed in a laboratory using a method that is National Glycohemoglobin Standardization Program (NGSP) certified and standardized to the Diabetes Control and Complications Trial (DCCT) assay. In the absence of unequivocal hyperglycemia, $HbA_{1c} > 6.5\%$ should be confirmed by repeat testing.

OR

2. Fasting plasma glucose (FPG) ≥ 126 mg/dL (7.0 mmol/L). Fasting is defined as no caloric intake for at least 8 hr. In the absence of unequivocal hyperglycemia, FPG ≥ 126 mg/dL (7.0 mmol/L) should be confirmed by repeat testing.

OR

3. 2-hr plasma glucose ≥ 200 mg/dL (11.1 mmol/L) during an Oral Glucose Tolerance Test (OGTT). The test should be performed as described by the World Health Organization, using a glucose load containing the equivalent of 75 g anhydrous glucose dissolved in water. In the absence of unequivocal hyperglycemia, 2-hr plasma glucose ≥ 200 mg/dL (11.1 mmol/L) during an Oral Glucose Tolerance Test (OGTT) should be confirmed by repeat testing.

OR

4. In a patient with classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose ≥ 200 mg/dL (11.1 mmol/L).

Exploratory Subgroup Analyses

Subgroup analyses of the primary and key secondary endpoints (as defined in the Statistical Analysis Plan) will be performed. All subgroup analyses will be conducted for the ITT population. No adjustments for multiple testing will be made.

Log-rank tests, treatment effects and the associated 95% confidence intervals for the primary and key secondary efficacy endpoints within each subgroup will be provided using the Cox proportional hazard model with treatment

US 10,568,861 B1

41

(AMR101 or placebo), and stratification as a factor (with the exception of the subgroup analyses of those subgroup variables related to the stratification factors, i.e., CV risk category that will not have stratification as a factor).

Subgroups including, but not limited to the following, will be explored. A complete list will be prospectively defined in the Statistical Analysis Plan.

Demographics:

Gender,

age (<65 yr and ≥65 yr),

race (white and nonwhite, or any other subset with at least 10% of the total number of patients),

geography (western vs. non-western)

Disease Parameters:

CV risk category,

the presence/absence of diabetes at baseline,

renal impairment

Treatment Parameters:

by statin intensity (statin type and regimen),

relevant concomitant medications,

Baseline Lipid and Lipoprotein Parameters:

LDL-C (by tertile),

HDL-C (by tertile),

TG (by tertile),

TG ≥150 mg/dL,

TG ≥200 mg/dL and TG <200 mg/dL, combined highest tertile for TG and lowest tertile for HDL-C,

hs-CRP (≤3 mg/L and >3 mg/L),

Apo B (by tertile),

non-HDL-C (by tertile)

The consistency of the treatment effects in subgroups will be assessed for the primary and key secondary efficacy endpoints. For each subgroup variable, a Cox proportional hazard model with terms for treatment, stratification factors (with the exception of those subgroup variables related to the stratification factors, i.e., CV risk category), subgroup, and treatment-by-subgroup interaction will be performed. The main treatment effect will not be tested with this model. P-values for testing the interaction terms will be provided.

Interim Efficacy Analysis

One interim analysis will be performed for the primary efficacy endpoint using best available data (adjudicated events and site reported endpoints) based on data when approximately 60% of the total number of primary endpoint events is reached. The interim analysis will be based on a group sequential design that includes early stopping rules for benefit while preserving the overall Type I error rate (O'Brien-Fleming). This allows for interim analysis and preserves the overall Type I error probability of $\alpha=0.05$ for the primary endpoint.

Approximately 1612 primary efficacy endpoint events are planned to be observed during the trial, based on sample size calculation assumptions. Therefore, the interim analysis will occur after at least 967 primary efficacy endpoint events have been observed. According to this boundary, the critical p-value at the interim analysis has to be $p \leq 0.0076$, resulting in the final evaluation p-value of 0.0476.

The interim results of the study will be monitored by an independent DMC. The analyses will be performed by the independent statistical group unblinded to the treatment assignment. The results will be reported only to the DMC. The unblinded information will not be released to sponsor under any circumstance before the completion of the study. Specific statistical guidelines for data monitoring will be discussed and formalized in a separate Interim Statistical Analysis Plan and DMC Charter.

42

Analysis of Safety

All analyses of safety will be conducted on the safety population, which is defined as all randomized patients who receive at least one dose of study drug. The safety assessment will be based on the frequency of adverse events, physical exams, vital signs and safety laboratory tests.

Adverse events with new onset during the study between the initiation of study drug and 30 days after the last dose of study drug for each patient will be considered treatment-emergent (TEAEs). This will include any AE with onset prior to initiation of study drug and increased severity after the treatment initiation.

Treatment-emergent adverse events will be summarized by system organ class and preferred term, and by treatment.

This will include overall incidence rates (regardless of severity and relationship to study drug), and incidence rates for moderate or severe adverse events. A summary of SAEs and adverse events leading to early discontinuation from the study will be presented through data listings.

Safety laboratory tests and vital signs will be summarized by post-treatment change from baseline for each of the parameters using descriptive statistics by treatment group. Those patients with significant laboratory abnormalities will be identified in data listings. Additional safety parameters will be summarized in data listings.

Sample Size Determination

Sample size estimation is based on the assumption that the primary composite endpoint (time from randomization to the first occurrence of CV death, non-fatal MI, non-fatal stroke, coronary revascularization, or unstable angina requiring hospitalization) would be relatively reduced by 15%, from an event rate by 4 years of 23.6% in the placebo group to 20.5% in the AMR101 group. It is expected that a minimum of 1612 primary efficacy endpoint events will be required during the study. A total of approximately 6990 patients are needed to be able to detect this difference at 4.76% significance level (because of the interim analysis described in Section 12.2.4.6) and with 90% power, assuming an 18-month enrollment period and a median follow-up of 4 years. The current sample size calculation is based on an estimated placebo yearly event rate of 5.9% (23.6% over 4 years). To protect against the possibility that the actual placebo event rate is lower than estimated, an extra 1000 patients will be enrolled (approximately 7990 patients in total). By adding the extra 1000 patients, the event rate in the placebo group could be 5.2% per year (20.8% over 4 years) without having to modify the other sample size assumptions.

Since this is an events-driven trial, the 'sample size' is the number of events rather than the number of patients. The number of events that occur depends primarily on three factors: how many patients are enrolled, the combined group event rate, and how long the patients are followed. Because of the difficulty in predicting the combined event rate, the sponsor will monitor that event rate as the trial progresses. If the combined event rate is less than anticipated, either increasing the number of patients, extending the length of follow-up, or a balance of adjusting both factors may be necessary to achieve the sample size of 1612 events.

Before completing the enrollment phase of the trial, i.e. approximately 3- to 6-months prior to the projected enrollment of the 7990th patient, the actual event rate based on pooled, blinded accumulation of primary efficacy endpoint events will be calculated and plotted. If those analyses suggest the number of patients with at least 1 adjudicated, primary event (and appropriately accounting for patients with potential primary events for which the adjudication process is then incomplete) is consistent with projections,

US 10,568,861 B1

43

then the study could continue toward the protocol-specified target enrollment of 7990 patients. However, if the number of such events appears less than, and inconsistent with projections, the Sponsor will consider (under blinded conditions) re-calculating the number of patients needed to achieve the target number of events within the desired timeline or extend the follow-up period. If the projected increase in number of patients is $\leq 25\%$ of the original 7990 target population, the Sponsor may, with documented approval of both the REDUCE-IT Steering Committee (SC) and the Data Monitoring Committee (DMC), extend enrollment to the revised target number without need for an additional protocol amendment. Under those conditions, all principal investigators, ethics committees, and regulatory authorities associated with the protocol will be promptly notified of the action. Should the projected increase in number of patients be more than 25% above the original 7990 target (i.e. more than 1998 additional patients) a formal protocol amendment will be initiated.

If the number of patients to be studied is increased, the enrollment phase will be extended to allow enrollment of the additional patients.

At completion of study enrollment, the actual number of patients randomized may vary from the target number (either original or revised) as a result of the inherent lag between the date the last patient started screening and the date the last patient was randomized.

Monitoring, Data Management, and Record Keeping

Data Management

Data Handling

Data will be recorded at the site on CRFs. All entries on a CRF are ultimately the responsibility of the Investigator, who is expected to review each form for completeness and accuracy before signing. A CRF must be completed for each randomized patient. The CRFs and source documents must be made available to the Sponsor and/or its designee.

Record Keeping

The Investigator must maintain all documents and records, originals or certified copies of original records, relating to the conduct of this trial, and necessary for the evaluation and reconstruction of the clinical trial. This documentation includes, but is not limited to protocol, CRFs, AE reports, patient source data (including records of patients, patient visit logs, clinical observations and findings), correspondence with health authorities and IRB, consent forms, inventory of study product, Investigator's curriculum vitae, monitor visit logs, laboratory reference ranges

44

and laboratory certification or quality control procedures, and laboratory director curriculum vitae.

The Investigator and affiliated institution should maintain the trial documents as required by the applicable regulations. The Investigator and affiliated institution should take measures to prevent accidental or premature destruction of documents. Clinical trial documents must be kept in the clinical site's archives indefinitely, unless written authorization is obtained from the Sponsor.

Direct Access to Source Data/Documents

The investigator and research institution agree that the Sponsor, their representatives and designees, the IRB or IEC, and representatives from worldwide regulatory agencies will have the right, both during and after the clinical trial, to review and inspect pertinent medical records related to the clinical trial.

Quality Control and Quality Assurance

The Sponsor and/or its designee(s) will perform quality control and quality assurance checks of all clinical trials that it sponsors. Before the enrollment of any patient in this study, the Sponsor or its designee will review with the investigator and site personnel the following documents: protocol, Investigator's Brochure, CRFs and procedures for their completion, the informed consent process, and the procedure for reporting SAEs. Site visits will be performed by the Sponsor and/or its designees. During these visits, information recorded on the CRFs will be verified against source documents and requests for clarification or correction may be made. After the CRF data is entered by the site, the Sponsor or designee will review for safety information, completeness, accuracy, and logical consistency. Computer programs that identify data inconsistencies may be used to help monitor the clinical trial. If necessary, requests for clarification or correction will be sent to investigators.

By signing the protocol, the Sponsor agrees directly or through its designee(s) to be responsible for implementing and maintaining quality control and quality assurance systems with written standard operating procedures to ensure that trials are conducted and data are generated, documented, and reported in compliance with the protocol, accepted standards of Good Clinical Practice (GCP), International Conference on Harmonization (ICH) and other applicable regulations.

Completion of Study

The end of the study will be at the time of the last patient-last visit of the follow-up period of the study. The IRB and IEC will be notified about the end of the study according to country-specific regulatory requirements.

TABLE 1

SCHEDULE OF PROCEDURES										
Screening										
Study Day	Up to	If a Visit 1.1 takes place, Visit 1 may occur up to 60 days before Day 0 ²	Follow-Up (FU) ¹³							
	42		0	120	360	720	1080	1440	1800	
	days			±	±	±	±	±	±	+
	before									
	Day 0		0	10	10	10	10	10	10	30
Months of FU			0	4	12	24	36	48	60	
Years of FU			0	0.33	1	2	3	4	5	
Visit #	1	1.1	2	3	4	5	6	7	LV ¹⁴	

US 10,568,861 B1

45

46

TABLE 1-continued

SCHEDULE OF PROCEDURES									
Study Day	Screening		Follow-Up (FU) ¹³						
	Up to 42 days before Day 0	If a Visit 1.1 takes place, Visit 1 may occur up to 60 days before Day 0 ²	0	120 ± 10	360 ± 10	720 ± 10	1080 ± 10	1440 ± 10	1800 ± 30
Study Procedures:									
Informed Consent	X								
Medical, Surgical & Family History	X								
Demographics	X								
Evaluate inclusion/exclusion criteria	X ¹	X ³	X						
Physical Examination			X	X	X	X	X	X	X
Weight, Height ⁴	X		X	X	X	X	X	X	X
Vital Signs ⁵	X	X	X	X	X	X	X	X	X
Waist Circumference			X		X	X	X	X	X
12-Lead ECG	X		X		X	X	X	X	X
Urine pregnancy test ⁶	X		X		X	X	X	X	X
Concomitant Meds	X	X	X	X	X	X	X	X	X
Randomization			X						
Dosing at the Research Site ⁷			X	X	X	X	X	X	
Efficacy events				X	X	X	X	X	X
AE Evaluations			X	X	X	X	X	X	X
Compliance Check ⁸				X	X	X	X	X	X
Chemistry and hematology ⁹	X	X ³	X	X	X	X	X	X	X
Fasting lipid profile ¹⁰	X	X ³	X	X	X	X	X	X	X
Genetic testing ¹¹			X						
Biomarkers: hs-CRP, apo B, hsTNT			X			X			X
Fasting blood sample for archiving ¹²			X		X	X	X	X	X

What is claimed is:

1. A method of reducing risk of cardiovascular death in a subject with established cardiovascular disease, the method comprising administering to said subject about 4 g of ethyl icosapentate per day for a period effective to reduce risk of cardiovascular death in the subject.
2. The method of claim 1, wherein the subject has a fasting baseline triglyceride level of about 135 mg/dL to about 500 mg/dL and a fasting baseline LDL-C level of about 40 mg/dL to about 100 mg/dL.
3. The method of claim 1, wherein the ethyl icosapentate is present in a pharmaceutical composition and the ethyl

icosapentate comprises at least about 96 wt. % of all omega-3 fatty acids in the pharmaceutical composition.

4. The method of claim 3, wherein about 1 g of the pharmaceutical composition is present in each of 4 capsules.

5. The method of claim 1, wherein said period ends at least 2 years after initial administration of the ethyl icosapentate to the subject.

6. The method of claim 1, wherein the subject is on statin therapy.

7. The method of claim 1, wherein the subject has a triglyceride level of at least 135 mg/dL and is on statin therapy.

* * * * *

FORM 19. Certificate of Compliance with Type-Volume Limitations

Form 19
July 2020

**UNITED STATES COURT OF APPEALS
FOR THE FEDERAL CIRCUIT**

CERTIFICATE OF COMPLIANCE WITH TYPE-VOLUME LIMITATIONS

Case Number: 23-1169

Short Case Caption: Amarin Pharma, Inc. v. Hikma Pharmaceuticals USA Inc.

Instructions: When computing a word, line, or page count, you may exclude any items listed as exempted under Fed. R. App. P. 5(c), Fed. R. App. P. 21(d), Fed. R. App. P. 27(d)(2), Fed. R. App. P. 32(f), or Fed. Cir. R. 32(b)(2).

The foregoing filing complies with the relevant type-volume limitation of the Federal Rules of Appellate Procedure and Federal Circuit Rules because it meets one of the following:

- ☒ the filing has been prepared using a proportionally-spaced typeface and includes 11,888 words.
- ☐ the filing has been prepared using a monospaced typeface and includes _____ lines of text.
- ☐ the filing contains _____ pages / _____ words / _____ lines of text, which does not exceed the maximum authorized by this court's order (ECF No. _____).

Date: 03/21/2023

Signature: /s/Nathan K. Kelley

Name: Nathan K. Kelley